

# Mario Fichera

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6993392/publications.pdf>

Version: 2024-02-01

16  
papers

295  
citations

1040056

9  
h-index

1058476

14  
g-index

16  
all docs

16  
docs citations

16  
times ranked

619  
citing authors

#	ARTICLE	IF	CITATIONS
1	Cognitive Reserve in Early Manifest Huntington Disease Patients: Leisure Time Is Associated with Lower Cognitive and Functional Impairment. <i>Journal of Personalized Medicine</i> , 2022, 12, 36.	2.5	7
2	Hsa-miR223-3p circulating level is upregulated in Friedreich's ataxia and inversely associated with HCLS1 associated protein X-1, HAX-1. <i>Human Molecular Genetics</i> , 2022, , .	2.9	1
3	Neuro-telehealth for fragile patients in a tertiary referral neurological institute during the COVID-19 pandemic in Milan, Lombardy. <i>Neurological Sciences</i> , 2021, 42, 2637-2644.	1.9	18
4	Progression characteristics of the European Friedreich's Ataxia Consortium for Translational Studies (EFACTS): a 4-year cohort study. <i>Lancet Neurology</i> , The, 2021, 20, 362-372.	10.2	53
5	Frataxin deficiency in Friedreich's ataxia is associated with reduced levels of HAX-1, a regulator of cardiomyocyte death and survival. <i>Human Molecular Genetics</i> , 2020, 29, 471-482.	2.9	8
6	Bilateral Repetitive Transcranial Magnetic Stimulation With the H-Coil in Parkinson's Disease: A Randomized, Sham-Controlled Study. <i>Frontiers in Neurology</i> , 2020, 11, 584713.	2.4	13
7	NfL and pNfH are increased in Friedreich's ataxia. <i>Journal of Neurology</i> , 2020, 267, 1420-1430.	3.6	17
8	Protocol of a randomized, double-blind, placebo-controlled, parallel-group, multicentre study of the efficacy and safety of nicotinamide in patients with Friedreich ataxia (NICOFA). <i>Neurological Research and Practice</i> , 2019, 1, 33.	2.0	14
9	Bi-hemispheric repetitive transcranial magnetic stimulation for upper limb motor recovery in chronic stroke: A feasibility study. <i>Brain Stimulation</i> , 2018, 11, 932-934.	1.6	4
10	Which patients discontinue? Issues on Levodopa/carbidopa intestinal gel treatment: Italian multicentre survey of 905 patients with long-term follow-up. <i>Parkinsonism and Related Disorders</i> , 2017, 38, 90-92.	2.2	44
11	Weight gain after subthalamic nucleus deep brain stimulation in Parkinson's disease is influenced by dyskinesias reduction and electrodes position. <i>Neurological Sciences</i> , 2017, 38, 2123-2129.	1.9	19
12	Drawing in Alzheimer-type dementia. <i>Neurology</i> , 2015, 85, 386-387.	1.1	0
13	Excitatory Deep Repetitive Transcranial Magnetic Stimulation With H-coil as Add-on Treatment of Motor Symptoms in Parkinson's Disease: An Open Label, Pilot Study. <i>Brain Stimulation</i> , 2014, 7, 297-300.	1.6	38
14	Secondary cervical dystonic tremor after Japanese encephalitis. <i>Neurological Sciences</i> , 2014, 35, 491-493.	1.9	3
15	Deep magnetic stimulation in a progressive supranuclear palsy patient with speech involvement. <i>Journal of Neurology</i> , 2013, 260, 670-673.	3.6	10
16	Interhemispheric Balance in Parkinson's Disease: A Transcranial Magnetic Stimulation Study. <i>Brain Stimulation</i> , 2013, 6, 892-897.	1.6	46