

Ursula Zimmer-Strobl

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7068611/publications.pdf>

Version: 2024-02-01

52
papers

3,432
citations

136950

32
h-index

175258

52
g-index

56
all docs

56
docs citations

56
times ranked

5727
citing authors

#	ARTICLE	IF	CITATIONS
1	PARP14 is a novel target in STAT6 mutant follicular lymphoma. <i>Leukemia</i> , 2022, 36, 2281-2292.	7.2	11
2	Notch2-mediated plasticity between marginal zone and follicular B cells. <i>Nature Communications</i> , 2021, 12, 1111.	12.8	26
3	ERK phosphorylation is RAF independent in naïve and activated B cells but RAF dependent in plasma cell differentiation. <i>Science Signaling</i> , 2021, 14, .	3.6	7
4	Context-dependent regulation of immunoglobulin mutagenesis by p53. <i>Molecular Immunology</i> , 2021, 138, 128-136.	2.2	1
5	Pre-clinical blocking of PD-L1 molecule, which expression is down regulated by NF- κ B, JAK1/JAK2 and BTK inhibitors, induces regression of activated B-cell lymphoma. <i>Cell Communication and Signaling</i> , 2019, 17, 89.	6.5	19
6	Chronic CD30 signaling in B cells results in lymphomagenesis by driving the expansion of plasmablasts and B1 cells. <i>Blood</i> , 2019, 133, 2597-2609.	1.4	14
7	PiggyBac transposon tools for recessive screening identify B-cell lymphoma drivers in mice. <i>Nature Communications</i> , 2019, 10, 1415.	12.8	37
8	Reproducing indolent B-cell lymphoma transformation with T-cell immunosuppression in LMP1/CD40-expressing mice. <i>Cellular and Molecular Immunology</i> , 2019, 16, 412-414.	10.5	6
9	Notch2 Signaling Maintains NSC Quiescence in the Murine Ventricular-Subventricular Zone. <i>Cell Reports</i> , 2018, 22, 992-1002.	6.4	93
10	Notch2 controls non-autonomous Wnt-signalling in chronic lymphocytic leukaemia. <i>Nature Communications</i> , 2018, 9, 3839.	12.8	51
11	Jagged1/Notch2 controls kidney fibrosis via Tfam-mediated metabolic reprogramming. <i>PLoS Biology</i> , 2018, 16, e2005233.	5.6	51
12	LAG-3 Inhibitory Receptor Expression Identifies Immunosuppressive Natural Regulatory Plasma Cells. <i>Immunity</i> , 2018, 49, 120-133.e9.	14.3	190
13	CD40-signalling abrogates induction of ROR γ t+ Treg cells by intestinal CD103+ DCs and causes fatal colitis. <i>Nature Communications</i> , 2017, 8, 14715.	12.8	36
14	c-Myc dysregulation is a co-transforming event for nuclear factor- κ B activated B cells. <i>Haematologica</i> , 2017, 102, 883-894.	3.5	17
15	HSP90 inhibition overcomes ibrutinib resistance in mantle cell lymphoma. <i>Blood</i> , 2016, 128, 2517-2526.	1.4	37
16	Regulation of monocyte cell fate by blood vessels mediated by Notch signalling. <i>Nature Communications</i> , 2016, 7, 12597.	12.8	115
17	Checkpoint kinase 1 negatively regulates somatic hypermutation. <i>Nucleic Acids Research</i> , 2014, 42, 3666-3674.	14.5	12
18	B-cell Expansion and Lymphomagenesis Induced by Chronic CD40 Signaling Is Strictly Dependent on CD19. <i>Cancer Research</i> , 2014, 74, 4318-4328.	0.9	13

#	ARTICLE	IF	CITATIONS
19	Canonical Notch2 signaling determines biliary cell fates of embryonic hepatoblasts and adult hepatocytes independent of Hes1. <i>Hepatology</i> , 2013, 57, 2469-2479.	7.3	85
20	An RNAi-Based Approach to Down-Regulate a Gene Family In Vivo. <i>PLoS ONE</i> , 2013, 8, e80312.	2.5	2
21	Humanized c-Myc Mouse. <i>PLoS ONE</i> , 2012, 7, e42021.	2.5	4
22	Immune modulation by Fas ligand reverse signaling: lymphocyte proliferation is attenuated by the intracellular Fas ligand domain. <i>Blood</i> , 2011, 117, 519-529.	1.4	26
23	Programming of marginal zone B-cell fate by basic KrÄppel-like factor (BKLK/KLF3). <i>Blood</i> , 2011, 117, 3780-3792.	1.4	26
24	CD19-independent instruction of murine marginal zone B-cell development by constitutive Notch2 signaling. <i>Blood</i> , 2011, 118, 6321-6331.	1.4	69
25	Asymmetric Arginine dimethylation of Epstein-Barr virus nuclear antigen 2 promotes DNA targeting. <i>Virology</i> , 2010, 397, 299-310.	2.4	16
26	Identification of Epidermal Pdx1 Expression Discloses Different Roles of Notch1 and Notch2 in Murine KrasG12D-Induced Skin Carcinogenesis In Vivo. <i>PLoS ONE</i> , 2010, 5, e13578.	2.5	36
27	The fusion kinase ITK-SYK mimics a T cell receptor signal and drives oncogenesis in conditional mouse models of peripheral T cell lymphoma. <i>Journal of Experimental Medicine</i> , 2010, 207, 1031-1044.	8.5	134
28	Notch2 is required for progression of pancreatic intraepithelial neoplasia and development of pancreatic ductal adenocarcinoma. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2010, 107, 13438-13443.	7.1	190
29	The fusion kinase ITK-SYK mimics a T cell receptor signal and drives oncogenesis in conditional mouse models of peripheral T cell lymphoma. <i>Journal of Cell Biology</i> , 2010, 189, i10-i10.	5.2	0
30	Notch1, Notch2, and Epstein-Barr virus-encoded nuclear antigen 2 signaling differentially affects proliferation and survival of Epstein-Barr virus-infected B cells. <i>Blood</i> , 2009, 113, 5506-5515.	1.4	31
31	Liver-specific inactivation of <i>Notch2</i> , but not <i>Notch1</i> , compromises intrahepatic bile duct development in mice. <i>Hepatology</i> , 2008, 48, 607-616.	7.3	194
32	Loss of intestinal crypt progenitor cells owing to inactivation of both Notch1 and Notch2 is accompanied by derepression of CDK inhibitors p27 ^{Kip1} and p57 ^{Kip2} . <i>EMBO Reports</i> , 2008, 9, 377-383.	4.5	362
33	Molecular Basis of Cytotoxicity of Epstein-Barr Virus (EBV) Latent Membrane Protein 1 (LMP1) in EBV Latency III B Cells: LMP1 Induces Type II Ligand-Independent Autoactivation of CD95/Fas with Caspase 8-Mediated Apoptosis. <i>Journal of Virology</i> , 2008, 82, 6721-6733.	3.4	49
34	Conditional ablation of Notch signaling in pancreatic development. <i>Development (Cambridge)</i> , 2008, 135, 2757-2765.	2.5	75
35	Constitutive CD40 signaling in B cells selectively activates the noncanonical NF-ÎB pathway and promotes lymphomagenesis. <i>Journal of Experimental Medicine</i> , 2008, 205, 1317-1329.	8.5	117
36	LMP1 signaling can replace CD40 signaling in B cells in vivo and has unique features of inducing class-switch recombination to IgG1. <i>Blood</i> , 2008, 111, 1448-1455.	1.4	96

#	ARTICLE	IF	CITATIONS
37	Restricted Expression of Epstein-Barr Virus Latent Genes in Murine B Cells Derived from Embryonic Stem Cells. PLoS ONE, 2008, 3, e1996.	2.5	2
38	Hierarchy of Notch-Delta interactions promoting T cell lineage commitment and maturation. Journal of Experimental Medicine, 2007, 204, 331-343.	8.5	161
39	B- and T-cell-specific inactivation of thioredoxin reductase 2 does not impair lymphocyte development and maintenance. Biological Chemistry, 2007, 388, 1083-1090.	2.5	16
40	Notch1 and Notch2 receptors influence progressive hair graying in a dose-dependent manner. Developmental Dynamics, 2007, 236, 282-289.	1.8	115
41	Control of Epstein-Barr virus reactivation by activated CD40 and viral latent membrane protein 1. Proceedings of the National Academy of Sciences of the United States of America, 2002, 99, 437-442.	7.1	75
42	Magnetic DNA Affinity Purification of a Cellular Transcription Factor. , 2001, 174, 271-277.		1
43	EBNA2 and Notch signalling in Epstein-Barr virus mediated immortalization of B lymphocytes. Seminars in Cancer Biology, 2001, 11, 423-434.	9.6	119
44	Activated Notch1 Can Transiently Substitute for EBNA2 in the Maintenance of Proliferation of LMP1-Expressing Immortalized B Cells. Journal of Virology, 2001, 75, 2033-2040.	3.4	64
45	Activated Notch1 Modulates Gene Expression in B Cells Similarly to Epstein-Barr Viral Nuclear Antigen 2. Journal of Virology, 2000, 74, 1727-1735.	3.4	86
46	Activated Mouse Notch1 Transactivates Epstein-Barr Virus Nuclear Antigen 2-Regulated Viral Promoters. Journal of Virology, 1999, 73, 2770-2780.	3.4	44
47	Functional Replacement of the Intracellular Region of the Notch1 Receptor by Epstein-Barr Virus Nuclear Antigen 2. Journal of Virology, 1998, 72, 6034-6039.	3.4	67
48	RBP-L, a Transcription Factor Related to RBP-J δ . Molecular and Cellular Biology, 1997, 17, 2679-2687.	2.3	122
49	Both Epstein-Barr Viral Nuclear Antigen 2 (EBNA2) and Activated Notch1 Transactivate Genes by Interacting with the Cellular Protein RBP-J δ . Immunobiology, 1997, 198, 299-306.	1.9	84
50	Epstein-Barr Virus Nuclear Antigen 2-Estrogen Receptor Fusion Proteins Transactivate Viral and Cellular Genes and Interact with RBP-J δ in a Conditional Fashion. Virology, 1995, 214, 675-679.	2.4	44
51	Crucial sequences within the Epstein-Barr virus TP1 promoter for EBNA2-mediated transactivation and interaction of EBNA2 with its responsive element. Journal of Virology, 1994, 68, 7497-7506.	3.4	51
52	Epstein-Barr virus nuclear antigen 2 exerts its transactivating function through interaction with recombination signal binding protein RBP-J kappa, the homologue of Drosophila Suppressor of Hairless. EMBO Journal, 1994, 13, 4973-82.	7.8	129