
## David A Simpson

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/7070224/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Ibrutinib as Initial Therapy for Patients with Chronic Lymphocytic Leukemia. New England Journal of Medicine, 2015, 373, 2425-2437.	27.0	1,261
2	Rituximab maintenance for 2 years in patients with high tumour burden follicular lymphoma responding to rituximab plus chemotherapy (PRIMA): a phase 3, randomised controlled trial. Lancet, The, 2011, 377, 42-51.	13.7	957
3	International, evidence-based consensus diagnostic criteria for HHV-8–negative/idiopathic multicentric Castleman disease. Blood, 2017, 129, 1646-1657.	1.4	381
4	Siltuximab for multicentric Castleman's disease: a randomised, double-blind, placebo-controlled trial. Lancet Oncology, The, 2014, 15, 966-974.	10.7	345
5	Long-term efficacy and safety of first-line ibrutinib treatment for patients with CLL/SLL: 5 years of follow-up from the phase 3 RESONATE-2 study. Leukemia, 2020, 34, 787-798.	7.2	321
6	Phase 1 study of the selective BTK inhibitor zanubrutinib in B-cell malignancies and safety and efficacy evaluation in CLL. Blood, 2019, 134, 851-859.	1.4	259
7	International, evidence-based consensus treatment guidelines for idiopathic multicentric Castleman disease. Blood, 2018, 132, 2115-2124.	1.4	232
8	Idelalisib or placebo in combination with bendamustine and rituximab in patients with relapsed or refractory chronic lymphocytic leukaemia: interim results from a phase 3, randomised, double-blind, placebo-controlled trial. Lancet Oncology, The, 2017, 18, 297-311.	10.7	219
9	Ponatinib versus imatinib for newly diagnosed chronic myeloid leukaemia: an international, randomised, open-label, phase 3 trial. Lancet Oncology, The, 2016, 17, 612-621.	10.7	214
10	First-Line Treatment of Patients With Indolent Non-Hodgkin Lymphoma or Mantle-Cell Lymphoma With Bendamustine Plus Rituximab Versus R-CHOP or R-CVP: Results of the BRIGHT 5-Year Follow-Up Study. Journal of Clinical Oncology, 2019, 37, 984-991.	1.6	183
11	Pembrolizumab plus pomalidomide and dexamethasone for patients with relapsed or refractory multiple myeloma (KEYNOTE-183): a randomised, open-label, phase 3 trial. Lancet Haematology,the, 2019, 6, e459-e469.	4.6	174
12	Sustained Progression-Free Survival Benefit of Rituximab Maintenance in Patients With Follicular Lymphoma: Long-Term Results of the PRIMA Study. Journal of Clinical Oncology, 2019, 37, 2815-2824.	1.6	173
13	Detection of atomic spin labels in a lipid bilayer using a single-spin nanodiamond probe. Proceedings of the National Academy of Sciences of the United States of America, 2013, 110, 10894-10898.	7.1	113
14	Non-Neurotoxic Nanodiamond Probes for Intraneuronal Temperature Mapping. ACS Nano, 2017, 11, 12077-12086.	14.6	113
15	Sustained efficacy and detailed clinical follow-up of first-line ibrutinib treatment in older patients with chronic lymphocytic leukemia: extended phase 3 results from RESONATE-2. Haematologica, 2018, 103, 1502-1510.	3.5	111
16	Scanning Nanospin Ensemble Microscope for Nanoscale Magnetic and Thermal Imaging. Nano Letters, 2016, 16, 326-333.	9.1	79
17	Zanubrutinib for the treatment of patients with Waldenström macroglobulinemia: 3 years of follow-up. Blood, 2020, 136, 2027-2037.	1.4	78
18	Ibrutinib for the treatment of Bing-Neel syndrome: a multicenter study. Blood, 2019, 133, 299-305.	1.4	69

DAVID A SIMPSON

#	Article	IF	CITATIONS
19	Updated 6 Year Follow-Up Of The PRIMA Study Confirms The Benefit Of 2-Year Rituximab Maintenance In Follicular Lymphoma Patients Responding To Frontline Immunochemotherapy. Blood, 2013, 122, 509-509.	1.4	66
20	Epidemiology of Castleman Disease. Hematology/Oncology Clinics of North America, 2018, 32, 1-10.	2.2	62
21	Zanubrutinib monotherapy for patients with treatment-naÃ <sup>-</sup> ve chronic lymphocytic leukemia and 17p deletion. Haematologica, 2021, 106, 2354-2363.	3.5	62
22	Zanubrutinib for the treatment of relapsed or refractory mantle cell lymphoma. Blood Advances, 2021, 5, 2577-2585.	5.2	60
23	Not All Fluorescent Nanodiamonds Are Created Equal: A Comparative Study. Particle and Particle Systems Characterization, 2019, 36, 1900009.	2.3	56
24	Electrospun Nanodiamond–Silk Fibroin Membranes: A Multifunctional Platform for Biosensing and Wound-Healing Applications. ACS Applied Materials & Interfaces, 2020, 12, 48408-48419.	8.0	50
25	Magnetically sensitive nanodiamond-doped tellurite glass fibers. Scientific Reports, 2018, 8, 1268.	3.3	44
26	Impact of Surface Functionalization on the Quantum Coherence of Nitrogen-Vacancy Centers in Nanodiamonds. ACS Applied Materials & Interfaces, 2018, 10, 13143-13149.	8.0	36
27	Updated Efficacy and Safety from the Phase 3 Resonate-2 Study: Ibrutinib As First-Line Treatment Option in Patients 65 Years and Older with Chronic Lymphocytic Leukemia/Small Lymphocytic Leukemia. Blood, 2016, 128, 234-234.	1.4	36
28	Enhanced Widefield Quantum Sensing with Nitrogen-Vacancy Ensembles Using Diamond Nanopillar Arrays. ACS Applied Materials & Interfaces, 2020, 12, 13421-13427.	8.0	33
29	Lenalidomide maintenance therapy in previously treated chronic lymphocytic leukaemia (CONTINUUM): a randomised, double-blind, placebo-controlled, phase 3 trial. Lancet Haematology,the, 2017, 4, e534-e543.	4.6	31
30	T-Cell Depleting Antibodies. BioDrugs, 2003, 17, 147-154.	4.6	30
31	Epic: A Phase 3 Trial of Ponatinib Compared with Imatinib in Patients with Newly Diagnosed Chronic Myeloid Leukemia in Chronic Phase (CP-CML). Blood, 2014, 124, 519-519.	1.4	30
32	Subgroup analysis of ICARIAâ€MM study in relapsed/refractory multiple myeloma patients with highâ€risk cytogenetics. British Journal of Haematology, 2021, 194, 120-131.	2.5	27
33	Singleâ€agent ibrutinib versus chemoimmunotherapy regimens for treatmentâ€naÃ⁻ve patients with chronic lymphocytic leukemia: A crossâ€trial comparison of phase 3 studies. American Journal of Hematology, 2018, 93, 1402-1410.	4.1	24
34	Zanubrutinib for treatmentâ€naÃ⁻ve and relapsed/refractory chronic lymphocytic leukaemia: longâ€ŧerm followâ€up of the phase I/II AUâ€003 study. British Journal of Haematology, 2022, 196, 1209-1218.	2.5	24
35	A practical guide to laboratory investigations at diagnosis and follow up in WaldenstrA¶m macroglobulinaemia: recommendations from the Medical and Scientific Advisory Group, Myeloma Australia, the Pathology Sub-committee of the Lymphoma and Related Diseases Registry and the Australasian Association of Clinical Biochemists Monoclonal Gammopathy Working Group.	0.6	23
36	Pathology, 2020, 52, 167-178. Efficacy and Safety of Zanubrutinib in Patients with Treatment-Naive Chronic Lymphocytic Leukemia (CLL) or Small Lymphocytic Lymphoma (SLL) with Del(17p): Initial Results from Arm C of the Sequoia (BGB-3111-304) Trial. Blood, 2019, 134, 499-499.	1.4	23

DAVID A SIMPSON

#	Article	IF	CITATIONS
37	Twice Daily Dosing with the Highly Specific BTK Inhibitor, Bgb-3111, Achieves Complete and Continuous BTK Occupancy in Lymph Nodes, and Is Associated with Durable Responses in Patients (pts) with Chronic Lymphocytic Leukemia (CLL)/Small Lymphocytic Lymphoma (SLL). Blood, 2016, 128, 642-642.	1.4	23
38	Zanubrutinib monotherapy in relapsed/refractory mantle cell lymphoma: a pooled analysis of two clinical trials. Journal of Hematology and Oncology, 2021, 14, 167.	17.0	21
39	New developments in the prophylaxis and treatment of graft versus host disease. Expert Opinion on Pharmacotherapy, 2001, 2, 1109-1117.	1.8	20
40	The effect of tyrosine kinase inhibitor interruption and interferon use on pregnancy outcomes and long-term disease control in chronic myeloid leukemia. Leukemia and Lymphoma, 2019, 60, 1796-1802.	1.3	20
41	An Open-Label, Randomized Study of Bendamustine and Rituximab (BR) Compared with Rituximab, Cyclophosphamide, Vincristine, and Prednisone (R-CVP) or Rituximab, Cyclophosphamide, Doxorubicin, Vincristine, and Prednisone (R-CHOP) in First-Line Treatment of Patients with Advanced Indolent Non-Hodgkin's Lymphoma (NHL) or Mantle Cell Lymphoma (MCL): The Bright Study. Blood, 2012, 120,	1.4	19
42	902:902 Treatment with the Bruton Tyrosine Kinase Inhibitor Zanubrutinib (BGB-3111) Demonstrates High Overall Response Rate and Durable Responses in Patients with Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma (CLL/SLL): Updated Results from a Phase 1/2 Trial. Blood, 2019, 134, 500-500.	1.4	18
43	First-line treatment of iNHL or MCL patients with BR or R-CHOP/R-CVP: Results of the BRIGHT 5-year follow-up study Journal of Clinical Oncology, 2017, 35, 7500-7500.	1.6	18
44	Differences in Quality of Life Between Bendamustine-Rituximab and R-CHOP/R-CVP in Patients With Previously Untreated Advanced Indolent Non-Hodgkin Lymphoma or Mantle Cell Lymphoma. Clinical Lymphoma, Myeloma and Leukemia, 2016, 16, 182-190.e1.	0.4	15
45	Optimizing Optical Tweezers Experiments for Magnetic Resonance Sensing with Nanodiamonds. ACS Photonics, 2021, 8, 1214-1221.	6.6	13
46	A Randomised Dose De-Escalation Study of Oral Fludarabine, ±Oral Cyclophosphamide and Intravenous Rituximab As First-Line Therapy of Fit Patients with Chronic Lymphocytic Leukaemia (CLL) Aged ≥65 Years: Final Analysis of Response and Toxicity. Blood, 2014, 124, 3325-3325.	1.4	13
47	Results of the Phase 3 Study of Lenalidomide Versus Placebo As Maintenance Therapy Following Second-Line Treatment for Patients with Chronic Lymphocytic Leukemia (the CONTINUUM Trial). Blood, 2016, 128, 230-230.	1.4	11
48	High Major Response Rate, Including Very Good Partial Responses (VGPR), in Patients (pts) with Waldenstrom Macroglobulinemia (WM) Treated with the Highly Specific BTK Inhibitor Bgb-3111: Expansion Phase Results from an Ongoing Phase I Study. Blood, 2016, 128, 1216-1216.	1.4	9
49	A Multicenter, Randomized, Double-Blind, Placebo-Controlled Study Of The Efficacy and Safety Of Siltuximab, An Anti-Interleukin-6 Monoclonal Antibody, In Patients With Multicentric Castleman's Disease. Blood, 2013, 122, 505-505.	1.4	7
50	Tresperimus: a new agent for transplant tolerance induction. Expert Opinion on Investigational Drugs, 2001, 10, 1381-1386.	4.1	6
51	Updated Safety and Activity of the Investigational Bruton Tyrosine Kinase Inhibitor Zanubrutinib (BGB-3111) in Patients with Mantle Cell Lymphoma. Blood, 2018, 132, 1592-1592.	1.4	6
52	A Randomised Dose De-Escalation Safety Study of Oral Fludarabine, ±Oral Cyclophosphamide and Intravenous Rituximab (OFOCIR) As First-Line Therapy of Fit Patients with Chronic Lymphocytic Leukaemia (CLL) Aged ≥65 Years – End of Recruitment Analysis of Response and Toxicity of the Australasian Leukaemia and Lymphoma Group (ALLG) and CLL Australian Research Consortium (CLLARC)	1.4	6
53	CLL5 Study. Blood, 2012, 120, 436-436. A cross-trial comparison of single-agent ibrutinib versus chlorambucil-obinutuzumab in previously untreated patients with chronic lymphocytic leukemia or small lymphocytic lymphoma. Haematologica, 2020, 105, e164-e168.	3.5	5
54	Differences in Quality of Life Between Bendamustine Plus Rituximab Compared with Standard First-Line Treatments in Patients with Previously Untreated Advanced Indolent Non-Hodgkin's Lymphoma or Mantle Cell Lymphoma. Blood, 2012, 120, 155-155.	1.4	5

DAVID A SIMPSON

#	Article	IF	CITATIONS
55	Three-year follow-up of treatment-naÃ⁻ve and previously treated patients with Waldenström macroglobulinemia (WM) receiving single-agent zanubrutinib Journal of Clinical Oncology, 2020, 38, 8051-8051.	1.6	5
56	Updated Analysis of Overall Survival in Randomized Phase III Study of Idelalisib in Combination with Bendamustine and Rituximab in Patients with Relapsed/Refractory CLL. Blood, 2016, 128, 231-231.	1.4	4
57	Secondary efficacy subanalysis by histology from the phase III BRIGHT study: First-line bendamustine-rituximab (BR) compared with standard R-CHOP/R-CVP for patients with advanced indolent non-Hodgkin lymphoma (NHL) or mantle cell lymphoma (MCL) Journal of Clinical Oncology, 2013, 31, 8537-8537.	1.6	4
58	Outcomes of First-Line Ibrutinib in Patients with Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma (CLL/SLL) and High-Risk Genomic Features with up to 6.5 Years Follow-up: Integrated Analysis of Two Phase 3 Studies (RESONATE-2 and iLLUMINATE). Blood, 2020, 136, 25-26.	1.4	4
59	Single-Agent Ibrutinib Versus Chlorambucil-Obinutuzumab As First-Line Treatment in Patients with Chronic Lymphocytic Leukemia or Small Lymphocytic Lymphoma (CLL/SLL): Results of a Cross-Trial Comparison. Blood, 2018, 132, 5565-5565.	1.4	3
60	Updated Results of the PRIMA Study Confirms the Benefit of 2-Years Rituximab Maintenance In Follicular Lymphoma Patients Responding to Immunochemotherapy Blood, 2010, 116, 1788-1788.	1.4	3
61	Teniposide, Cyclophosphamide, and Prednisone in Myeloma after Bortezomib and Imid Failure Is a Useful Option. Blood, 2014, 124, 5767-5767.	1.4	3
62	EPIC: A phase III trial of ponatinib (PON) versus imatinib (IM) in patients (pts) with newly diagnosed CP-CML Journal of Clinical Oncology, 2014, 32, 7023-7023.	1.6	3
63	Once Weekly Subcutanous Bortezomib with Cyclophosphamide and Dexamethasone Is Well Tolerated and Effective As Initial Treatment in Symptomatic Multiple Myeloma. Blood, 2012, 120, 4049-4049.	1.4	3
64	Single Centre Experience in Treating Elderly Patients with Acute Myeloid Leukaemia or High Risk Myelodysplastic Syndrome with a Combination of Low Dose Cytarabine, Venetoclax and Fluconazole (VeLDAC-F). Blood, 2018, 132, 5220-5220.	1.4	2
65	Polynesian Variant Of Idiopathic Multicentric Castleman Disease. Blood, 2013, 122, 5127-5127.	1.4	2
66	Results from the International, Randomized Phase 3 Study of Ibrutinib Versus Chlorambucil in Patients 65 Years and Older with Treatment-NaÃ-ve CLL/SLL (RESONATE-2TM). Blood, 2015, 126, 495-495.	1.4	2
67	Efficacy of siltuximab in patients with previously treated multicentric Castleman's disease (MCD) Journal of Clinical Oncology, 2014, 32, 8514-8514.	1.6	2
68	Ibrutinib for the Treatment of Bing-Neel Syndrome. Blood, 2018, 132, 1609-1609.	1.4	2
69	Bing Neel Syndrome: Retrospective Australasian Experience of a Rare Treatable Complication of WaldenstrA¶m Macroglobulinaemia/ Lymphoplasmacytic Lymphoma. Blood, 2018, 132, 5325-5325.	1.4	1
70	Different safety profiles of first-line bendamustine-rituximab (BR), R-CHOP, and R-CVP in an open-label, randomized study of indolent non-Hodgkin lymphoma (NHL) and mantle cell lymphoma (MCL): The BRIGHT study Journal of Clinical Oncology, 2013, 31, 8565-8565.	1.6	1
71	Phase 3 study of oral lenalidomide as maintenance therapy for patients with B-cell chronic lymphocytic leukemia (CLL) Journal of Clinical Oncology, 2014, 32, TPS7125-TPS7125.	1.6	1
72	Fixed Dose Carfilzomib Triplets in Lenalidomide Refractory Myeloma. Blood, 2018, 132, 3306-3306.	1.4	1

#	Article	IF	CITATIONS
73	Thalidomide: An Active Agent in AILT Lymphoma. Blood, 2008, 112, 4960-4960.	1.4	Ο
74	Renal Amyloidosis - Tempo of Proteinuria Resolution in Bortezomib Era. Blood, 2014, 124, 5651-5651.	1.4	0
75	Durable Responses in Fit Elderly Patients with Chronic Lymphocytic Leukemia (CLL) in a Randomised, Fludarabine-Based, Immunochemotherapy Dose De-Escalation Study - Long-Term Follow-up By Treatment Arm and Mutational Status. Blood, 2018, 132, 4432-4432.	1.4	0