

Tomasz Rakowski

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/707467/publications.pdf>

Version: 2024-02-01

154
papers

2,146
citations

304743

22
h-index

315739

38
g-index

159
all docs

159
docs citations

159
times ranked

2519
citing authors

#	ARTICLE	IF	CITATIONS
1	Early glycoprotein IIb/IIIa inhibitors in primary angioplasty (EGYPT) cooperation: an individual patient data meta-analysis. <i>Heart</i> , 2008, 94, 1548-1558.	2.9	135
2	Early glycoprotein IIb/IIIa inhibitors in primary angioplastyâ€”abciximab longâ€”term results (EGYPTâ€”ALT) cooperation: individual patientâ€™s data metaâ€”analysis. <i>Journal of Thrombosis and Haemostasis</i> , 2011, 9, 2361-2370.	3.8	115
3	Impact of Multivessel Coronary Artery Disease and Noninfarct-Related Artery Revascularization on Outcome of Patients With ST-Elevation Myocardial Infarction Transferred for Primary Percutaneous Coronary Intervention (from the EUROTRANSFER Registry). <i>American Journal of Cardiology</i> , 2010, 106, 342-347.	1.6	109
4	Elevated plasma asymmetric dimethyl-L-arginine levels are linked to endothelial progenitor cell depletion and carotid atherosclerosis in rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , 2007, 56, 809-819.	6.7	105
5	Diabetes mellitus is associated with distal embolization, impaired myocardial perfusion, and higher mortality in patients with ST-segment elevation myocardial infarction treated with primary angioplasty and glycoprotein IIb/IIIa inhibitors. <i>Atherosclerosis</i> , 2009, 207, 181-185.	0.8	85
6	Thrombus aspiration followed by direct stenting: A novel strategy of primary percutaneous coronary intervention in ST-segment elevation myocardial infarction. Results of the Polish-Italian-Hungarian RANdomized ThrombEctomy Trial (PIHRATE Trial). <i>American Heart Journal</i> , 2010, 160, 966-972.	2.7	83
7	European registry on patients with ST-elevation myocardial infarction transferred for mechanical reperfusion with a special focus on early administration of abciximabâ€”EUROTRANSFER Registry. <i>American Heart Journal</i> , 2008, 156, 1147-1154.	2.7	60
8	Early abciximab administration before primary percutaneous coronary intervention improves infarct-related artery patency and left ventricular function in high-risk patients with anterior wall myocardial infarction: A randomized study. <i>American Heart Journal</i> , 2007, 153, 360-365.	2.7	52
9	Impact of infarct-related artery patency before primary PCI on outcome in patients with ST-segment elevation myocardial infarction: the HORIZONS-AMI trial. <i>EuroIntervention</i> , 2013, 8, 1307-1314.	3.2	42
10	Gender-related differences in outcome after ST-segment elevation myocardial infarction treated by primary angioplasty and glycoprotein IIb/IIIa inhibitors: insights from the EGYPT cooperation. <i>Journal of Thrombosis and Thrombolysis</i> , 2010, 30, 342-346.	2.1	38
11	Early abciximab administration before transfer for primary percutaneous coronary interventions for ST-elevation myocardial infarction reduces 1-year mortality in patients with high-risk profile. Results from EUROTRANSFER Registry. <i>American Heart Journal</i> , 2009, 158, 569-575.	2.7	35
12	Impact of smoking status on outcome in patients with ST-segment elevation myocardial infarction treated with primary percutaneous coronary intervention. <i>Journal of Thrombosis and Thrombolysis</i> , 2012, 34, 397-403.	2.1	33
13	Mesh covered stent in ST-segment elevation myocardial infarction. <i>EuroIntervention</i> , 2010, 6, 582-589.	3.2	33
14	Transradial approach in patients with ST-elevation myocardial infarction treated with abciximab results in fewer bleeding complications: data from EUROTRANSFER registry. <i>Coronary Artery Disease</i> , 2010, 21, 292-297.	0.7	31
15	Primary Angioplasty in Patient with St-Segment Elevation Myocardial Infarction in the Setting of Intentional Carbon Monoxide Poisoning. <i>Journal of Emergency Medicine</i> , 2013, 45, 831-834.	0.7	28
16	Admission glucose level and in-hospital outcomes in diabetic and non-diabetic patients with acute myocardial infarction. <i>Clinical Research in Cardiology</i> , 2010, 99, 715-721.	3.3	27
17	Impact of direct stenting on outcome of patients with STâ€”elevation myocardial infarction transferred for primary percutaneous coronary intervention (from the EUROTRANSFER registry). <i>Catheterization and Cardiovascular Interventions</i> , 2014, 84, 925-931.	1.7	27
18	Characteristics of patients presenting with myocardial infarction with non-obstructive coronary arteries (MINOCA) in Poland: data from the ORPKI national registry. <i>Journal of Thrombosis and Thrombolysis</i> , 2019, 47, 462-466.	2.1	27

#	ARTICLE	IF	CITATIONS
19	Plasma asymmetric dimethylarginine is related to anticitrullinated protein antibodies in rheumatoid arthritis of short duration. <i>Metabolism: Clinical and Experimental</i> , 2009, 58, 316-318.	3.4	26
20	Age-related differences in treatment strategies and clinical outcomes in unselected cohort of patients with ST-segment elevation myocardial infarction transferred for primary angioplasty. <i>Journal of Thrombosis and Thrombolysis</i> , 2012, 34, 214-221.	2.1	26
21	Out-of-hospital cardiac arrest in patients treated with primary PCI for STEMI. Long-term follow up data from EUROTRANSFER registry. <i>Resuscitation</i> , 2012, 83, 303-306.	3.0	24
22	Association between endothelial progenitor cell depletion in blood and mild-to-moderate renal insufficiency in stable angina. <i>Nephrology Dialysis Transplantation</i> , 2008, 23, 2265-2273.	0.7	22
23	Association between advanced Killip class at presentation and impaired myocardial perfusion among patients with ST-segment elevation myocardial infarction treated with primary angioplasty and adjunctive glycoprotein IIb/IIIa inhibitors. <i>American Heart Journal</i> , 2009, 158, 416-421.	2.7	21
24	Inter-individual variability in response to clopidogrel in patients with coronary artery disease. <i>Kardiologia Polska</i> , 2005, 62, 108-117; discussion 118.	0.6	21
25	Radial Approach Expertise and Clinical Outcomes of Percutaneous Coronary Interventions Performed Using Femoral Approach. <i>Journal of Clinical Medicine</i> , 2019, 8, 1484.	2.4	20
26	Percutaneous coronary intervention during on- and off-hours in patients with ST-segment elevation myocardial infarction. <i>Hellenic Journal of Cardiology</i> , 2021, 62, 212-218.	1.0	20
27	Asymmetric dimethylarginine predicts decline of glucose tolerance in men with stable coronary artery disease: a 4.5-year follow-up study. <i>Cardiovascular Diabetology</i> , 2013, 12, 64.	6.8	19
28	Acute and long-term outcomes of percutaneous balloon aortic valvuloplasty for the treatment of severe aortic stenosis. <i>Catheterization and Cardiovascular Interventions</i> , 2017, 90, 303-310.	1.7	19
29	Acute myocardial infarction and a new ABCC6 mutation in a 16-year-old boy with pseudoxanthoma elasticum. <i>International Journal of Cardiology</i> , 2007, 116, 261-262.	1.7	18
30	Impact of advanced age on myocardial perfusion, distal embolization, and mortality patients with ST-segment elevation myocardial infarction treated by primary angioplasty and glycoprotein IIb/IIIa inhibitors. <i>Heart and Vessels</i> , 2014, 29, 15-20.	1.2	18
31	Cardioembolic acute myocardial infarction and stroke in a patient with persistent atrial fibrillation. <i>International Journal of Cardiology</i> , 2012, 161, e46-e47.	1.7	17
32	Prevalence and Predictors of Coronary Artery Perforation During Percutaneous Coronary Interventions (from the ORPKI National Registry in Poland). <i>American Journal of Cardiology</i> , 2019, 124, 1186-1189.	1.6	17
33	Impact of bifurcation target lesion on angiographic, electrocardiographic, and clinical outcomes of patients undergoing primary percutaneous coronary intervention (from the Harmonizing Outcomes) <i>Tj ETQq1 1 0.784314 rgBT /Overbo</i> <i>EuroIntervention</i> , 2013, 9, 817-823.	3.2	17
34	Determinants of stroke following percutaneous coronary intervention in acute myocardial infarction (from ORPKI Polish National Registry). <i>International Journal of Cardiology</i> , 2016, 223, 236-238.	1.7	16
35	Prognostic significance of new onset atrial fibrillation in acute coronary syndrome patients treated conservatively. <i>Cardiology Journal</i> , 2010, 17, 57-64.	1.2	16
36	Patency of infarct related artery after pharmacological reperfusion during transfer to primary percutaneous coronary intervention influences left ventricular function and one-year clinical outcome. <i>International Journal of Cardiology</i> , 2008, 124, 326-331.	1.7	15

#	ARTICLE	IF	CITATIONS
37	Impact of Time from Symptom Onset to Drug Administration on Outcome in Patients Undergoing Glycoprotein IIb/IIIa Facilitated Primary Angioplasty (from the EGYPT Cooperation). <i>American Journal of Cardiology</i> , 2015, 115, 711-715.	1.6	15
38	Mesh-Covered Embolic Protection Stent Implantation in ST-Segment Elevation Myocardial Infarction. <i>Circulation: Cardiovascular Interventions</i> , 2015, 8, e001484.	3.9	15
39	Impact of Admission Glucose Level and Presence of Diabetes Mellitus on Mortality in Patients With Non-ST-Segment Elevation Acute Coronary Syndrome Treated Conservatively. <i>American Journal of Cardiology</i> , 2009, 103, 954-958.	1.6	14
40	Ultrasound-guided thrombin injection in the treatment of iatrogenic arterial pseudoaneurysms: Single-center experience. <i>Journal of Clinical Ultrasound</i> , 2014, 42, 24-26.	0.8	14
41	Interval From Initiation of Prasugrel to Coronary Angiography in Patients With Non-ST-Segment Elevation Myocardial Infarction. <i>Journal of the American College of Cardiology</i> , 2019, 73, 906-914.	2.8	14
42	Synergistic effects of asymmetrical dimethyl-L-arginine accumulation and endothelial progenitor cell deficiency on renal function decline during a 2-year follow-up in stable angina. <i>Nephrology Dialysis Transplantation</i> , 2010, 25, 2576-2583.	0.7	13
43	Borderline trend towards long-term mortality benefit from drug eluting stents implantation in ST-elevation myocardial infarction patients in Poland—data from NRDES registry. <i>Catheterization and Cardiovascular Interventions</i> , 2014, 83, 436-442.	1.7	13
44	Contemporary use of P2Y12 inhibitors in patients with ST-segment elevation myocardial infarction referred to primary percutaneous coronary interventions in Poland: Data from ORPKI national registry. <i>Journal of Thrombosis and Thrombolysis</i> , 2018, 45, 151-157.	2.1	13
45	Predictive utility of NT-pro BNP for infarct size and left ventricle function after acute myocardial infarction in long-term follow-up. <i>Disease Markers</i> , 2013, 34, 199-204.	1.3	13
46	Time delay in primary angioplasty: how relevant is it?. <i>Heart</i> , 2007, 93, 1164-1166.	2.9	12
47	More aggressive pharmacological treatment may improve clinical outcome in patients with non-ST-elevation acute coronary syndromes treated conservatively. <i>Coronary Artery Disease</i> , 2007, 18, 299-303.	0.7	12
48	Benefits of pharmacological facilitation with glycoprotein IIb/IIIa inhibitors in diabetic patients undergoing primary angioplasty for STEMI. A subanalysis of the EGYPT cooperation. <i>Journal of Thrombosis and Thrombolysis</i> , 2009, 28, 288-298.	2.1	12
49	Impact of distal embolization on myocardial perfusion and survival among patients undergoing primary angioplasty with glycoprotein IIb/IIIa inhibitors: insights from the EGYPT cooperation. <i>Journal of Thrombosis and Thrombolysis</i> , 2010, 30, 23-28.	2.1	12
50	Predictors and in-hospital outcomes of cardiogenic shock on admission in patients with acute coronary syndromes admitted to hospitals without on-site invasive facilities. <i>Acute Cardiac Care</i> , 2010, 12, 3-9.	0.2	12
51	Drug-eluting versus bare-metal stents in ST-segment elevation myocardial infarction: a mortality analysis from the EUROTRANSFER Registry. <i>Clinical Research in Cardiology</i> , 2011, 100, 139-145.	3.3	12
52	Impact of Hypertension on Distal Embolization, Myocardial Perfusion, and Mortality in Patients With ST Segment Elevation Myocardial Infarction Undergoing Primary Angioplasty. <i>American Journal of Cardiology</i> , 2013, 112, 1083-1086.	1.6	12
53	Impact of advanced age on the safety and effectiveness of paclitaxel-eluting stent implantation in patients with ST-segment elevation myocardial infarction undergoing primary angioplasty. <i>Catheterization and Cardiovascular Interventions</i> , 2013, 82, 869-877.	1.7	12
54	Predictive Utility of NT-pro BNP for Infarct Size and Left Ventricle Function after Acute Myocardial Infarction in Long-Term Follow-Up. <i>Disease Markers</i> , 2013, 34, 199-204.	1.3	12

#	ARTICLE	IF	CITATIONS
55	The network of invasive cardiology facilities in Poland in 2016 (data from the ORPKI Polish National) Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 50 217	0.6	12
56	Effect of introducing a regional 24/7 primary percutaneous coronary intervention service network on treatment outcomes in patients with ST segment elevation myocardial infarction. <i>Kardiologia Polska</i> , 2015, 73, 323-330.	0.6	12
57	Management and mortality in patients with non-ST-segment elevation vs. ST-segment elevation myocardial infarction. Data from the Malopolska Registry of Acute Coronary Syndromes. <i>Kardiologia Polska</i> , 2009, 67, 115-20; discussion 121-2.	0.6	12
58	ST-segment resolution assessed immediately after primary percutaneous coronary intervention correlates with infarct size and left ventricular function in cardiac magnetic resonance at 1-year follow-up. <i>Journal of Electrocardiology</i> , 2009, 42, 152-156.	0.9	11
59	Inverted takotsubo cardiomyopathy in a patient with essential thrombocythemia exposed to anagrelide and phentermine. <i>International Journal of Cardiology</i> , 2012, 160, e31-e32.	1.7	11
60	Predictors of Coronary and Carotid Atherosclerosis in Patients with Severe Degenerative Aortic Stenosis. <i>International Journal of Medical Sciences</i> , 2013, 10, 1361-1366.	2.5	11
61	Abciximab in the management of acute myocardial infarction with ST-segment elevation: evidence-based treatment, current clinical use, and future perspectives. <i>Therapeutics and Clinical Risk Management</i> , 2014, 10, 567.	2.0	11
62	Direct Rapid Left Ventricular Wire Pacing during Balloon Aortic Valvuloplasty. <i>Journal of Clinical Medicine</i> , 2020, 9, 1017.	2.4	11
63	Chronic obstructive pulmonary disease affects the angiographic presentation and outcomes of patients with coronary artery disease treated with percutaneous coronary interventions. <i>Polish Archives of Internal Medicine</i> , 2017, 128, 24-34.	0.4	11
64	Impact of time-to-treatment on myocardial perfusion after primary percutaneous coronary intervention with Gp IIb/IIIa inhibitors. <i>Journal of Cardiovascular Medicine</i> , 2013, 14, 815-820.	1.5	10
65	Circulating biomarkers as predictors of left ventricular remodeling after myocardial infarction. <i>Postepy W Kardiologii Interwencyjnej</i> , 2021, 17, 21-32.	0.2	10
66	Treatment Delay and Clinical Outcomes in Patients with ST-Segment Elevation Myocardial Infarction during the COVID-19 Pandemic. <i>Journal of Clinical Medicine</i> , 2021, 10, 3920.	2.4	10
67	Early abciximab use in ST-elevation myocardial infarction treated with primary percutaneous coronary intervention improves long-term outcome. Data from EUROTRANSFER Registry. <i>Kardiologia Polska</i> , 2010, 68, 539-43.	0.6	10
68	Early administration of abciximab reduces mortality in female patients with ST-elevation myocardial infarction undergoing primary percutaneous coronary intervention (from the EUROTRANSFER) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 217	0.6	10
69	Hospitalization Length after Myocardial Infarction: Risk-Assessment-Based Time of Hospital Discharge vs. Real Life Practice. <i>Journal of Clinical Medicine</i> , 2018, 7, 564.	2.4	9
70	Long-term follow-up of mesh-covered stent implantation in patients with ST-segment elevation myocardial infarction. <i>Kardiologia Polska</i> , 2014, 72, 140-145.	0.6	9
71	Patients with non-ST-elevation myocardial infarction and without chest pain are treated less aggressively and experience higher in-hospital mortality. <i>Kardiologia Polska</i> , 2007, 65, 769-75; discussion 776-7.	0.6	9
72	Relationship between chronic obstructive pulmonary disease and in-hospital management and outcomes in patients with acute myocardial infarction. <i>Kardiologia Polska</i> , 2010, 68, 294-301.	0.6	9

#	ARTICLE	IF	CITATIONS
73	Creatine kinase-MB assessed in patients with acute myocardial infarction correlates with cardiac magnetic resonance infarct size at 6-month follow up. <i>Hellenic Journal of Cardiology</i> , 2014, 55, 4-8.	1.0	9
74	Impact of acute infarct-related artery patency before percutaneous coronary intervention on 30-day outcomes in patients with ST-segment elevation myocardial infarction treated with primary percutaneous coronary intervention in the EUROMAX trial. <i>European Heart Journal: Acute Cardiovascular Care</i> , 2018, 7, 514-521.	1.0	8
75	Clinical outcomes in nonagenarians undergoing a percutaneous coronary intervention. <i>Coronary Artery Disease</i> , 2018, 29, 573-578.	0.7	8
76	Impact of Coronary Artery Disease and Diabetes Mellitus on the Long-Term Follow-Up in Patients after Retrograde Recanalization of the Femoropopliteal Arterial Region. <i>Journal of Diabetes Research</i> , 2019, 2019, 1-6.	2.3	8
77	Radial approach reduces mortality in ST-segment elevation myocardial infarction with cardiogenic shock. <i>Polish Archives of Internal Medicine</i> , 2021, 131, 421-428.	0.4	8
78	Predictors of periprocedural complications in patients undergoing percutaneous coronary interventions within coronary artery bypass grafts. <i>Cardiology Journal</i> , 2020, 26, 633-644.	1.2	8
79	Renal insufficiency increases mortality in acute coronary syndromes regardless of TIMI risk score. <i>Kardiologia Polska</i> , 2008, 66, 28-34; discussion 35-6.	0.6	8
80	In-hospital management and mortality in elderly patients with non-ST-segment elevation acute coronary syndromes treated in centers without on-site invasive facilities. <i>Cardiology Journal</i> , 2008, 15, 451-7.	1.2	8
81	Primary percutaneous coronary intervention during on- vs off-hours in patients with ST-elevation myocardial infarction. Results from EUROTRANSFER Registry. <i>Kardiologia Polska</i> , 2011, 69, 1017-22.	0.6	8
82	Long-Term Follow-up After Retrograde Recanalization of Superficial Femoral Artery Chronic Total Occlusion. <i>Journal of Invasive Cardiology</i> , 2017, 29, 336-339.	0.4	8
83	Lipoma of the aortic valve in a patient with acute myocardial infarction. <i>International Journal of Cardiology</i> , 2007, 115, E36-E38.	1.7	7
84	Early abciximab administration before primary percutaneous coronary intervention improves clinical outcome in elderly patients transferred with ST-elevation myocardial infarction. <i>International Journal of Cardiology</i> , 2010, 143, 147-153.	1.7	7
85	Effects of short-term anti-inflammatory therapy on endothelial function in patients with non-ST-segment elevation acute coronary syndrome. <i>Cardiovascular Revascularization Medicine</i> , 2011, 12, 2-9.	0.8	7
86	Imaging of inflamed carotid artery atherosclerotic plaques with the use of ^{99m} Tc-HYNIC-IL-2 scintigraphy in end-stage renal disease patients. <i>European Journal of Nuclear Medicine and Molecular Imaging</i> , 2012, 39, 673-682.	6.4	7
87	Twelve months follow-up after retrograde recanalization of superficial femoral artery chronic total occlusion. <i>Postępy W Kardiologii Interwencyjnej</i> , 2017, 1, 47-52.	0.2	7
88	Bailout rotational atherectomy in patients with myocardial infarction is not associated with an increased periprocedural complication rate or poorer angiographic outcomes in comparison to elective procedures (from the ORPKI Polish National Registry 2015-2016). <i>Postępy W Kardiologii Interwencyjnej</i> , 2018, 14, 135-143.	0.2	7
89	Impact of On-Site Surgical Backup on Periprocedural Outcomes of Primary Percutaneous Interventions in Patients Presenting With ST-Segment Elevation Myocardial Infarction (From the ORPKI). <i>Tj ETQq1 1106784314rgBT /Ove</i>		
90	Etnografia przedtekstowa. Fenomenologiczne korzenie interpretacji antropologicznej. <i>Teksty Drugie</i> , 2017, 1, 16-39.	0.1	7

#	ARTICLE	IF	CITATIONS
91	Mid-regional pro-adrenomedullin and lactate dehydrogenase as predictors of left ventricular remodeling in patients with myocardial infarction treated with percutaneous coronary intervention. Polish Archives of Internal Medicine, 2021, , .	0.4	7
92	Sex-Related Differences in Outcomes After Percutaneous Balloon Aortic Valvuloplasty. Journal of Invasive Cardiology, 2017, 29, 188-194.	0.4	7
93	Impact of early abciximab administration on myocardial reperfusion in patients with ST-segment elevation myocardial infarction pretreated with 600Åmg of clopidogrel before percutaneous coronary intervention. Journal of Thrombosis and Thrombolysis, 2010, 30, 347-353.	2.1	6
94	Impact of multivessel disease on myocardial perfusion and survival among patients undergoing primary percutaneous coronary intervention with glycoprotein IIb/IIIa inhibitors. Archives of Cardiovascular Diseases, 2013, 106, 155-161.	1.6	6
95	Impact of intra-aortic balloon pump on long-term mortality of unselected patients with ST-segment elevation myocardial infarction complicated by cardiogenic shock. Postępy W Kardiologii Interwencyjnej, 2014, 3, 175-180.	0.2	6
96	No long-term clinical benefit from manual aspiration thrombectomy in ST-elevation myocardial infarction patients. Data from NRDES registry. Catheterization and Cardiovascular Interventions, 2015, 85, E16-22.	1.7	6
97	Circulatory support with Impella CP device during high-risk percutaneous coronary interventions: initial experience in Poland. Postępy W Kardiologii Interwencyjnej, 2016, 3, 254-257.	0.2	6
98	CHA2DS2-VASc and R2-CHA2DS2-VASc scores predict in-hospital and post-discharge outcome in patients with myocardial infarction. Postępy W Kardiologii Interwencyjnej, 2018, 14, 391-398.	0.2	6
99	Chronic obstructive pulmonary disease and periprocedural complications in patients undergoing percutaneous coronary interventions. PLoS ONE, 2018, 13, e0204257.	2.5	6
100	Management of myocardial infarction with ST-segment elevation in district hospitals without catheterisation laboratory–Acute Coronary Syndromes Registry of MaÅ,opolska 2002-2003. Kardiologia Polska, 2006, 64, 1053-60; discussion 1061-2.	0.6	6
101	Local hospital networks for STEMI treatment for a population of half a million inhabitants increase the use of invasive treatment of acute coronary syndromes to the European recommended level. The MaÅ,opolska Registry of Acute Coronary Syndromes 2005-2006. Kardiologia Polska, 2008, 66, 489-97, discussion 498-9.	0.6	6
102	Early abciximab administration before primary percutaneous coronary intervention improves clinical outcome in diabetic patients with ST-segment elevation myocardial infarction (EUROTRANSFER) Tj ETQq0 0 0 rgBT0 Overlocks10 Tf 50 2	0.2	6
103	Efficacy of an Embolic Protection Stent as a Function of Delay to Reperfusion in ST-Segment Elevation Myocardial Infarction (from the MASTER Trial). American Journal of Cardiology, 2014, 114, 1485-1489.	1.6	5
104	The ACEF (age, creatinine, ejection fraction) score predicts ischemic and bleeding outcomes of patients with acute coronary syndromes treated conservatively. Postępy W Kardiologii Interwencyjnej, 2017, 2, 160-164.	0.2	5
105	Coincidence of Moderately Elevated N-Terminal Pro-“B-Type Natriuretic Peptide, Endothelial Progenitor Cells Deficiency and Propensity to Exercise-Induced Myocardial Ischemia in Stable Angina. Disease Markers, 2010, 28, 101-114.	1.3	4
106	Transportation with very long transfer delays (>90 min) for facilitated PCI with reduced-dose fibrinolysis in patients with ST-segment elevation myocardial infarction. International Journal of Cardiology, 2010, 139, 218-227.	1.7	4
107	Diabetes and periprocedural outcomes in patients treated with rotablation during percutaneous coronary interventions. Cardiology Journal, 2020, 27, 152-161.	1.2	4
108	Predictors of syncope in patients with severe aortic stenosis: The role of orthostatic unload test. Cardiology Journal, 2020, 27, 749-755.	1.2	4

#	ARTICLE	IF	CITATIONS
109	Parental history of premature coronary artery disease does not affect plasma levels of asymmetric dimethylarginine in young healthy adults. <i>Polish Archives of Internal Medicine</i> , 2012, 122, 487-493.	0.4	4
110	Comparación de seguridad y efectividad entre los accesos radiales derecho e izquierdo en la intervención coronaria percutánea. <i>Revista Espanola De Cardiologia</i> , 2022, 75, 119-128.	1.2	4
111	Synergistic adverse prognostic effects of asymmetric dimethylarginine and endothelial progenitor-related cells deficiency after elective coronary angioplasty. <i>International Journal of Cardiology</i> , 2011, 152, 400-403.	1.7	3
112	Comparison of safety and effectiveness between the right and left radial artery approach in percutaneous coronary intervention. <i>Revista Espanola De Cardiologia (English Ed)</i> , 2020, 75, 119-119.	0.6	3
113	Long-term outcomes of percutaneous coronary interventions within coronary artery bypass grafts. <i>Archives of Medical Science</i> , 2021, 17, 628-637.	0.9	3
114	Time-related impact of distal embolisation on myocardial perfusion and survival among patients undergoing primary angioplasty with glycoprotein IIb/IIIa inhibitors: insights from the EGYPT cooperation. <i>EuroIntervention</i> , 2012, 8, 470-476.	3.2	3
115	Impact of coronary artery disease presence on the long-term follow-up of carotid artery stenting. <i>Kardiologia Polska</i> , 2015, 73, 274-279.	0.6	3
116	Primary coronary angioplasty in patients with ST segment elevation acute myocardial infarction and diabetes. <i>Kardiologia Polska</i> , 2004, 61, 232-41; discussion 242.	0.6	3
117	Percutaneous peripheral interventions in patients with multivessel coronary artery disease. <i>Kardiologia Polska</i> , 2010, 68, 1115-21.	0.6	3
118	Local administration of abciximab using a ClearWay RX infusion catheter in a patient with acute coronary syndrome caused by late in-stent thrombosis. <i>Kardiologia Polska</i> , 2012, 70, 1199-201.	0.6	3
119	Prehospital Clopidogrel Administration in Patients With ST-Segment Elevation Myocardial Infarction Treated With Primary PCI: Real-Life Experience From the Multicenter NRDES Registry. <i>Journal of Invasive Cardiology</i> , 2016, 28, E56-8.	0.4	3
120	Angiographic perfusion score assessed in patients with acute myocardial infarction is correlated with cardiac magnetic resonance infarct size and N-terminal pro-brain natriuretic peptide in 6-month follow-up. <i>Journal of Thrombosis and Thrombolysis</i> , 2010, 30, 441-445.	2.1	2
121	Methods and techniques Comparison of radiation dose exposure in patients undergoing percutaneous coronary intervention vs. peripheral intervention. <i>Postepy W Kardiologii Interwencyjnej</i> , 2014, 4, 308-313.	0.2	2
122	Introduction of new oral antiplatelet drugs in myocardial infarction hospital network: initial experience. <i>Journal of Thrombosis and Thrombolysis</i> , 2014, 37, 243-245.	2.1	2
123	Chronic obstructive pulmonary disease affects angiographic presentation and outcomes. Authors' reply.. <i>Polish Archives of Internal Medicine</i> , 2018, 128, 195-196.	0.4	2
124	Direct Absorb bioresorbable scaffold implantation in acute coronary syndrome. <i>Kardiologia Polska</i> , 2018, 76, 1434-1440.	0.6	2
125	Percutaneous peripheral interventions in patients with non-ST elevation acute coronary syndromes performed by interventional cardiologists: rationale and results. <i>Kardiologia Polska</i> , 2008, 66, 135-41; discussion 142-3.	0.6	2
126	Effects of early abciximab administration before primary percutaneous coronary intervention on left ventricular function assessed by cardiac magnetic resonance. <i>Kardiologia Polska</i> , 2008, 66, 617-22; discussion 623.	0.6	2

#	ARTICLE	IF	CITATIONS
127	Impact of infarct related artery patency after early abciximab administration on one-year mortality in patients with ST-segment elevation myocardial infarction (data from the EUROTRANSFER Registry). <i>Kardiologia Polska</i> , 2012, 70, 215-21.	0.6	2
128	Risk factors of contrast-induced nephropathy in patients with acute coronary syndrome. <i>Kardiologia Polska</i> , 2022, 80, 760-764.	0.6	2
129	Knowledge and Prevalence of Risk Factors for Coronary Artery Disease in Patients after Percutaneous Coronary Intervention and Coronary Artery Bypass Grafting. <i>Healthcare (Switzerland)</i> , 2022, 10, 1142.	2.0	2
130	Recanalization of peripheral arteries by interventional cardiologists: Rationale and results. <i>International Journal of Cardiology</i> , 2008, 129, 304-306.	1.7	1
131	Long-term follow-up of percutaneous peripheral interventions in lower limb arteries in patients with acute coronary syndrome and diabetes. <i>Postepy W Kardiologii Interwencyjnej</i> , 2010, 3, 117-121.	0.2	1
132	Successful primary angioplasty in patient with ST-segment elevation myocardial infarction caused by large septal branch occlusion. <i>International Journal of Cardiology</i> , 2012, 160, e5-e7.	1.7	1
133	Comment on: Tessari et al. Roles of Insulin, Age, and Asymmetric Dimethylarginine on Nitric Oxide Synthesis In Vivo. <i>Diabetes</i> 2013;62:2699-2708. <i>Diabetes</i> , 2013, 62, e23-e23.	0.6	1
134	Endothelial Progenitor Cells and Long-Term Prognosis in Patients With Stable Angina Treated With Percutaneous Coronary Intervention. <i>Circulation Journal</i> , 2013, 77, 2415.	1.6	1
135	Complete infarct-related artery revascularization in acute myocardial infarction patients. CORAMI Registry. <i>Postepy W Kardiologii Interwencyjnej</i> , 2015, 2, 84-88.	0.2	1
136	Transradial access and the risk of periprocedural stroke. <i>American Heart Journal</i> , 2017, 186, e5-e6.	2.7	1
137	Predictors of mortality and outcomes after retrograde endovascular angioplasty in patients with peripheral artery disease. <i>Postepy W Kardiologii Interwencyjnej</i> , 2019, 15, 234-239.	0.2	1
138	TCT-250 Prevalence and Predictors of Coronary Artery Perforation During Percutaneous Coronary Interventions: Data From the ORPKI National Registry in Poland. <i>Journal of the American College of Cardiology</i> , 2019, 74, B249.	2.8	1
139	Safety of bivalirudin versus unfractionated heparin in endovascular revascularization of peripheral arteries in short- and long-term follow-up. <i>Postepy W Kardiologii Interwencyjnej</i> , 2019, 15, 91-97.	0.2	1
140	Concomitant multi-vessel disease is associated with a lower procedural death rate in patients treated with percutaneous coronary interventions within the left main coronary artery (from the ORPKI). <i>Tj ETQq0 0 0 rgBT (Overlock 10 Tf 50 2</i>		
141	Urgent Pericardiocentesis Is More Frequently Needed After Left Circumflex Coronary Artery Perforation. <i>Journal of Clinical Medicine</i> , 2020, 9, 3043.	2.4	1
142	Etnografia i eksperymenty artystyczne. O powstawaniu nowych p ³ l poznawczych we wsp ³ Å,czesnej antropologii. <i>Teksty Drugie</i> , 2017, 1, 91-110.	0.1	1
143	From pharmacologically assisted early transfer to a universal primary angioplasty service: the experience of the Malopolska region. <i>EuroIntervention</i> , 2012, 8, P51-P54.	3.2	1
144	Etnografia, pami TM Åž, eksperyment: w stron TM alternatywnej historii spo ^Å ecznej. <i>Teksty Drugie</i> , 2016, 6, 268-281.	0.1	1

#	ARTICLE	IF	CITATIONS
145	Sztuka w przestrzeniach wiejskich i eksperymenty etnograficzne. PoÅ¼eganieni kultury zawstydzienia: jednoczasowoÅ, zwrot ku sobie, proto-socjologia. Teksty Drugie, 2016, 4, 66-87.	0.1	1
146	The impact of multiple stent implantation in the infarct-related artery on one-year clinical outcomes of patients with ST-elevation myocardial infarction undergoing primary percutaneous coronary intervention. Data from the Polish NRDES Registry. Kardiologia Polska, 2016, 74, 717-725.	0.6	1
147	Time-to-reperfusion therapy influences outcome of patients with myocardial infarction subjected to facilitated PCI. EuroIntervention, 2005, 1, 309-14.	3.2	1
148	Predictors of infarct-related artery patency following combined lytic therapy in patients with ST-segment elevation myocardial infarction treated with immediate percutaneous coronary intervention. Kardiologia Polska, 2011, 69, 452-7.	0.6	1
149	Use of aspiration thrombectomy in a 102-year-old patient with acute inferior ST-segment elevation myocardial infarction. International Journal of Cardiology, 2012, 160, e46-e47.	1.7	0
150	A 24-year-old male with acute coronary syndrome due to the circumflex coronary artery thrombosis. Diagnostic challenge in everyday practice. International Journal of Cardiology, 2015, 198, 131-133.	1.7	0
151	No clinical benefit from manual thrombus aspiration in patients with non-ST-elevation myocardial infarction. Postepy W Kardiologii Interwencyjnej, 2016, 1, 32-40.	0.2	0
152	Authorsâ€™ response. Kardiologia Polska, 2014, 72, 476-477.	0.6	0
153	Long-term clinical outcomes of direct absorb bioresorbable vascular scaffold implantation in acute coronary syndrome. Minerva Cardioangiologica, 2019, 67, 374-379.	1.2	0
154	Frailty as a Predictor of In-Hospital Outcome in Patients with Myocardial Infarction. Journal of Cardiovascular Development and Disease, 2022, 9, 145.	1.6	0