

Chyung-Ru Wang

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7102114/publications.pdf>

Version: 2024-02-01

83
papers

7,244
citations

76326

40
h-index

60623

81
g-index

88
all docs

88
docs citations

88
times ranked

7953
citing authors

#	ARTICLE	IF	CITATIONS
1	Structure and dynamics of major histocompatibility class Ib molecule H2-M3 complexed with mitochondrial-derived peptides. <i>Journal of Biomolecular Structure and Dynamics</i> , 2022, 40, 10300-10312.	3.5	1
2	Role of Group 1 CD1-restricted T Cells in Host Defense and Inflammatory Diseases. <i>Critical Reviews in Immunology</i> , 2021, 41, 1-21.	0.5	3
3	Mitochondrial metabolism is essential for invariant natural killer T cell development and function. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2021, 118, .	7.1	20
4	CD1-Restricted T Cells in Inflammatory Skin Diseases. <i>Journal of Investigative Dermatology</i> , 2021, , .	0.7	2
5	Type II Natural Killer T Cells Contribute to Protection Against Systemic Methicillin-Resistant <i>Staphylococcus aureus</i> Infection. <i>Frontiers in Immunology</i> , 2020, 11, 610010.	4.8	8
6	USP22 controls iNKT immunity through MED1 suppression of histone H2A monoubiquitination. <i>Journal of Experimental Medicine</i> , 2020, 217, .	8.5	15
7	Group 1 CD1-restricted T cells contribute to control of systemic <i>Staphylococcus aureus</i> infection. <i>PLoS Pathogens</i> , 2020, 16, e1008443.	4.7	11
8	Invariant Natural Killer T-Cells and Total CD1d Restricted Cells Differentially Influence Lipid Metabolism and Atherosclerosis in Low Density Receptor Deficient Mice. <i>International Journal of Molecular Sciences</i> , 2019, 20, 4566.	4.1	5
9	Non-classical Immunity Controls Microbiota Impact on Skin Immunity and Tissue Repair. <i>Cell</i> , 2018, 172, 784-796.e18.	28.9	323
10	Induction of Mycobacterium Tuberculosis Lipid-Specific T Cell Responses by Pulmonary Delivery of Mycolic Acid-Loaded Polymeric Micellar Nanocarriers. <i>Frontiers in Immunology</i> , 2018, 9, 2709.	4.8	37
11	Linking CD1-Restricted T Cells With Autoimmunity and Dyslipidemia: Lipid Levels Matter. <i>Frontiers in Immunology</i> , 2018, 9, 1616.	4.8	17
12	The Lysine Acetyltransferase GCN5 Is Required for iNKT Cell Development through EGR2 Acetylation. <i>Cell Reports</i> , 2017, 20, 600-612.	6.4	30
13	Crosstalk between type II NKT cells and T cells leads to spontaneous chronic inflammatory liver disease. <i>Journal of Hepatology</i> , 2017, 67, 791-800.	3.7	31
14	CD1b-autoreactive T cells contribute to hyperlipidemia-induced skin inflammation in mice. <i>Journal of Clinical Investigation</i> , 2017, 127, 2339-2352.	8.2	59
15	MHC Ib molecule Qa-1 presents Mycobacterium tuberculosis peptide antigens to CD8+ T cells and contributes to protection against infection. <i>PLoS Pathogens</i> , 2017, 13, e1006384.	4.7	47
16	CD1b-autoreactive T cells recognize phospholipid antigens and contribute to antitumor immunity against a CD1b ⁺ T cell lymphoma. <i>OncImmunology</i> , 2016, 5, e1213932.	4.6	22
17	Nonclassical MHC Ib-restricted CD8+ T Cells Recognize Mycobacterium tuberculosis-Derived Protein Antigens and Contribute to Protection Against Infection. <i>PLoS Pathogens</i> , 2016, 12, e1005688.	4.7	20
18	Role of Group 1 CD1-Restricted T Cells in Infectious Disease. <i>Frontiers in Immunology</i> , 2015, 6, 337.	4.8	39

#	ARTICLE	IF	CITATIONS
19	Pak2 Controls Acquisition of NKT Cell Fate by Regulating Expression of the Transcription Factors PLZF and Egr2. <i>Journal of Immunology</i> , 2015, 195, 5272-5284.	0.8	8
20	Mycolic acid-specific T cells protect against <i>Mycobacterium tuberculosis</i> infection in a humanized transgenic mouse model. <i>ELife</i> , 2015, 4, .	6.0	55
21	The adaptor protein SAP regulates type II NKT cell development, cytokine production, and cytotoxicity against lymphoma. <i>European Journal of Immunology</i> , 2014, 44, 3646-3657.	2.9	11
22	Type II natural killer T cells foster the antitumor activity of CpG-oligodeoxynucleotides. <i>Oncolmmunology</i> , 2014, 3, e28977.	4.6	5
23	Polyclonal type II natural killer T cells require PLZF and SAP for their development and contribute to CpG-mediated antitumor response. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2014, 111, 2674-2679.	7.1	61
24	Mitochondria Are Required for Antigen-Specific T Cell Activation through Reactive Oxygen Species Signaling. <i>Immunity</i> , 2013, 38, 225-236.	14.3	981
25	The Functions of Type I and Type II Natural Killer T Cells in Inflammatory Bowel Diseases. <i>Inflammatory Bowel Diseases</i> , 2013, 19, 1330-1338.	1.9	53
26	SAP Is Required for the Development of Innate Phenotype in H2-M3-Restricted CD8+ T Cells. <i>Journal of Immunology</i> , 2012, 189, 4787-4796.	0.8	10
27	Dysregulation of CD1d-Restricted Type II Natural Killer T Cells Leads to Spontaneous Development of Colitis in Mice. <i>Gastroenterology</i> , 2012, 142, 326-334.e2.	1.3	65
28	Recognition of the nonclassical MHC class I molecule H2-M3 by the receptor Ly49A regulates the licensing and activation of NK cells. <i>Nature Immunology</i> , 2012, 13, 1171-1177.	14.5	49
29	Differential requirements for the Ets transcription factor Elf-1 in the development of NKT cells and NK cells. <i>Blood</i> , 2011, 117, 1880-1887.	1.4	48
30	Autoreactive CD1b-restricted T cells: a new innate-like T-cell population that contributes to immunity against infection. <i>Blood</i> , 2011, 118, 3870-3878.	1.4	38
31	Positive selecting cell type determines the phenotype of MHC class Ib-restricted CD8 ⁺ T cells. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2011, 108, 13241-13246.	7.1	27
32	Nonconventional CD8+ T Cell Responses to <i>Listeria</i> Infection in Mice Lacking MHC Class Ia and H2-M3. <i>Journal of Immunology</i> , 2011, 186, 489-498.	0.8	17
33	CD1d-Expressing Breast Cancer Cells Modulate NKT Cell-Mediated Antitumor Immunity in a Murine Model of Breast Cancer Metastasis. <i>PLoS ONE</i> , 2011, 6, e20702.	2.5	85
34	CD1d Expression in Paneth Cells and Rat Exocrine Pancreas Revealed by Novel Monoclonal Antibodies Which Differentially Affect NKT Cell Activation. <i>PLoS ONE</i> , 2010, 5, e13089.	2.5	15
35	Abstract B21: CD1 d-expressing breast cancer cells promote iNKT-mediated antitumor immunity in a mouse model of breast cancer bone metastasis. , 2010, , .		0
36	Polymorphisms in CD1d affect antigen presentation and the activation of CD1d-restricted T cells. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2009, 106, 1909-1914.	7.1	33

#	ARTICLE	IF	CITATIONS
37	CD1-restricted adaptive immune responses to <i>Mycobacteria</i> in human group 1 CD1 transgenic mice. <i>Journal of Experimental Medicine</i> , 2009, 206, 2497-2509.	8.5	105
38	Antagonistic Effect of Toll-Like Receptor Signaling and Bacterial Infections on Transplantation Tolerance. <i>Transplantation</i> , 2009, 87, S77-S79.	1.0	17
39	Prevention of Allograft Tolerance by Bacterial Infection with <i>Listeria monocytogenes</i> . <i>Journal of Immunology</i> , 2008, 180, 5991-5999.	0.8	83
40	An MHC class II-restricted CD8 T cell response confers antiviral immunity. <i>Journal of Experimental Medicine</i> , 2008, 205, 1647-1657.	8.5	30
41	Bacterial infection alters the kinetics and function of iNKT cell responses. <i>Journal of Leukocyte Biology</i> , 2008, 84, 1462-1471.	3.3	22
42	Modulation of NKT Cell Development by B7-CD28 Interaction: An Expanding Horizon for Costimulation. <i>PLoS ONE</i> , 2008, 3, e2703.	2.5	24
43	Characterization of the Natural Killer T-Cell Response in an Adoptive Transfer Model of Atherosclerosis. <i>American Journal of Pathology</i> , 2007, 170, 1100-1107.	3.8	71
44	Serine Protease Inhibitor 6 Protects Cytotoxic T Cells from Self-Inflicted Injury by Ensuring the Integrity of Cytotoxic Granules. <i>Immunity</i> , 2006, 24, 451-461.	14.3	107
45	H2-M3-restricted T cell response to infection. <i>Microbes and Infection</i> , 2006, 8, 2277-2283.	1.9	25
46	Impaired response to <i>Listeria</i> in H2-M3-deficient mice reveals a nonredundant role of MHC class II-specific T cells in host defense. <i>Journal of Experimental Medicine</i> , 2006, 203, 449-459.	8.5	52
47	A Cell-Type Specific CD1d Expression Program Modulates Invariant NKT Cell Development and Function. <i>Journal of Immunology</i> , 2006, 176, 1421-1430.	0.8	40
48	Activating Transcription Factor/cAMP Response Element Binding Protein Family Member Regulated Transcription of CD1A. <i>Journal of Immunology</i> , 2006, 177, 7024-7032.	0.8	17
49	IFN- γ -Mediated Up-Regulation of CD1d in Bacteria-Infected APCs. <i>Journal of Immunology</i> , 2006, 177, 7841-7848.	0.8	43
50	Essential role of TNF family molecule LIGHT as a cytokine in the pathogenesis of hepatitis. <i>Journal of Clinical Investigation</i> , 2006, 116, 1045-1051.	8.2	62
51	Long-term loss of canonical NKT cells following an acute virus infection. <i>European Journal of Immunology</i> , 2005, 35, 879-889.	2.9	45
52	The natural killer T cell ligand β -galactosylceramide prevents or promotes pristane-induced lupus in mice. <i>European Journal of Immunology</i> , 2005, 35, 1143-1154.	2.9	81
53	Transcriptional Regulation of CD1D1 by Ets Family Transcription Factors. <i>Journal of Immunology</i> , 2005, 175, 1022-1029.	0.8	32
54	Glycolipid antigen induces long-term natural killer T cell anergy in mice. <i>Journal of Clinical Investigation</i> , 2005, 115, 2572-2583.	8.2	386

#	ARTICLE	IF	CITATIONS
55	Quantitative and Qualitative Differences in the In Vivo Response of NKT Cells to Distinct $\hat{1}\pm$ - and $\hat{1}^2$ -Anomeric Glycolipids. <i>Journal of Immunology</i> , 2004, 173, 3693-3706.	0.8	136
56	Selective inhibition of anthrax edema factor by adefovir, a drug for chronic hepatitis B virus infection. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2004, 101, 3242-3247.	7.1	109
57	CD1d deficiency exacerbates inflammatory dermatitis in MRL- <i>lpr/lpr</i> mice. <i>European Journal of Immunology</i> , 2004, 34, 1723-1732.	2.9	58
58	The response of natural killer T cells to glycolipid antigens is characterized by surface receptor down-modulation and expansion. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2003, 100, 10913-10918.	7.1	306
59	Expression of CD1d Under the Control of a MHC Class Ia Promoter Skews the Development of NKT Cells, But Not CD8+ T Cells. <i>Journal of Immunology</i> , 2003, 171, 4105-4112.	0.8	25
60	CD1d-expressing Dendritic Cells but Not Thymic Epithelial Cells Can Mediate Negative Selection of NKT Cells. <i>Journal of Experimental Medicine</i> , 2003, 197, 907-918.	8.5	122
61	Functional Roles of TAP and Tapasin in the Assembly of M3- <i>N</i> -Formylated Peptide Complexes. <i>Journal of Immunology</i> , 2001, 167, 1507-1514.	0.8	32
62	Cd1-Restricted Nk T Cells Protect Nonobese Diabetic Mice from Developing Diabetes. <i>Journal of Experimental Medicine</i> , 2001, 194, 313-320.	8.5	251
63	Human CD1d Functions as a Transplantation Antigen and a Restriction Element in Mice. <i>Journal of Immunology</i> , 2001, 166, 3829-3836.	0.8	15
64	Tapasin Enhances Peptide-Induced Expression of H2-M3 Molecules, but Is Not Required for the Retention of Open Conformers. <i>Journal of Immunology</i> , 2001, 167, 2097-2105.	0.8	37
65	Induction of M3-Restricted Cytotoxic T Lymphocyte Responses by N-Formylated Peptides Derived from <i>Mycobacterium tuberculosis</i> . <i>Journal of Experimental Medicine</i> , 2001, 193, 1213-1220.	8.5	65
66	CD1d-Specific NK1.1+ T Cells with a Transgenic Variant TCR. <i>Journal of Immunology</i> , 2000, 165, 168-174.	0.8	74
67	Tracking the Response of Natural Killer T Cells to a Glycolipid Antigen Using Cd1d Tetramers. <i>Journal of Experimental Medicine</i> , 2000, 192, 741-754.	8.5	818
68	Comparative Contribution of CD1 on the Development of CD4+ and CD8+ T Cell Compartments. <i>Journal of Immunology</i> , 2000, 164, 739-745.	0.8	10
69	Cutting Edge: The Ets1 Transcription Factor Is Required for the Development of NK T Cells in Mice. <i>Journal of Immunology</i> , 2000, 164, 2857-2860.	0.8	86
70	MHC Class Ib-Restricted CTL Provide Protection Against Primary and Secondary <i>Listeria monocytogenes</i> Infection. <i>Journal of Immunology</i> , 2000, 165, 5192-5201.	0.8	73
71	Affinity of thymic self-peptides for the TCR determines the selection of CD8+ T lymphocytes in the thymus. <i>International Immunology</i> , 2000, 12, 1353-1363.	4.0	19
72	The Selection of M3-Restricted T Cells Is Dependent on M3 Expression and Presentation of N-Formylated Peptides in the Thymus. <i>Journal of Experimental Medicine</i> , 1999, 190, 1869-1878.	8.5	39

#	ARTICLE	IF	CITATIONS
73	Susceptibility of Mice Deficient in CD1D or TAP1 to Infection with Mycobacterium tuberculosis. Journal of Experimental Medicine, 1999, 189, 1973-1980.	8.5	329
74	The Majority of H2-M3 Is Retained Intracellularly in a Peptide-Receptive State and Traffics to the Cell Surface in the Presence of N-Formylated Peptides. Journal of Experimental Medicine, 1999, 190, 423-434.	8.5	64
75	Selection and Expansion of CD8 α^+ β^+ T Cell Receptor β^+ Intestinal Intraepithelial Lymphocytes in the Absence of Both Classical Major Histocompatibility Complex Class I and Nonclassical Cd1 Molecules. Journal of Experimental Medicine, 1999, 190, 885-890.	8.5	92
76	Tissue distribution, regulation and intracellular localization of murine CD1 molecules. Molecular Immunology, 1998, 35, 525-536.	2.2	82
77	H2-M3, A FULL-SERVICE CLASS I HISTOCOMPATIBILITY ANTIGEN. Annual Review of Immunology, 1997, 15, 851-879.	21.8	125
78	Impaired NK1+ T Cell Development and Early IL-4 Production in CD1-Deficient Mice. Immunity, 1997, 6, 459-467.	14.3	440
79	Identification, expression, and crystallization of the protease-resistant conserved domain of synapsin I. Protein Science, 1997, 6, 2264-2267.	7.6	7
80	Rat RT1 orthologs of mouse H2-M class I genes. Immunogenetics, 1995, 42, 63-67.	2.4	29
81	Nonclassical binding of formylated peptide in crystal structure of the MHC class I molecule H2-M3. Cell, 1995, 82, 655-664.	28.9	151
82	Organization and structure of the H-2M4-M8 class I genes in the mouse major histocompatibility complex. Immunogenetics, 1993, 38, 258-71.	2.4	31
83	H-2M3 encodes the MHC Class I molecule presenting the maternally transmitted antigen of the mouse. Cell, 1991, 66, 335-345.	28.9	117