

# Nabil G Seidah

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/719608/publications.pdf>

Version: 2024-02-01

609  
papers

43,318  
citations

1994

101  
h-index

3732

179  
g-index

633  
all docs

633  
docs citations

633  
times ranked

28861  
citing authors

#	ARTICLE	IF	CITATIONS
1	The Multifaceted Biology of PCSK9. <i>Endocrine Reviews</i> , 2022, 43, 558-582.	20.1	75
2	Caffeine blocks SREBP2-induced hepatic PCSK9 expression to enhance LDLR-mediated cholesterol clearance. <i>Nature Communications</i> , 2022, 13, 770.	12.8	47
3	Distinctive Roles of Furin and TMPRSS2 in SARS-CoV-2 Infectivity. <i>Journal of Virology</i> , 2022, 96, e0012822.	3.4	64
4	Sortilin enhances secretion of apolipoprotein(a) through effects on apolipoprotein B secretion and promotes uptake of lipoprotein(a). <i>Journal of Lipid Research</i> , 2022, 63, 100216.	4.2	4
5	Inhibitory Antibodies against PCSK9 Reduce Surface CD36 and Mitigate Diet-Induced Renal Lipotoxicity. <i>Kidney360</i> , 2022, 3, 1394-1410.	2.1	10
6	PCSK9 Contributes to the Cholesterol, Glucose, and Insulin2 Homeostasis in Seminiferous Tubules and Maintenance of Immunotolerance in Testis. <i>Frontiers in Cell and Developmental Biology</i> , 2022, 10, 889972.	3.7	2
7	Erratum for Essalmani et al., "Distinctive Roles of Furin and TMPRSS2 in SARS-CoV-2 Infectivity", <i>Journal of Virology</i> , 2022, 96, .	3.4	3
8	Post-Transcriptional Effects of miRNAs on PCSK7 Expression and Function: miR-125a-5p, miR-143-3p, and miR-409-3p as Negative Regulators. <i>Metabolites</i> , 2022, 12, 588.	2.9	2
9	The loss-of-function PCSK9Q152H variant increases ER chaperones GRP78 and GRP94 and protects against liver injury. <i>Journal of Clinical Investigation</i> , 2021, 131, .	8.2	29
10	Mutational Spectrum of LDLR and PCSK9 Genes Identified in Iranian Patients With Premature Coronary Artery Disease and Familial Hypercholesterolemia. <i>Frontiers in Genetics</i> , 2021, 12, 625959.	2.3	7
11	L-Carnitine Tartrate Downregulates the ACE2 Receptor and Limits SARS-CoV-2 Infection. <i>Nutrients</i> , 2021, 13, 1297.	4.1	15
12	In Vivo Analysis of the Contribution of Proprotein Convertases to the Processing of FGF23. <i>Frontiers in Endocrinology</i> , 2021, 12, 690681.	3.5	8
13	How Do Enveloped Viruses Exploit the Secretory Proprotein Convertases to Regulate Infectivity and Spread?. <i>Viruses</i> , 2021, 13, 1229.	3.3	18
14	Substantial PCSK9 inactivation in Î²-cells does not modify glucose homeostasis or insulin secretion in mice. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2021, 1866, 158968.	2.4	24
15	Proprotein convertase subtilisin/kexin Type 9 is required for Ahnak-mediated metastasis of melanoma into lung epithelial cells. <i>Neoplasia</i> , 2021, 23, 993-1001.	5.3	12
16	Asialoglycoprotein receptor 1 is a novel PCSK9-independent ligand of liver LDLR cleaved by furin. <i>Journal of Biological Chemistry</i> , 2021, 297, 101177.	3.4	15
17	PCSK9 is not secreted from mature differentiated intestinal cells. <i>Journal of Lipid Research</i> , 2021, 62, 100096.	4.2	4
18	The PCSK9 discovery, an inactive protease with varied functions in hypercholesterolemia, viral infections, and cancer. <i>Journal of Lipid Research</i> , 2021, 62, 100130.	4.2	32

#	ARTICLE	IF	CITATIONS
19	PCSK9 regulates the NODAL signaling pathway and cellular proliferation in hiPSCs. <i>Stem Cell Reports</i> , 2021, 16, 2958-2972.	4.8	7
20	Functional analysis of natural PCSK9 mutants in modern and archaic humans. <i>FEBS Journal</i> , 2020, 287, 515-528.	4.7	8
21	Angiopoietin1 Deficiency in Hepatocytes Affects the Growth of Colorectal Cancer Liver Metastases (CRCLM). <i>Cancers</i> , 2020, 12, 35.	3.7	15
22	Circulating Rather Than Intestinal PCSK9 (Proprotein Convertase Subtilisin Kexin Type 9) Regulates Postprandial Lipemia in Mice. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2020, 40, 2084-2094.	2.4	18
23	Shedding of cancer susceptibility candidate 4 by the convertases PC7/furin unravels a novel secretory protein implicated in cancer progression. <i>Cell Death and Disease</i> , 2020, 11, 665.	6.3	10
24	Molecular evolution of the proopiomelanocortin system in Barn owl species. <i>PLoS ONE</i> , 2020, 15, e0231163.	2.5	3
25	Hypolipidaemia among patients with PMM2-CDG is associated with low circulating PCSK9 levels: a case report followed by observational and experimental studies. <i>Journal of Medical Genetics</i> , 2020, 57, 11-17.	3.2	8
26	The motif EXEXXL in the cytosolic tail of the secretory human proprotein convertase PC7 regulates its trafficking and cleavage activity. <i>Journal of Biological Chemistry</i> , 2020, 295, 2068-2083.	3.4	5
27	Circulating PCSK9 is associated with liver biomarkers and hepatic steatosis. <i>Clinical Biochemistry</i> , 2020, 77, 20-25.	1.9	26
28	Proprotein convertase 7 (PCSK7) reduces apoA levels. <i>FEBS Journal</i> , 2020, 287, 3565-3578.	4.7	13
29	Dengue virus induces PCSK9 expression to alter antiviral responses and disease outcomes. <i>Journal of Clinical Investigation</i> , 2020, 130, 5223-5234.	8.2	41
30	A novel cell-based sensor detecting the activity of individual basic proprotein convertases. <i>FEBS Journal</i> , 2019, 286, 4597-4620.	4.7	4
31	Novel strategies to target proprotein convertase subtilisin kexin 9: beyond monoclonal antibodies. <i>Cardiovascular Research</i> , 2019, 115, 510-518.	3.8	63
32	Ser-Phosphorylation of PCSK9 (Proprotein Convertase Subtilisin-Kexin 9) by Fam20C (Family With Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50	2.4	36
33	HIV-induced neuroinflammation: impact of PAR1 and PAR2 processing by Furin. <i>Cell Death and Differentiation</i> , 2019, 26, 1942-1954.	11.2	11
34	The Elusive Inhibitory Function of the Acidic N-Terminal Segment of the Prodomain of PCSK9: The Plot Thickens. <i>Journal of Molecular Biology</i> , 2019, 431, 904-907.	4.2	10
35	Diet-induced hepatic steatosis abrogates cell-surface LDLR by inducing de novo PCSK9 expression in mice. <i>Journal of Biological Chemistry</i> , 2019, 294, 9037-9047.	3.4	40
36	Pcsk9 knockout exacerbates diet-induced non-alcoholic steatohepatitis, fibrosis and liver injury in mice. <i>JHEP Reports</i> , 2019, 1, 418-429.	4.9	51

#	ARTICLE	IF	CITATIONS
37	Variation in Serum PCSK9 (Proprotein Convertase Subtilisin/Kexin Type 9), Cardiovascular Disease Risk, and an Investigation of Potential Unanticipated Effects of PCSK9 Inhibition. <i>Circulation Genomic and Precision Medicine</i> , 2019, 12, e002335.	3.6	11
38	New Sequencing technologies help revealing unexpected mutations in Autosomal Dominant Hypercholesterolemia. <i>Scientific Reports</i> , 2018, 8, 1943.	3.3	25
39	The ever-expanding saga of the proprotein convertases and their roles in body homeostasis. <i>Current Opinion in Lipidology</i> , 2018, 29, 144-150.	2.7	30
40	Posttranslational modification of proprotein convertase subtilisin/kexin type 9 is differentially regulated in response to distinct cardiometabolic treatments as revealed by targeted proteomics. <i>Journal of Clinical Lipidology</i> , 2018, 12, 1027-1038.	1.5	10
41	Loss-of-function PCSK9 mutants evade the unfolded protein response sensor GRP78 and fail to induce endoplasmic reticulum stress when retained. <i>Journal of Biological Chemistry</i> , 2018, 293, 7329-7343.	3.4	29
42	Site-specific O-glycosylation of members of the low-density lipoprotein receptor superfamily enhances ligand interactions. <i>Journal of Biological Chemistry</i> , 2018, 293, 7408-7422.	3.4	57
43	Osteopontin as a novel substrate for the proprotein convertase 5/6 (PCSK5) in bone. <i>Bone</i> , 2018, 107, 45-55.	2.9	14
44	Transcriptome Analysis Reveals Nonfoamy Rather Than Foamy Plaque Macrophages Are Proinflammatory in Atherosclerotic Murine Models. <i>Circulation Research</i> , 2018, 123, 1127-1142.	4.5	275
45	Functional Analysis of Novel PCSK9 Mutants from Sapiens (R96L, R105W and P174S) and Denisovan (H449L). <i>Atherosclerosis Supplements</i> , 2018, 32, 60.	1.2	0
46	A single domain antibody against the Cys- and His-rich domain of PCSK9 and evolocumab exhibit different inhibition mechanisms in humanized PCSK9 mice. <i>Biological Chemistry</i> , 2018, 399, 1363-1374.	2.5	10
47	Low-density lipoprotein (LDL)-dependent uptake of Gram-positive lipoteichoic acid and Gram-negative lipopolysaccharide occurs through LDL receptor. <i>Scientific Reports</i> , 2018, 8, 10496.	3.3	47
48	PCSK9 REDUCES HEPATIC LIPID CONTENT AND CONFERS PROTECTION AGAINST ER STRESS AND ROS IN HEPG2 CELLS. <i>FASEB Journal</i> , 2018, 32, 539.8.	0.5	0
49	Loss-of-function PCSK9 mutants evade the unfolded protein response sensor, GRP78, and fail to induce endoplasmic reticulum stress when retained. <i>FASEB Journal</i> , 2018, 32, 793.7.	0.5	0
50	Prohormone convertase 7 is necessary for the normal processing of cholecystokinin in mouse brain. <i>Biochemical and Biophysical Research Communications</i> , 2017, 482, 1190-1193.	2.1	2
51	Thrombin activation of protein C requires prior processing by a liver proprotein convertase. <i>Journal of Biological Chemistry</i> , 2017, 292, 10564-10573.	3.4	10
52	Can iPCSK9-induced hypocholesterolemia starve cancer cells?. <i>Journal of Clinical Lipidology</i> , 2017, 11, 600-601.	1.5	5
53	Endoplasmic Reticulum Stress and Ca <sup>2+</sup> Depletion Differentially Modulate the Sterol Regulatory Protein PCSK9 to Control Lipid Metabolism. <i>Journal of Biological Chemistry</i> , 2017, 292, 1510-1523.	3.4	31
54	The Proprotein Convertases in Hypercholesterolemia and Cardiovascular Diseases: Emphasis on Proprotein Convertase Subtilisin/Kexin 9. <i>Pharmacological Reviews</i> , 2017, 69, 33-52.	16.0	90

#	ARTICLE	IF	CITATIONS
55	The Proprotein Convertase Subtilisin/Kexin Type 9-resistant R410S Low Density Lipoprotein Receptor Mutation. <i>Journal of Biological Chemistry</i> , 2017, 292, 1573-1590.	3.4	30
56	Insights into a PCSK9 structural groove: a harbinger of new drugs to reduce LDL-cholesterol. <i>Nature Structural and Molecular Biology</i> , 2017, 24, 785-786.	8.2	8
57	Effect of the p.Arg357His mutation of PCSK9 on basal and postprandial lipoprotein metabolism. <i>Atherosclerosis</i> , 2017, 263, e2.	0.8	0
58	A case of hypocholesterolemia and steatosis in a carrier of a PCSK9 loss-of-function mutation and polymorphisms predisposing to nonalcoholic fatty liver disease. <i>Journal of Clinical Lipidology</i> , 2017, 11, 1101-1105.	1.5	10
59	Association Between Plasma Proprotein Convertase Subtilisin/Kexin Type 9 and the Presence of Metabolic Syndrome in a Predominantly Rural-Based Sub-Saharan African Population. <i>Metabolic Syndrome and Related Disorders</i> , 2017, 15, 423-429.	1.3	13
60	Pcsk5 is required in the early cranio-cardiac mesoderm for heart development. <i>BMC Developmental Biology</i> , 2017, 17, 6.	2.1	10
61	Post-transcriptional Regulation of PCSK9 by miR-191, miR-222, and miR-224. <i>Frontiers in Genetics</i> , 2017, 8, 189.	2.3	40
62	New insights of altered lipid profile in Fragile X Syndrome. <i>PLoS ONE</i> , 2017, 12, e0174301.	2.5	26
63	The PCSK9 revolution and the potential of PCSK9-based therapies to reduce LDL-cholesterol. <i>Global Cardiology Science &amp; Practice</i> , 2017, 2017, e201702.	0.4	31
64	Proprotein convertase furin regulates osteocalcin and bone endocrine function. <i>Journal of Clinical Investigation</i> , 2017, 127, 4104-4117.	8.2	55
65	Roles of the low density lipoprotein receptor and related receptors in inhibition of lipoprotein(a) internalization by proprotein convertase subtilisin/kexin type 9. <i>PLoS ONE</i> , 2017, 12, e0180869.	2.5	40
66	Proprotein Convertase Subtilisin/Kexin type 9 affects insulin but not lipid metabolism in cystic fibrosis. <i>Clinical and Investigative Medicine</i> , 2017, 40, 59.	0.6	1
67	Neuroendocrine Peptide Processing $\alpha$ - $\tau$ . , 2017, , .		0
68	Role of Ostm1 Cytosolic Complex with Kinesin 5B in Intracellular Dispersion and Trafficking. <i>Molecular and Cellular Biology</i> , 2016, 36, 507-521.	2.3	18
69	New developments in proprotein convertase subtilisin/kexin 9's biology and clinical implications. <i>Current Opinion in Lipidology</i> , 2016, 27, 274-281.	2.7	47
70	Deferoxamine stimulates LDLR expression and LDL uptake in HepG2 cells. <i>Molecular Nutrition and Food Research</i> , 2016, 60, 600-608.	3.3	11
71	Back cover: Deferoxamine stimulates LDLR expression and LDL uptake in HepG2 cells. <i>Molecular Nutrition and Food Research</i> , 2016, 60, NA.	3.3	0
72	Proprotein convertase subtilisin / kexin 9 (PCSK9) inhibitors and the future of dyslipidemia therapy: an updated patent review (2011-2015). <i>Expert Opinion on Therapeutic Patents</i> , 2016, 26, 1377-1392.	5.0	23

#	ARTICLE	IF	CITATIONS
73	Differential Expression of PCSK9 Modulates Infection, Inflammation, and Coagulation in a Murine Model of Sepsis. <i>Shock</i> , 2016, 46, 672-680.	2.1	110
74	Mechanism of Folding and Activation of Subtilisin Kexin Isozyme-1 (SKI-1)/Site-1 Protease (S1P). <i>Journal of Biological Chemistry</i> , 2016, 291, 2055-2066.	3.4	13
75	PCSK9 inhibition-mediated reduction in Lp(a) with evolocumab: an analysis of 10 clinical trials and the LDL receptor's role. <i>Journal of Lipid Research</i> , 2016, 57, 1086-1096.	4.2	180
76	An Unbiased Mass Spectrometry Approach Identifies Glypican-3 as an Interactor of Proprotein Convertase Subtilisin/Kexin Type 9 (PCSK9) and Low Density Lipoprotein Receptor (LDLR) in Hepatocellular Carcinoma Cells. <i>Journal of Biological Chemistry</i> , 2016, 291, 24676-24687.	3.4	14
77	Proprotein Convertase Subtilisin/Kexin Type 9 (PCSK9) Single Domain Antibodies Are Potent Inhibitors of Low Density Lipoprotein Receptor Degradation. <i>Journal of Biological Chemistry</i> , 2016, 291, 16659-16671.	3.4	28
78	Evaluating The Roles Of PCSK9 And Specific Receptors In Lipoprotein(A) Catabolism. <i>Journal of Clinical Lipidology</i> , 2016, 10, 720-721.	1.5	3
79	Reducing Vascular Calcification by Anti-IL-1 $\beta$ Monoclonal Antibody in a Mouse Model of Familial Hypercholesterolemia. <i>Angiology</i> , 2016, 67, 157-167.	1.8	44
80	Statins can exert dual, concentration dependent effects on HCV entry in vitro. <i>Antiviral Research</i> , 2016, 128, 43-48.	4.1	10
81	Deletion of Mbtps1 (Pcsk8, S1p, Ski-1) Gene in Osteocytes Stimulates Soleus Muscle Regeneration and Increased Size and Contractile Force with Age. <i>Journal of Biological Chemistry</i> , 2016, 291, 4308-4322.	3.4	42
82	The PCSK9 revolution and the potential of PCSK9-based therapies to reduce LDL-cholesterol. <i>Global Cardiology Science &amp; Practice</i> , 2015, 2015, 59.	0.4	2
83	PACE4 (PCSK6): another proprotein convertase link to iron homeostasis?. <i>Haematologica</i> , 2015, 100, e377-e377.	3.5	22
84	Decreased PCSK9 expression in human hepatocellular carcinoma. <i>BMC Gastroenterology</i> , 2015, 15, 176.	2.0	46
85	Cleavage of a Neuroinvasive Human Respiratory Virus Spike Glycoprotein by Proprotein Convertases Modulates Neurovirulence and Virus Spread within the Central Nervous System. <i>PLoS Pathogens</i> , 2015, 11, e1005261.	4.7	62
86	Liver-Specific Inactivation of the Proprotein Convertase FURIN Leads to Increased Hepatocellular Carcinoma Growth. <i>BioMed Research International</i> , 2015, 2015, 1-8.	1.9	15
87	Lipoprotein(a) Catabolism Is Regulated by Proprotein Convertase Subtilisin/Kexin Type 9 through the Low Density Lipoprotein Receptor. <i>Journal of Biological Chemistry</i> , 2015, 290, 11649-11662.	3.4	176
88	MBTPS1/SKI-1/S1P proprotein convertase is required for ECM signaling and axial elongation during somitogenesis and vertebral development. <i>Human Molecular Genetics</i> , 2015, 24, 2884-2898.	2.9	23
89	High-fructose feeding promotes accelerated degradation of hepatic LDL receptor and hypercholesterolemia in hamsters via elevated circulating PCSK9 levels. <i>Atherosclerosis</i> , 2015, 239, 364-374.	0.8	29
90	SKI-1/S1P inhibitor PF-429242 impairs the onset of HCV infection. <i>Antiviral Research</i> , 2015, 115, 94-104.	4.1	31

#	ARTICLE	IF	CITATIONS
91	Familial hypercholesterolemia mutations in the Middle Eastern and North African region: A need for a national registry. <i>Journal of Clinical Lipidology</i> , 2015, 9, 187-194.	1.5	44
92	A semi-automated mass spectrometric immunoassay coupled to selected reaction monitoring (MSIA- $\epsilon$ -SRM) reveals novel relationships between circulating PCSK9 and metabolic phenotypes in patient cohorts. <i>Methods</i> , 2015, 81, 66-73.	3.8	23
93	Amyloid Precursor-like Protein 2 and Sortilin Do Not Regulate the PCSK9 Convertase-mediated Low Density Lipoprotein Receptor Degradation but Interact with Each Other. <i>Journal of Biological Chemistry</i> , 2015, 290, 18609-18620.	3.4	47
94	Plasma Membrane Tetraspanin CD81 Complexes with Proprotein Convertase Subtilisin/Kexin Type 9 (PCSK9) and Low Density Lipoprotein Receptor (LDLR), and Its Levels Are Reduced by PCSK9. <i>Journal of Biological Chemistry</i> , 2015, 290, 23385-23400.	3.4	22
95	PCSK9 deficiency unmasks a sex- and tissue-specific subcellular distribution of the LDL and VLDL receptors in mice. <i>Journal of Lipid Research</i> , 2015, 56, 2133-2142.	4.2	45
96	A novel assay uncovers an unexpected role for SR-BI in LDL transcytosis. <i>Cardiovascular Research</i> , 2015, 108, 268-277.	3.8	112
97	Neuroinflammation-Induced Interactions between Protease-Activated Receptor 1 and Proprotein Convertases in HIV-Associated Neurocognitive Disorder. <i>Molecular and Cellular Biology</i> , 2015, 35, 3684-3700.	2.3	29
98	PCSK9, apolipoprotein E and lipoviral particles in chronic hepatitis C genotype 3: Evidence for genotype-specific regulation of lipoprotein metabolism. <i>Journal of Hepatology</i> , 2015, 62, 763-770.	3.7	33
99	Zymogen Activation and Subcellular Activity of Subtilisin Kexin Isozyme 1/Site 1 Protease. <i>Journal of Biological Chemistry</i> , 2014, 289, 35743-35756.	3.4	18
100	Is there a link between proprotein convertase PC7 activity and human lipid homeostasis?. <i>FEBS Open Bio</i> , 2014, 4, 741-745.	2.3	9
101	Clearance of Plasma Proprotein Convertase Subtilisin/Kexin 9 by Low-Density Lipoprotein Apheresis. <i>Circulation Research</i> , 2014, 115, e3-4.	4.5	6
102	The effect of insulin on circulating PCSK9 in postmenopausal obese women. <i>Clinical Biochemistry</i> , 2014, 47, 1033-1039.	1.9	47
103	Prodomain of the proprotein convertase subtilisin/kexin Furin (ppFurin) protects from tumor progression and metastasis. <i>Carcinogenesis</i> , 2014, 35, 528-536.	2.8	22
104	Low Density Lipoprotein Receptor Class A Repeats Are O-Glycosylated in Linker Regions. <i>Journal of Biological Chemistry</i> , 2014, 289, 17312-17324.	3.4	46
105	PCSK9. <i>Circulation Research</i> , 2014, 114, 1022-1036.	4.5	495
106	Annexin A2 Reduces PCSK9 Protein Levels via a Translational Mechanism and Interacts with the M1 and M2 Domains of PCSK9. <i>Journal of Biological Chemistry</i> , 2014, 289, 17732-17746.	3.4	40
107	SORCS1 is necessary for normal insulin secretory granule biogenesis in metabolically stressed $\beta^2$ cells. <i>Journal of Clinical Investigation</i> , 2014, 124, 4240-4256.	8.2	53
108	Furin Is the Major Proprotein Convertase Required for KISS1-to-Kisspeptin Processing. <i>PLoS ONE</i> , 2014, 9, e84958.	2.5	21

#	ARTICLE	IF	CITATIONS
109	Abstract 135: Phosphorylation and furin-mediated processing are critical posttranslational modifications of the KISS1 metastasis suppressor. , 2014, , .		0
110	Implication of the proprotein convertases in iron homeostasis: Proprotein convertase 7 sheds human transferrin receptor 1 and furin activates hepcidin. Hepatology, 2013, 57, 2514-2524.	7.3	57
111	Aortic calcification: Novel insights from familial hypercholesterolemia and potential role for the low-density lipoprotein receptor. Atherosclerosis, 2013, 226, 9-15.	0.8	130
112	Processing of Human Toll-like Receptor 7 by Furin-like Proprotein Convertases Is Required for Its Accumulation and Activity in Endosomes. Immunity, 2013, 39, 711-721.	14.3	77
113	Beyond LDL-C lowering: Distinct molecular sphingolipids are good indicators of proprotein convertase subtilisin/kexin type 9 (PCSK9) deficiency. Atherosclerosis, 2013, 228, 380-385.	0.8	34
114	APOE p.Leu167del mutation in familial hypercholesterolemia. Atherosclerosis, 2013, 231, 218-222.	0.8	84
115	Rapid development of sensitive, high-throughput, quantitative and highly selective mass spectrometric targeted immunoassays for clinically important proteins in human plasma and serum. Clinical Biochemistry, 2013, 46, 399-410.	1.9	98
116	PCSK9 plays a significant role in cholesterol homeostasis and lipid transport in intestinal epithelial cells. Atherosclerosis, 2013, 227, 297-306.	0.8	118
117	Regional Distribution and Metabolic Effect of PCSK9 insLEU and R46L Gene Mutations and apoE Genotype. Canadian Journal of Cardiology, 2013, 29, 927-933.	1.7	32
118	Viral envelope glycoprotein processing by proprotein convertases. Antiviral Research, 2013, 99, 49-60.	4.1	22
119	Furin Is the Primary in Vivo Convertase of Angiopoietin-like 3 and Endothelial Lipase in Hepatocytes. Journal of Biological Chemistry, 2013, 288, 26410-26418.	3.4	43
120	The Cytosolic Adaptor <math>AP-1A</math> Is Essential for the Trafficking and Function of Niemann-Pick Type C Proteins. Traffic, 2013, 14, 458-469.	2.7	17
121	Differential Recognition of Old World and New World Arenavirus Envelope Glycoproteins by Subtilisin Kexin Isozyme 1 (SKI-1)/Site 1 Protease (S1P). Journal of Virology, 2013, 87, 6406-6414.	3.4	18
122	The Multifaceted Proprotein Convertases: Their Unique, Redundant, Complementary, and Opposite Functions. Journal of Biological Chemistry, 2013, 288, 21473-21481.	3.4	151
123	Disruption of the expression of the proprotein convertase PC7 reduces BDNF production and affects learning and memory in mice. Proceedings of the National Academy of Sciences of the United States of America, 2013, 110, 17362-17367.	7.1	74
124	Reply:. Hepatology, 2013, 58, 1861-1862.	7.3	1
125	Proprotein Convertase Furin and Proprotein Convertase PC5/6. , 2013, , 1803-1811.		1
126	Proprotein Convertases PC4, PACE4, and PC7. , 2013, , 1812-1820.		0



#	ARTICLE	IF	CITATIONS
127	Proprotein Convertases SKI-1/S1P and PCSK9. , 2013, , 1821-1828.		0
128	Site-1 Protease. , 2013, , 3265-3270.		2
129	&lt;em>In utero&lt;/em> Measurement of Heart Rate in Mouse by Noninvasive M-mode Echocardiography. Journal of Visualized Experiments, 2013, , e50994.	0.3	4
130	Modulation of Protease Activated Receptor 1 Influences Human Metapneumovirus Disease Severity in a Mouse Model. PLoS ONE, 2013, 8, e72529.	2.5	33
131	Proprotein Convertase Subtilisin Kexin 9 (PCSK9) Inhibitors in the Treatment of Hypercholesterolemia and other Pathologies. Current Pharmaceutical Design, 2013, 19, 3161-3172.	1.9	70
132	Evidence from a Randomized Trial That Simvastatin, but Not Ezetimibe, Upregulates Circulating PCSK9 Levels. PLoS ONE, 2013, 8, e60095.	2.5	54
133	Proprotein Convertase Subtilisin/Kexin Type 9 (PCSK9) Can Mediate Degradation of the Low Density Lipoprotein Receptor-Related Protein 1 (LRP-1). PLoS ONE, 2013, 8, e64145.	2.5	183
134	PCSK9 Prosegment Chimera as Novel Inhibitors of LDLR Degradation. PLoS ONE, 2013, 8, e72113.	2.5	20
135	Proprotein Convertase 1/3. , 2013, , 3286-3290.		0
136	Proprotein Convertase 5. , 2013, , 3305-3310.		0
137	Proprotein Convertase 2. , 2013, , 3290-3295.		6
138	Proprotein Convertase 7. , 2013, , 3310-3314.		0
139	Proprotein Convertase PCSK9. , 2013, , 3315-3322.		1
140	Proprotein Convertase 4. , 2013, , 3295-3299.		0
141	Abstract 3860: Furin is required for processing KISS1 to kisspeptins.. , 2013, , .		0
142	Differential expression and processing of secretogranin II in relation to the status of pheochromocytoma: implications for the production of the tumoral marker EM66. Journal of Molecular Endocrinology, 2012, 48, 115-127.	2.5	11
143	Loss- and Gain-of-function PCSK9 Variants. Journal of Biological Chemistry, 2012, 287, 33745-33755.	3.4	71
144	Disruption of Proprotein Convertase 1/3 (PC1/3) Expression in Mice Causes Innate Immune Defects and Uncontrolled Cytokine Secretion. Journal of Biological Chemistry, 2012, 287, 14703-14717.	3.4	32

#	ARTICLE	IF	CITATIONS
145	The M2 Module of the Cys-His-rich Domain (CHRD) of PCSK9 Protein Is Needed for the Extracellular Low-density Lipoprotein Receptor (LDLR) Degradation Pathway. <i>Journal of Biological Chemistry</i> , 2012, 287, 43492-43501.	3.4	62
146	Rosuvastatin, Proprotein Convertase Subtilisin/Kexin Type 9 Concentrations, and LDL Cholesterol Response: the JUPITER Trial. <i>Clinical Chemistry</i> , 2012, 58, 183-189.	3.2	133
147	Molecular Characterization of the Processing of Arenavirus Envelope Glycoprotein Precursors by Subtilisin Kexin Isozyme-1/Site-1 Protease. <i>Journal of Virology</i> , 2012, 86, 4935-4946.	3.4	34
148	Mouse Model of Intraluminal MCAO: Cerebral Infarct Evaluation by Cresyl Violet Staining. <i>Journal of Visualized Experiments</i> , 2012, , .	0.3	62
149	Identification and characterization of new gain-of-function mutations in the PCSK9 gene responsible for autosomal dominant hypercholesterolemia. <i>Atherosclerosis</i> , 2012, 223, 394-400.	0.8	92
150	Automated design of ligands to polypharmacological profiles. <i>Nature</i> , 2012, 492, 215-220.	27.8	698
151	Effect of the Mediterranean diet with and without weight loss on surrogate markers of cholesterol homeostasis in men with the metabolic syndrome. <i>British Journal of Nutrition</i> , 2012, 107, 705-711.	2.3	58
152	Proprotein Convertase Subtilisin/Kexin Type 9 Deficiency Reduces Melanoma Metastasis in Liver. <i>Neoplasia</i> , 2012, 14, 1122-IN5.	5.3	94
153	SKI-1 and Furin Generate Multiple RGMa Fragments that Regulate Axonal Growth. <i>Developmental Cell</i> , 2012, 22, 391-402.	7.0	56
154	Opposite Roles of Furin and PC5A in N-Cadherin Processing. <i>Neoplasia</i> , 2012, 14, 880-IN3.	5.3	23
155	The biology and therapeutic targeting of the proprotein convertases. <i>Nature Reviews Drug Discovery</i> , 2012, 11, 367-383.	46.4	647
156	Loss of Endothelial Furin Leads to Cardiac Malformation and Early Postnatal Death. <i>Molecular and Cellular Biology</i> , 2012, 32, 3382-3391.	2.3	43
157	Gene Inactivation of Proprotein Convertase Subtilisin/Kexin Type 9 Reduces Atherosclerosis in Mice. <i>Circulation</i> , 2012, 125, 894-901.	1.6	193
158	SKI-1/S1P inhibition: A promising surrogate to statins to block Hepatitis C virus replication. <i>Antiviral Research</i> , 2012, 95, 159-166.	4.1	27
159	5â€ŒThiomannosides Block the Biosynthesis of Dolicholâ€ŒLinked Oligosaccharides and Mimic Class I Congenital Disorders of Glycosylation. <i>ChemBioChem</i> , 2012, 13, 392-401.	2.6	2
160	Annexin A2 Is a Natural Extrahepatic Inhibitor of the PCSK9-Induced LDL Receptor Degradation. <i>PLoS ONE</i> , 2012, 7, e41865.	2.5	98
161	Quantitative Proteomic Analysis of PCSK9 Gain of Function in Human Hepatic HuH7 Cells. <i>Journal of Proteome Research</i> , 2011, 10, 2011-2026.	3.7	15
162	The LDLR deficient mouse as a model for aortic calcification and quantification by micro-computed tomography. <i>Atherosclerosis</i> , 2011, 219, 455-462.	0.8	54

#	ARTICLE	IF	CITATIONS
163	What lies ahead for the proprotein convertases?. <i>Annals of the New York Academy of Sciences</i> , 2011, 1220, 149-161.	3.8	74
164	Arenavirus envelope glycoproteins mimic autoprocessing sites of the cellular proprotein convertase subtilisin kexin isozyme-1/site-1 protease. <i>Virology</i> , 2011, 417, 18-26.	2.4	23
165	Inactivation of endothelial proprotein convertase 5/6 decreases collagen deposition in the cardiovascular system: role of fibroblast autophagy. <i>Journal of Molecular Medicine</i> , 2011, 89, 1103-1111.	3.9	25
166	N-Glycosylation controls trafficking, zymogen activation and substrate processing of proprotein convertases PC1/3 and subtilisin kexin isozyme-1. <i>Glycobiology</i> , 2011, 21, 1290-1300.	2.5	19
167	Atorvastatin increases intestinal expression of NPC1L1 in hyperlipidemic men. <i>Journal of Lipid Research</i> , 2011, 52, 558-565.	4.2	92
168	PCSK9 reduces the protein levels of the LDL receptor in mouse brain during development and after ischemic stroke. <i>Journal of Lipid Research</i> , 2011, 52, 1383-1391.	4.2	77
169	Proprotein Convertase PC7 Enhances the Activation of the EGF Receptor Pathway through Processing of the EGF Precursor. <i>Journal of Biological Chemistry</i> , 2011, 286, 9185-9195.	3.4	26
170	Furin Is the Major Processing Enzyme of the Cardiac-specific Growth Factor Bone Morphogenetic Protein 10. <i>Journal of Biological Chemistry</i> , 2011, 286, 22785-22794.	3.4	52
171	In Vivo Evidence That Furin from Hepatocytes Inactivates PCSK9. <i>Journal of Biological Chemistry</i> , 2011, 286, 4257-4263.	3.4	132
172	The Proprotein Convertase PC7. <i>Journal of Biological Chemistry</i> , 2011, 286, 2728-2738.	3.4	47
173	A Novel Mouse Model of Alzheimer's Disease with Chronic Estrogen Deficiency Leads to Glial Cell Activation and Hypertrophy. <i>Journal of Aging Research</i> , 2011, 2011, 1-12.	0.9	21
174	Inhibition of Proprotein Convertase SKI-1 Blocks Transcription of Key Extracellular Matrix Genes Regulating Osteoblastic Mineralization. <i>Journal of Biological Chemistry</i> , 2011, 286, 1836-1849.	3.4	37
175	Circulating Proprotein Convertase Subtilisin/Kexin 9 (PCSK9) Regulates VLDLR Protein and Triglyceride Accumulation in Visceral Adipose Tissue. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2011, 31, 785-791.	2.4	220
176	Novel Loss-of-Function PCSK9 Variant Is Associated with Low Plasma LDL Cholesterol in a French-Canadian Family and with Impaired Processing and Secretion in Cell Culture. <i>Clinical Chemistry</i> , 2011, 57, 1415-1423.	3.2	101
177	Latent Transforming Growth Factor $\beta$ 2-Binding Proteins-2 and -3 Inhibit the Proprotein Convertase 5/6A. <i>Journal of Biological Chemistry</i> , 2011, 286, 29063-29073.	3.4	20
178	The Proprotein Convertases, 20 Years Later. <i>Methods in Molecular Biology</i> , 2011, 768, 23-57.	0.9	140
179	The Influence of PCSK9 Polymorphisms on Serum Low-Density Lipoprotein Cholesterol and Risk of Atherosclerosis. <i>Current Atherosclerosis Reports</i> , 2010, 12, 308-315.	4.8	71
180	PCSK9-deficient mice exhibit impaired glucose tolerance and pancreatic islet abnormalities. <i>FEBS Letters</i> , 2010, 584, 701-706.	2.8	165

#	ARTICLE	IF	CITATIONS
181	Molecular cloning and embryonic expression of zebrafish PCSK5 orthologues: Functional assessment during lateral line development. <i>Developmental Dynamics</i> , 2010, 239, 2933-2946.	1.8	16
182	A new method for measurement of total plasma PCSK9: clinical applications. <i>Journal of Lipid Research</i> , 2010, 51, 140-149.	4.2	197
183	Strong induction of PCSK9 gene expression through HNF1 $\alpha$ and SREBP2: mechanism for the resistance to LDL-cholesterol lowering effect of statins in dyslipidemic hamsters. <i>Journal of Lipid Research</i> , 2010, 51, 1486-1495.	4.2	208
184	ACAT1 gene ablation increases 24(S)-hydroxycholesterol content in the brain and ameliorates amyloid pathology in mice with AD. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2010, 107, 3081-3086.	7.1	170
185	Effects of the Prosegment and pH on the Activity of PCSK9. <i>Journal of Biological Chemistry</i> , 2010, 285, 40965-40978.	3.4	60
186	Intestinal and Hepatic Cholesterol Carriers in Diabetic <i>Psammomys obesus</i> . <i>Endocrinology</i> , 2010, 151, 958-970.	2.8	26
187	Surface Expression of Precursor N-cadherin Promotes Tumor Cell Invasion. <i>Neoplasia</i> , 2010, 12, 1066-1073.	5.3	56
188	Hypocholesterolaemia in chronic Hepatitis C virus (HCV) infection – Reduced production or increased LDL clearance?. <i>Atherosclerosis</i> , 2010, 213, e4.	0.8	0
189	A Locked Nucleic Acid Antisense Oligonucleotide (LNA) Silences PCSK9 and Enhances LDLR Expression In Vitro and In Vivo. <i>PLoS ONE</i> , 2010, 5, e10682.	2.5	167
190	Zebrafish ProVEGF-C Expression, Proteolytic Processing and Inhibitory Effect of Unprocessed ProVEGF-C during Fin Regeneration. <i>PLoS ONE</i> , 2010, 5, e11438.	2.5	20
191	Alzheimer Disease A $\beta$ Production in the Absence of S-Palmitoylation-dependent Targeting of BACE1 to Lipid Rafts. <i>Journal of Biological Chemistry</i> , 2009, 284, 3793-3803.	3.4	137
192	Potential Role of Proprotein Convertase SKI-1 in the Mineralization of Primary Bone. <i>Cells Tissues Organs</i> , 2009, 189, 25-32.	2.3	21
193	Regulation of the proprotein convertase subtilisin/kexin type 9 in intestinal epithelial cells. <i>American Journal of Physiology - Renal Physiology</i> , 2009, 296, G805-G815.	3.4	26
194	Dissection of the Endogenous Cellular Pathways of PCSK9-induced Low Density Lipoprotein Receptor Degradation. <i>Journal of Biological Chemistry</i> , 2009, 284, 28856-28864.	3.4	228
195	Genetic Variation at the Proprotein Convertase Subtilisin/Kexin Type 5 Gene Modulates High-Density Lipoprotein Cholesterol Levels. <i>Circulation: Cardiovascular Genetics</i> , 2009, 2, 467-475.	5.1	33
196	PCSK9 impedes hepatitis C virus infection <i>in vitro</i> and modulates liver CD81 expression. <i>Hepatology</i> , 2009, 50, 17-24.	7.3	129
197	The proprotein convertase PC5/6 is protective against intestinal tumorigenesis: in vivo mouse model. <i>Molecular Cancer</i> , 2009, 8, 73.	19.2	29
198	PCSK9 as a therapeutic target of dyslipidemia. <i>Expert Opinion on Therapeutic Targets</i> , 2009, 13, 19-28.	3.4	110

#	ARTICLE	IF	CITATIONS
199	Targeted production of proprotein convertase PC1 enhances mammary development and tumorigenesis in transgenic mice. This article is one of a selection of papers published in a special issue celebrating the 125th anniversary of the Faculty of Medicine at the University of Manitoba.. Canadian Journal of Physiology and Pharmacology, 2009, 87, 831-838.	1.4	4
200	The H5N1 Influenza Variant Fujian-Like Hemagglutinin Selected Following Vaccination Exhibits a Compromised Furin Cleavage. Journal of Molecular Neuroscience, 2008, 35, 339-343.	2.3	6
201	Proprotein convertase subtilisin/kexin type 9 (PCSK9): Hepatocyte-specific low-density lipoprotein receptor degradation and critical role in mouse liver regeneration. Hepatology, 2008, 48, 646-654.	7.3	354
202	PCSK9 is phosphorylated by a Golgi casein kinase-like kinase <i>in vivo</i> and circulates as a phosphoprotein in humans. FEBS Journal, 2008, 275, 3480-3493.	4.7	58
203	Cell-surface processing of extracellular human immunodeficiency virus type 1 Vpr by proprotein convertases. Virology, 2008, 372, 384-397.	2.4	38
204	The activation and physiological functions of the proprotein convertases. International Journal of Biochemistry and Cell Biology, 2008, 40, 1111-1125.	2.8	285
205	Proprotein convertases as therapeutic targets. Expert Opinion on Therapeutic Targets, 2008, 12, 1289-1300.	3.4	32
206	The Proprotein Convertase PCSK9 Induces the Degradation of Low Density Lipoprotein Receptor (LDLR) and Its Closest Family Members VLDLR and ApoER2. Journal of Biological Chemistry, 2008, 283, 2363-2372.	3.4	402
207	Inhibition of Chikungunya Virus Infection in Cultured Human Muscle Cells by Furin Inhibitors. Journal of Biological Chemistry, 2008, 283, 21899-21908.	3.4	114
208	<i>In vivo</i> functions of the proprotein convertase PC5/6 during mouse development: Gdf11 is a likely substrate. Proceedings of the National Academy of Sciences of the United States of America, 2008, 105, 5750-5755.	7.1	99
209	Annexin A2 Is a C-terminal PCSK9-binding Protein That Regulates Endogenous Low Density Lipoprotein Receptor Levels. Journal of Biological Chemistry, 2008, 283, 31791-31801.	3.4	132
210	The Regulated Cell Surface Zymogen Activation of the Proprotein Convertase PC5A Directs the Processing of Its Secretory Substrates. Journal of Biological Chemistry, 2008, 283, 2373-2384.	3.4	80
211	VACTERL/caudal regression/Currarino syndrome-like malformations in mice with mutation in the proprotein convertase <i>Pcsk5</i> . Genes and Development, 2008, 22, 1465-1477.	5.9	110
212	Selective inhibition of proprotein convertases represses the metastatic potential of human colorectal tumor cells. Journal of Clinical Investigation, 2008, 118, 352-363.	8.2	109
213	Role and regulation of the proprotein convertase subtilisin/kexin type 9 (PCSK9) in the small intestine. FASEB Journal, 2008, 22, 948.4.	0.5	0
214	Respiratory Distress and Neonatal Lethality in Mice Lacking Golgi $\beta$ 1,2-Mannosidase IB Involved in N-Glycan Maturation. Journal of Biological Chemistry, 2007, 282, 2558-2566.	3.4	31
215	The Proprotein Convertase SKI-1/S1P. Journal of Biological Chemistry, 2007, 282, 27402-27413.	3.4	30
216	Opposing Function of the Proprotein Convertases Furin and PACE4 on Breast Cancer Cells' Malignant Phenotypes: Role of Tissue Inhibitors of Metalloproteinase-1. Cancer Research, 2007, 67, 9030-9034.	0.9	57

#	ARTICLE	IF	CITATIONS
217	BEING PROUD OF ONESELF AS A PERSON OR BEING PROUD OF ONE'S PHYSICAL APPEARANCE: WHAT MATTERS FOR FEELING WELL IN ADOLESCENCE?. <i>Social Behavior and Personality</i> , 2007, 35, 255-268.	0.6	13
218	A targeted deletion/insertion in the mouse <i>Pcsk1</i> locus is associated with homozygous embryo preimplantation lethality, mutant allele preferential transmission and heterozygous female susceptibility to dietary fat. <i>Developmental Biology</i> , 2007, 306, 584-598.	2.0	34
219	Dysregulation of dynorphins in Alzheimer disease. <i>Neurobiology of Aging</i> , 2007, 28, 1700-1708.	3.1	60
220	Immunohistochemical expression and colocalization of somatostatin, carboxypeptidase-E and prohormone convertases 1 and 2 in rat brain. <i>Neuroscience</i> , 2007, 147, 403-418.	2.3	22
221	Plasma PCSK9 levels correlate with cholesterol in men but not in women. <i>Biochemical and Biophysical Research Communications</i> , 2007, 361, 451-456.	2.1	82
222	Of PCSK9, cholesterol homeostasis and parasitic infections: Possible survival benefits of loss-of-function PCSK9 genetic polymorphisms. <i>Medical Hypotheses</i> , 2007, 69, 1010-1017.	1.5	24
223	Heparin enhances the furin cleavage of HIV-1 gp160 peptides. <i>FEBS Letters</i> , 2007, 581, 5807-5813.	2.8	29
224	Gene Set Enrichment Analysis Reveals Several Globally Affected Pathways due to SKI-1/S1P Inhibition in HepG2 Cells. <i>DNA and Cell Biology</i> , 2007, 26, 765-772.	1.9	11
225	Three common alleles of KIR2DL4 (CD158d) encode constitutively expressed, inducible and secreted receptors in NK cells. <i>European Journal of Immunology</i> , 2007, 37, 199-211.	2.9	60
226	The Cellular Trafficking of the Secretory Proprotein Convertase PCSK9 and Its Dependence on the LDLR. <i>Traffic</i> , 2007, 8, 718-732.	2.7	213
227	PC1/3, PC2 and PC5/6A are targeted to dense core secretory granules by a common mechanism. <i>FEBS Journal</i> , 2007, 274, 4094-4102.	4.7	22
228	The proprotein convertases are potential targets in the treatment of dyslipidemia. <i>Journal of Molecular Medicine</i> , 2007, 85, 685-696.	3.9	145
229	Increased stress-induced analgesia in mice lacking the proneuropeptide convertase PC2. <i>Neuroscience Letters</i> , 2006, 406, 71-75.	2.1	19
230	Unexpected similarity between the cytosolic West Nile virus NS3 and the secretory furin-like serine proteinases. <i>Biochemical Journal</i> , 2006, 393, e1-3.	3.7	11
231	Implication of the proprotein convertase NARC-1/PCSK9 in the development of the nervous system. <i>Journal of Neurochemistry</i> , 2006, 98, 838-850.	3.9	99
232	Expression and Processing of the Neuroendocrine Protein Secretogranin II in Benign and Malignant Pheochromocytomas. <i>Annals of the New York Academy of Sciences</i> , 2006, 1073, 527-532.	3.8	19
233	The Proprotein Convertase SKI-1/S1P. <i>Journal of Biological Chemistry</i> , 2006, 281, 23471-23481.	3.4	57
234	The proprotein convertases and their implication in sterol and/or lipid metabolism. <i>Biological Chemistry</i> , 2006, 387, 871-7.	2.5	88

#	ARTICLE	IF	CITATIONS
235	Deletion of the Gene Encoding Proprotein Convertase 5/6 Causes Early Embryonic Lethality in the Mouse. <i>Molecular and Cellular Biology</i> , 2006, 26, 354-361.	2.3	73
236	Proprotein convertases: lessons from knockouts. <i>FASEB Journal</i> , 2006, 20, 1954-1963.	0.5	210
237	The Proprotein Convertase (PC) PCSK9 Is Inactivated by Furin and/or PC5/6A. <i>Journal of Biological Chemistry</i> , 2006, 281, 30561-30572.	3.4	246
238	Evaluation of Anti-Proprotein Convertase Activity of Diterpene Andrographolid Derived Products. , 2006, , 137-154.		5
239	Discovery of the Proprotein Convertases and their Inhibitors. , 2006, , 7-26.		0
240	beta-Amyloid protein converting enzyme 1 and brain-specific type II membrane protein BRI3: binding partners processed by furin. <i>Journal of Neurochemistry</i> , 2005, 92, 93-102.	3.9	39
241	Regulation of the stepwise proteolytic cleavage and secretion of PDGF-B by the proprotein convertases. <i>Oncogene</i> , 2005, 24, 6925-6935.	5.9	67
242	Translational control of hippocampal synaptic plasticity and memory by the eIF2 $\gamma$ kinase GCN2. <i>Nature</i> , 2005, 436, 1166-1170.	27.8	344
243	Abnormal expression and processing of the proprotein convertases PC1 and PC2 in human colorectal liver metastases. <i>BMC Cancer</i> , 2005, 5, 149.	2.6	26
244	Immunohistochemical localization of subtilisin/kexin-like proprotein convertases in human atherosclerosis. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2005, 446, 351-359.	2.8	28
245	Human Carcinoma Cell Growth and Invasiveness Is Impaired by the Propeptide of the Ubiquitous Proprotein Convertase Furin. <i>Cancer Research</i> , 2005, 65, 4162-4171.	0.9	47
246	Proprotein Convertases Are Responsible for Proteolysis and Inactivation of Endothelial Lipase. <i>Journal of Biological Chemistry</i> , 2005, 280, 36551-36559.	3.4	91
247	The Cysteine-rich Domain of the Secreted Proprotein Convertases PC5A and PACE4 Functions as a Cell Surface Anchor and Interacts with Tissue Inhibitors of Metalloproteinases. <i>Molecular Biology of the Cell</i> , 2005, 16, 5215-5226.	2.1	88
248	Wild-type PCSK9 inhibits LDL clearance but does not affect apoB-containing lipoprotein production in mouse and cultured cells. <i>Journal of Lipid Research</i> , 2005, 46, 1312-1319.	4.2	93
249	Furin-Like Proprotein Convertases Are Central Regulators of the Membrane Type Matrix Metalloproteinase $\alpha$ 2-Pro-Matrix Metalloproteinase-2 Proteolytic Cascade in Atherosclerosis. <i>Circulation</i> , 2005, 111, 2820-2827.	1.6	103
250	Implication of proprotein convertases in the processing and spread of severe acute respiratory syndrome coronavirus. <i>Biochemical and Biophysical Research Communications</i> , 2005, 326, 554-563.	2.1	71
251	Dual regulation of the LDL receptor $\alpha$ 2—Some clarity and new questions. <i>Cell Metabolism</i> , 2005, 1, 290-292.	16.2	87
252	Evidence for proprotein convertase activity in the endoplasmic reticulum/early Golgi. <i>FEBS Letters</i> , 2005, 579, 5621-5625.	2.8	26

#	ARTICLE	IF	CITATIONS
253	Chloroquine is a potent inhibitor of SARS coronavirus infection and spread. <i>Virology Journal</i> , 2005, 2, 69.	3.4	1,457
254	Statins Upregulate PCSK9, the Gene Encoding the Proprotein Convertase Neural Apoptosis-Regulated Convertase-1 Implicated in Familial Hypercholesterolemia. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2004, 24, 1454-1459.	2.4	557
255	Endothelium-Restricted Overexpression of Human Endothelin-1 Causes Vascular Remodeling and Endothelial Dysfunction. <i>Circulation</i> , 2004, 110, 2233-2240.	1.6	296
256	Development of Protein-based Inhibitors of the Proprotein of Convertase SKI-1/S1P. <i>Journal of Biological Chemistry</i> , 2004, 279, 17338-17347.	3.4	51
257	NARC-1/PCSK9 and Its Natural Mutants. <i>Journal of Biological Chemistry</i> , 2004, 279, 48865-48875.	3.4	544
258	Endoproteolytic Activation of $\alpha_5\beta_1$ Integrin by Proprotein Convertase PC5 Is Required for Vascular Smooth Muscle Cell Adhesion to Vitronectin and Integrin-Dependent Signaling. <i>Circulation</i> , 2004, 109, 770-776.	1.6	32
259	Regulation of matrix metalloproteinase MT1-MMP/MMP-2 in cardiac fibroblasts by TGF- $\beta_1$ involves furin-convertase. <i>Cardiovascular Research</i> , 2004, 63, 87-97.	3.8	100
260	Stepwise Posttranslational Processing of Progrowth Hormone-Releasing Hormone (proGHRH) Polypeptide by Furin and PC1. <i>Endocrine</i> , 2004, 23, 199-214.	2.2	24
261	Implications of proprotein Convertase 5 (PC5) in the arterial restenotic process in a porcine model. <i>Cardiovascular Pathology</i> , 2004, 13, 241-250.	1.6	8
262	Induction of immune activation by a novel immunomodulatory oligonucleotide without thymocyte apoptosis. <i>Biochemical and Biophysical Research Communications</i> , 2004, 318, 60-66.	2.1	8
263	Proprotein convertases regulate insulin-like growth factor 1-induced membrane-type 1 matrix metalloproteinase in VSMCs via endoproteolytic activation of the insulin-like growth factor-1 receptor. <i>Biochemical and Biophysical Research Communications</i> , 2004, 321, 531-538.	2.1	35
264	The metabolism of $\beta$ -amyloid converting enzyme and $\beta$ -amyloid precursor protein processing. <i>Biochemical and Biophysical Research Communications</i> , 2004, 325, 235-242.	2.1	10
265	Prosomatostatin is proteolytically processed at the amino terminal segment by subtilase SKI-1. <i>Regulatory Peptides</i> , 2004, 120, 133-140.	1.9	23
266	Inhibition of Proprotein Convertases Enhances Cell Migration and Metastases Development of Human Colon Carcinoma Cells in a Rat Model. <i>American Journal of Pathology</i> , 2004, 164, 1925-1933.	3.8	27
267	Coordinated regulation and colocalization of $\alpha_5\beta_1$ integrin and its activating enzyme proprotein convertase PC5 in vivo. <i>Histochemistry and Cell Biology</i> , 2003, 119, 239-245.	1.7	18
268	Mutations in PCSK9 cause autosomal dominant hypercholesterolemia. <i>Nature Genetics</i> , 2003, 34, 154-156.	21.4	2,532
269	Processing of $\alpha_4$ integrin by the proprotein convertases: histidine at position P6 regulates cleavage. <i>Biochemical Journal</i> , 2003, 373, 475-484.	3.7	56
270	The proprotein convertase PC2 is involved in the maturation of prosomatostatin to somatostatin-14 but not in the somatostatin deficit in Alzheimer's disease. <i>Neuroscience</i> , 2003, 122, 437-447.	2.3	31



#	ARTICLE	IF	CITATIONS
271	Proteolytic processing of chromogranin A by the prohormone convertase PC2. <i>Regulatory Peptides</i> , 2003, 111, 111-116.	1.9	18
272	Crimean-Congo Hemorrhagic Fever Virus Glycoprotein Proteolytic Processing by Subtilase SKI-1. <i>Journal of Virology</i> , 2003, 77, 8640-8649.	3.4	132
273	A Serine Protease Inhibitor Prevents Endoplasmic Reticulum Stress-induced Cleavage but Not Transport of the Membrane-bound Transcription Factor ATF6. <i>Journal of Biological Chemistry</i> , 2003, 278, 31024-31032.	3.4	194
274	Neurotrophin-4, Alone or Heterodimerized with Brain-derived Neurotrophic Factor, Is Sorted to the Constitutive Secretory Pathway. <i>Journal of Biological Chemistry</i> , 2003, 278, 48129-48136.	3.4	22
275	The secretory proprotein convertase neural apoptosis-regulated convertase 1 (NARC-1): Liver regeneration and neuronal differentiation. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2003, 100, 928-933.	7.1	1,012
276	Structure-Function Analysis of the Prosegment of the Proprotein Convertase PC5A. <i>Journal of Biological Chemistry</i> , 2003, 278, 2886-2895.	3.4	46
277	The Proprotein Convertase PC5A and a Metalloprotease Are Involved in the Proteolytic Processing of the Neural Adhesion Molecule L1. <i>Journal of Biological Chemistry</i> , 2003, 278, 10381-10388.	3.4	90
278	Down-regulation of alphaV/beta3 integrin via misrouting to lysosomes by overexpression of a beta3Lamp1 fusion protein. <i>Biochemical Journal</i> , 2003, 370, 703-711.	3.7	12
279	The secretory proprotein convertases furin, PC5, and PC7 activate VEGF-C to induce tumorigenesis. <i>Journal of Clinical Investigation</i> , 2003, 111, 1723-1732.	8.2	109
280	The secretory proprotein convertases furin, PC5, and PC7 activate VEGF-C to induce tumorigenesis. <i>Journal of Clinical Investigation</i> , 2003, 111, 1723-1732.	8.2	170
281	The proteolytic processing of pro-platelet-derived growth factor-A at RRKR(86) by members of the proprotein convertase family is functionally correlated to platelet-derived growth factor-A-induced functions and tumorigenicity. <i>Cancer Research</i> , 2003, 63, 1458-63.	0.9	64
282	Genetic Mapping of the Gene for SKI-1/S1P Protease (locus Symbol Mbtps 1) to Mouse Chromosome 8. <i>DNA Sequence</i> , 2002, 13, 109-111.	0.7	4
283	Biosynthesis and Cellular Trafficking of the Convertase SKI-1/S1P. <i>Journal of Biological Chemistry</i> , 2002, 277, 11265-11275.	3.4	100
284	The metalloendopeptidase nardilysin (NRDc) is potently inhibited by heparin-binding epidermal growth factor-like growth factor (HB-EGF). <i>Biochemical Journal</i> , 2002, 367, 229-238.	3.7	24
285	Effects of l- and d-REKR amino acid-containing peptides on HIV and SIV envelope glycoprotein precursor maturation and HIV and SIV replication. <i>Biochemical Journal</i> , 2002, 366, 863-872.	3.7	4
286	La mesure du perfectionnisme: Validation canadienne-française du Positive and Negative Perfectionism Scale.. <i>Canadian Journal of Behavioural Science</i> , 2002, 34, 168-171.	0.6	12
287	Mesure de l'estime de soi à l'adolescence: version canadienne-française du Self-Perception Profile for Adolescents de Harter.. <i>Canadian Journal of Behavioural Science</i> , 2002, 34, 158-162.	0.6	25
288	Role of proprotein convertases in the pathogenic processing of the amyloidosis-associated form of secretory gelsolin. <i>Amyloid: the International Journal of Experimental and Clinical Investigation: the Official Journal of the International Society of Amyloidosis</i> , 2002, 9, 83-87.	3.0	15

#	ARTICLE	IF	CITATIONS
289	Cellular Limited Proteolysis of Precursor Proteins and Peptides. <i>The Enzymes</i> , 2002, 22, 237-258.	1.7	4
290	Proprotein Convertases in Tumor Progression and Malignancy. <i>American Journal of Pathology</i> , 2002, 160, 1921-1935.	3.8	196
291	A rapid fluorometric assay for the proteolytic activity of SKI-1/S1P based on the surface glycoprotein of the hemorrhagic fever Lassa virus. <i>FEBS Letters</i> , 2002, 514, 333-339.	2.8	34
292	Characterization of a repressor element in the promoter region of proprotein convertase 2 (PC2) gene. <i>Molecular Brain Research</i> , 2002, 102, 35-47.	2.3	4
293	Altered processing of the neurotensin/neuromedinâ€fN precursor in PC2 knock down mice: a biochemical and immunohistochemical study. <i>Journal of Neurochemistry</i> , 2002, 82, 783-793.	3.9	47
294	Proteolytic Processing of Chromogranin B and Secretogranin II by Prohormone Convertases. <i>Journal of Neurochemistry</i> , 2002, 70, 374-383.	3.9	58
295	Synthetic proregion-related peptides of proprotein convertases, PC1 and furin, represent potent inhibitors of each protease. , 2002, , 676-678.		1
296	Precursor convertases in the secretory pathway, cytosol and extracellular milieu. <i>Essays in Biochemistry</i> , 2002, 38, 79-94.	4.7	190
297	Proprotein convertases are important mediators of the adipocyte differentiation of mouse 3T3-L1 cells. <i>Journal of Cell Science</i> , 2002, 115, 1203-1211.	2.0	19
298	Biosynthesis and enzymatic characterization of human SKI-1 and the processing of its inhibitory prosegment. , 2002, , 465-466.		0
299	Evidence that Furin Is an Authentic Transforming Growth Factor-Î²1-Converting Enzyme. <i>American Journal of Pathology</i> , 2001, 158, 305-316.	3.8	220
300	Biosynthesis and Post-translational Processing of the Precursor to Brain-derived Neurotrophic Factor. <i>Journal of Biological Chemistry</i> , 2001, 276, 12660-12666.	3.4	480
301	Implication of the proprotein convertases furin, PC5 and PC7 in the cleavage of surface glycoproteins of Hong Kong, Ebola and respiratory syncytial viruses: a comparative analysis with fluorogenic peptides. <i>Biochemical Journal</i> , 2001, 353, 537.	3.7	68
302	Neuroendocrine secretory protein 7B2: structure, expression and functions. <i>Biochemical Journal</i> , 2001, 357, 329.	3.7	123
303	Replication of HIV-1 viruses in the presence of the Portland Î±1-antitrypsin variant (Î±1-PDX) inhibitor. <i>Biochemical Journal</i> , 2001, 360, 127.	3.7	4
304	Implication of the proprotein convertases furin, PC5 and PC7 in the cleavage of surface glycoproteins of Hong Kong, Ebola and respiratory syncytial viruses: a comparative analysis with fluorogenic peptides. <i>Biochemical Journal</i> , 2001, 353, 537-545.	3.7	87
305	Neuroendocrine secretory protein 7B2: structure, expression and functions. <i>Biochemical Journal</i> , 2001, 357, 329-342.	3.7	155
306	Constitutive Î±-secretase cleavage of the Î²-amyloid precursor protein in the furin-deficient LoVo cell line: involvement of the pro-hormone convertase 7 and the disintegrin metalloprotease ADAM10. <i>Journal of Neurochemistry</i> , 2001, 76, 1532-1539.	3.9	113

#	ARTICLE	IF	CITATIONS
307	Inhibition of HIV-2ROD replication in a lymphoblastoid cell line by the $\hat{I}\pm 1$ -antitrypsin Portland variant ( $\hat{I}\pm 1$ -PDX) and the decRVKRcmk peptide: comparison with HIV-1LAI. <i>Microbes and Infection</i> , 2001, 3, 1073-1084.	1.9	3
308	pH-induced conformational transitions of a molten-globule-like state of the inhibitory prodomain of furin: Implications for zymogen activation. <i>Protein Science</i> , 2001, 10, 934-942.	7.6	26
309	The Lassa virus glycoprotein precursor GP-C is proteolytically processed by subtilase SKI-1/S1P. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2001, 98, 12701-12705.	7.1	316
310	Selective Expression of the Proprotein Convertases Furin, PC5, and PC7 in Proliferating Vascular Smooth Muscle Cells of the Rat Aorta In Vitro. <i>Journal of Histochemistry and Cytochemistry</i> , 2001, 49, 323-331.	2.5	28
311	Inhibition of Proprotein Convertases Is Associated with Loss of Growth and Tumorigenicity of HT-29 Human Colon Carcinoma Cells. <i>Journal of Biological Chemistry</i> , 2001, 276, 30686-30693.	3.4	156
312	Inhibitory Specificity and Potency of proSAAS-derived Peptides toward Proprotein Convertase 1. <i>Journal of Biological Chemistry</i> , 2001, 276, 32720-32728.	3.4	52
313	Post-translational Processing of $\hat{I}^2$ -Secretase ( $\hat{I}^2$ -Amyloid-converting Enzyme) and Its Ectodomain Shedding. <i>Journal of Biological Chemistry</i> , 2001, 276, 10879-10887.	3.4	273
314	Elevated expression of proprotein convertases alters breast cancer cell growth in response to estrogen and tamoxifen. <i>Journal of Molecular Endocrinology</i> , 2001, 26, 95-105.	2.5	23
315	PreproTRH178-199 and Two Novel Peptides (pFQ7 and pSE14) Derived from Its Processing, Which Are Produced in the Paraventricular Nucleus of the Rat Hypothalamus, Are Regulated during Suckling**This work was supported by the National Science Foundation (Grant No. IBN-9507952 to) Tj ETQq1 1 0.784314 rgBT /Over	2.8	38
316	A Highly Selective Cell Permeable Peptide Inhibitor of Proprotein Convertase 1: Design, Synthesis and Biological Evaluation in Cellular PC1-Mediated Proteolysis. , 2001, , 558-560.		1
317	PreproTRH178-199 and Two Novel Peptides (pFQ7 and pSE14) Derived from Its Processing, Which Are Produced in the Paraventricular Nucleus of the Rat Hypothalamus, Are Regulated during Suckling. <i>Endocrinology</i> , 2001, 142, 896-906.	2.8	15
318	Biosynthesis and Enzymatic Characterization of Human SKI-1/S1P and the Processing of Its Inhibitory Prosegment. <i>Journal of Biological Chemistry</i> , 2000, 275, 2349-2358.	3.4	79
319	Endoproteolytic processing of integrin pro- $\hat{I}\pm$ subunits involves the redundant function of furin and proprotein convertase (PC) 5A, but not paired basic amino acid converting enzyme (PACE) 4, PC5B or PC7. <i>Biochemical Journal</i> , 2000, 346, 133.	3.7	32
320	Molecular cloning and biochemical characterization of a new mouse testis soluble-zinc-metallopeptidase of the neprilysin family. <i>Biochemical Journal</i> , 2000, 347, 419.	3.7	49
321	N-arginine dibasic convertase (nardilysin) isoforms are soluble dibasic-specific metalloendopeptidases that localize in the cytoplasm and at the cell surface. <i>Biochemical Journal</i> , 2000, 349, 587.	3.7	29
322	N-arginine dibasic convertase (nardilysin) isoforms are soluble dibasic-specific metalloendopeptidases that localize in the cytoplasm and at the cell surface. <i>Biochemical Journal</i> , 2000, 349, 587-597.	3.7	34
323	Effect of alpha-1 antitrypsin Portland variant ( $\hat{I}\pm 1$ -PDX) on HIV-1 replication. <i>Biochemical Journal</i> , 2000, 352, 91.	3.7	3
324	Immunohistochemical evidence for the implication of PCI in the processing of proneurotensin in rat brain. <i>NeuroReport</i> , 2000, 11, 3443-3447.	1.2	18

#	ARTICLE	IF	CITATIONS
325	Regional and cellular localization of the neuroendocrine prohormone convertases PC1 and PC2 in the rat central nervous system. <i>Journal of Comparative Neurology</i> , 2000, 424, 439-460.	1.6	49
326	Immunohistochemical evidence for the involvement of protein convertases 5A and 2 in the processing of pro-neurotensin in rat brain. <i>Journal of Comparative Neurology</i> , 2000, 424, 461-475.	1.6	29
327	Pro-opiomelanocortin-Related Peptides, Prohormone Convertases 1 and 2 and the Regulatory Peptide 7B2 are Present in Melanosomes of Human Melanocytes. <i>Journal of Investigative Dermatology</i> , 2000, 114, 430-437.	0.7	52
328	Neurotrophin-3 Sorts to the Constitutive Secretory Pathway of Hippocampal Neurons and Is Diverted to the Regulated Secretory Pathway by Coexpression with Brain-Derived Neurotrophic Factor. <i>Journal of Neuroscience</i> , 2000, 20, 4059-4068.	3.6	114
329	Molecular Cloning Demonstrates Structural Features of Homologous Bovine Prohormone Convertases 1 and 2. <i>DNA and Cell Biology</i> , 2000, 19, 409-419.	1.9	2
330	Alternative Proteolytic Processing of Mouse Mammary Tumor Virus Superantigens. <i>Journal of Virology</i> , 2000, 74, 3067-3073.	3.4	8
331	The Kex2p Proregion Is Essential for the Biosynthesis of an Active Enzyme and Requires a C-terminal Basic Residue for Its Function. <i>Molecular Biology of the Cell</i> , 2000, 11, 1947-1957.	2.1	26
332	A Predicted $\alpha$ -Helix Mediates Targeting of the Proprotein Convertase PC1 to the Regulated Secretory Pathway. <i>Journal of Biological Chemistry</i> , 2000, 275, 40337-40343.	3.4	48
333	The C-terminal Region of proSAAS Is a Potent Inhibitor of Prohormone Convertase 1. <i>Journal of Biological Chemistry</i> , 2000, 275, 23596-23601.	3.4	96
334	Cathepsin-B Fusion Proteins Misroute Secretory Protein Partners Such as the Proprotein Convertase PC2-7B2 Complex toward the Lysosomal Degradation Pathways. <i>Biochemical and Biophysical Research Communications</i> , 2000, 276, 594-599.	2.1	4
335	Inhibitory Activity and Structural Characterization of a C-Terminal Peptide Fragment Derived from the Prosegment of the Proprotein Convertase PC7. <i>Biochemistry</i> , 2000, 39, 2868-2877.	2.5	27
336	Coordinated Expression of $\beta$ -Amyloid Precursor Protein and the Putative $\beta$ -Secretase BACE and $\gamma$ -Secretase ADAM10 in Mouse and Human Brain. <i>Journal of Neurochemistry</i> , 2000, 75, 2133-2143.	3.9	160
337	The RGD Motif and the C-terminal Segment of Proprotein Convertase 1 Are Critical for Its Cellular Trafficking but Not for Its Intracellular Binding to Integrin $\alpha$ 5 $\beta$ 1. <i>Journal of Biological Chemistry</i> , 1999, 274, 12461-12467.	3.4	39
338	Differential Sorting of Nerve Growth Factor and Brain-Derived Neurotrophic Factor in Hippocampal Neurons. <i>Journal of Neuroscience</i> , 1999, 19, 2069-2080.	3.6	299
339	Parathyroid Hormone Processing by the Proprotein Convertase-7: Comparison with Furin and Assessment of Modulation of Parathyroid Convertase Messenger Ribonucleic Acid Levels by Calcium and 1,25-Dihydroxyvitamin D3*. <i>Endocrinology</i> , 1999, 140, 3633-3642.	2.8	34
340	The Prosegments of Furin and PC7 as Potent Inhibitors of Proprotein Convertases. <i>Journal of Biological Chemistry</i> , 1999, 274, 33913-33920.	3.4	122
341	Cellular Localization and Role of Prohormone Convertases in the Processing of Pro-melanin Concentrating Hormone in Mammals. <i>Journal of Biological Chemistry</i> , 1999, 274, 6536-6545.	3.4	53
342	Mammalian subtilisin/kexin isozyme SKI-1: A widely expressed proprotein convertase with a unique cleavage specificity and cellular localization. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1999, 96, 1321-1326.	7.1	273

#	ARTICLE	IF	CITATIONS
343	Distribution and regulation of proconvertases PC1 and PC2 in human pituitary adenomas. <i>Pituitary</i> , 1999, 1, 187-195.	2.9	23
344	Proprotein and prohormone convertases: a family of subtilases generating diverse bioactive polypeptides1Published on the World Wide Web on 17 August 1999.1. <i>Brain Research</i> , 1999, 848, 45-62.	2.2	775
345	Molecular cloning, characterization of cDNA, and distribution of mRNA encoding the frog prohormone convertase PC1. , 1999, 405, 160-172.		17
346	Immunohistochemical distribution of the prohormone convertase PC5-A in rat brain. <i>Neuroscience</i> , 1999, 92, 641-654.	2.3	22
347	Occurrence of an HIV-1 gp160 endoproteolytic activity in low-density vesicles and evidence for a distinct density distribution from endogenously expressed furin and PC7/LPC convertases. <i>FEBS Letters</i> , 1999, 456, 97-102.	2.8	17
348	Enzymic characterization in vitro of recombinant proprotein convertase PC4. <i>Biochemical Journal</i> , 1999, 343, 29-37.	3.7	41
349	Inhibition of proprotein convertases-1, -7 and furin by diterpines of <i>Andrographis paniculata</i> and their succinoyl esters. <i>Biochemical Journal</i> , 1999, 338, 107.	3.7	38
350	Inhibition of proprotein convertases-1, -7 and furin by diterpines of <i>Andrographis paniculata</i> and their succinoyl esters. <i>Biochemical Journal</i> , 1999, 338, 107-113.	3.7	102
351	The Subtilisin/Kexin Family of Precursor Convertases: Emphasis on PC1, PC2/7B2, POMC and the Novel Enzyme SKIâ€¹. <i>Annals of the New York Academy of Sciences</i> , 1999, 885, 57-74.	3.8	130
352	Enzymic characterization in vitro of recombinant proprotein convertase PC4. <i>Biochemical Journal</i> , 1999, 343, 29.	3.7	20
353	Parathyroid Hormone Processing by the Proprotein Convertase-7: Comparison with Furin and Assessment of Modulation of Parathyroid Convertase Messenger Ribonucleic Acid Levels by Calcium and 1,25-Dihydroxyvitamin D3. <i>Endocrinology</i> , 1999, 140, 3633-3642.	2.8	9
354	Precursor Convertases: An Evolutionary Ancient, Cell-Specific, Combinatorial Mechanism Yielding Diverse Bioactive Peptides and Proteins. <i>Annals of the New York Academy of Sciences</i> , 1998, 839, 9-24.	3.8	187
355	The LIM homeobox protein mLIM3/Lhx3 induces expression of the prolactin gene by a Pit-1/GHF-1-independent pathway in corticotroph AtT20 cells. <i>FEBS Letters</i> , 1998, 431, 333-338.	2.8	15
356	Residues unique to the pro-hormone convertase PC2 modulate its autoactivation, binding to 7B2 and enzymatic activity. <i>FEBS Letters</i> , 1998, 428, 37-42.	2.8	31
357	Molecular Characterization, Enzymatic Analysis, and Purification of Murine Proprotein Convertase-1/3 (PC1/PC3) Secreted from Recombinant Baculovirus-Infected Insect Cells. <i>Protein Expression and Purification</i> , 1998, 14, 353-366.	1.3	31
358	Immunocytochemical Localization of the Prohormone Convertases PC1 and PC2 in Rat Prolactin Cells. <i>Journal of Histochemistry and Cytochemistry</i> , 1998, 46, 101-108.	2.5	10
359	Attenuation of the Polypeptide 7B2, Prohormone Convertase PC2, and Vasopressin in the Hypothalamus of Some Prader-Willi Patients: Indications for a Processing Defect. <i>Journal of Clinical Endocrinology and Metabolism</i> , 1998, 83, 591-599.	3.6	11
360	PC5-A-mediated Processing of Pro-neurotensin in Early Compartments of the Regulated Secretory Pathway of PC5-transfected PC12 Cells. <i>Journal of Biological Chemistry</i> , 1998, 273, 25339-25346.	3.4	46

#	ARTICLE	IF	CITATIONS
361	In Vitro Cleavage of Internally Quenched Fluorogenic Human Parathyroid Hormone and Parathyroid-related Peptide Substrates by Furin. <i>Journal of Biological Chemistry</i> , 1998, 273, 8572-8580.	3.4	62
362	The Pore-forming Toxin Proaerolysin Is Activated by Furin. <i>Journal of Biological Chemistry</i> , 1998, 273, 32656-32661.	3.4	130
363	Delta-1 Activation of Notch-1 Signaling Results in $\Delta$ HES-1 Transactivation. <i>Molecular and Cellular Biology</i> , 1998, 18, 7423-7431.	2.3	301
364	The Notch1 receptor is cleaved constitutively by a furin-like convertase. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1998, 95, 8108-8112.	7.1	661
365	The Vasopressin Precursor Is Not Processed in the Hypothalamus of Wolfram Syndrome Patients with Diabetes Insipidus: Evidence for the Involvement of PC2 and 7B2. <i>Journal of Clinical Endocrinology and Metabolism</i> , 1998, 83, 4026-4033.	3.6	16
366	In Vitro Characterization of the Novel Proprotein Convertase PC7. <i>Journal of Biological Chemistry</i> , 1997, 272, 19672-19681.	3.4	81
367	Processing of Prothyrotropin-releasing Hormone by the Family of Prohormone Convertases. <i>Journal of Biological Chemistry</i> , 1997, 272, 19958-19968.	3.4	71
368	$\Delta$ 1-Antitrypsin Portland Inhibits Processing of Precursors Mediated by Proprotein Convertases Primarily within the Constitutive Secretory Pathway. <i>Journal of Biological Chemistry</i> , 1997, 272, 26210-26218.	3.4	88
369	Prohormone Convertases in Mouse Submandibular Gland: Co-localization of Furin and Nerve Growth Factor. <i>Journal of Histochemistry and Cytochemistry</i> , 1997, 45, 795-804.	2.5	19
370	The integrity of the RRGDL sequence of the proprotein convertase PC1 is critical for its zymogen and C-terminal processing and for its cellular trafficking. <i>Biochemical Journal</i> , 1997, 326, 737-744.	3.7	48
371	Impaired fertility in mice deficient for the testicular germ-cell protease PC4. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1997, 94, 6842-6846.	7.1	156
372	The Developmental Expression in the Rat CNS and Peripheral Tissues of Proteases PC5 and PACE4 mRNAs: Comparison with Other Proprotein Processing Enzymes. <i>Developmental Biology</i> , 1997, 181, 268-283.	2.0	43
373	Comparative functional role of PC7 and furin in the processing of the HIV envelope glycoprotein gp160. <i>FEBS Letters</i> , 1997, 405, 68-72.	2.8	54
374	Comparative processing of bovine leukemia virus envelope glycoprotein gp72 by subtilisin/kexin-like mammalian convertases. <i>FEBS Letters</i> , 1997, 406, 205-210.	2.8	23
375	Immunohistochemical Localization of the Pro-Peptide Processing Enzymes PC1/PC3 and PC2 in the Human Anal Canal. <i>Peptides</i> , 1997, 18, 755-760.	2.4	7
376	Pilocarpine-induced seizures are accompanied by a transient elevation in the messenger RNA expression of the prohormone convertase PC1 in rat hippocampus: Comparison with nerve growth factor and brain-derived neurotrophic factor expression. <i>Neuroscience</i> , 1997, 76, 425-439.	2.3	27
377	Two Activation States of the Prohormone Convertase PC1 in the Secretory Pathway. <i>Journal of Biological Chemistry</i> , 1997, 272, 15184-15188.	3.4	65
378	Cellular Localization of the Prohormone Convertases in the Hypothalamic Paraventricular and Supraoptic Nuclei: Selective Regulation of PC1 in Corticotrophin-Releasing Hormone Parvocellular Neurons Mediated by Glucocorticoids. <i>Journal of Neuroscience</i> , 1997, 17, 563-575.	3.6	93

#	ARTICLE	IF	CITATIONS
379	Regulation by gastric acid of the processing of progastrin-derived peptides in rat antral mucosa. <i>Journal of Physiology</i> , 1997, 502, 409-419.	2.9	190
380	Comparative analysis of expression of the proprotein convertases furin, PACE4, PC1 and PC2 in human lung tumours. <i>British Journal of Cancer</i> , 1997, 75, 1509-1514.	6.4	116
381	Prorenin activation and prohormone convertases in the mouse As4.1 cell line. <i>Kidney International</i> , 1997, 51, 104-109.	5.2	21
382	Expression of the proprotein convertases PC1 and PC2 mRNAs in thyrotropin releasing hormone neurons of the rat paraventricular nucleus of hypothalamus. <i>Brain Research</i> , 1997, 761, 77-86.	2.2	22
383	Eukaryotic protein processing: endoproteolysis of precursor proteins. <i>Current Opinion in Biotechnology</i> , 1997, 8, 602-607.	6.6	271
384	Pro-protein convertase gene expression in human breast cancer. , 1997, 71, 966-971.		135
385	Histidine-rich human salivary peptides are inhibitors of proprotein convertases furin and PC7 but act as substrates for PC1. <i>Chemical Biology and Drug Design</i> , 1997, 49, 596-603.	1.1	26
386	Pro-protein convertase gene expression in human breast cancer. <i>International Journal of Cancer</i> , 1997, 71, 966-971.	5.1	2
387	Increased Proteolytic Processing of Protein Tyrosine Phosphatase $\frac{1}{4}$ in Confluent Vascular Endothelial Cells: The Role of PC5, a Member of the Subtilisin Family. <i>Biochemistry</i> , 1996, 35, 3797-3802.	2.5	64
388	Cellular processing of the neurotrophin precursors of NT3 and BDNF by the mammalian proprotein convertases. <i>FEBS Letters</i> , 1996, 379, 247-250.	2.8	309
389	Functional analysis of human PACE4-A and PACE4-C isoforms: identification of a new PACE4-CS isoform. <i>FEBS Letters</i> , 1996, 396, 31-36.	2.8	25
390	cDNA structure, tissue distribution, and chromosomal localization of rat PC7, a novel mammalian proprotein convertase closest to yeast kexin-like proteinases.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1996, 93, 3388-3393.	7.1	250
391	Comparative cellular processing of the human immunodeficiency virus (HIV-1) envelope glycoprotein gp160 by the mammalian subtilisin/kexin-like convertases. <i>Biochemical Journal</i> , 1996, 314, 521-532.	3.7	105
392	Cellular processing of the nerve growth factor precursor by the mammalian pro-protein convertases. <i>Biochemical Journal</i> , 1996, 314, 951-960.	3.7	258
393	Lack of integrin $\beta$ -chain endoproteolytic cleavage in furin-deficient human colon adenocarcinoma cells LoVo. <i>Biochemical Journal</i> , 1996, 317, 803-809.	3.7	60
394	Characterization of structural determinants and molecular mechanisms involved in pro-stromelysin-3 activation by 4-aminophenylmercuric acetate and furin-type convertases. <i>Biochemical Journal</i> , 1996, 315, 953-958.	3.7	73
395	The isoforms of proprotein convertase PC5 are sorted to different subcellular compartments.. <i>Journal of Cell Biology</i> , 1996, 135, 1261-1275.	5.2	152
396	Identification of the Paired Basic Convertases Implicated in HIV gp160 Processing Based on in Vitro Assays and Expression in CD4+ Cell Lines. <i>Journal of Biological Chemistry</i> , 1996, 271, 30442-30450.	3.4	109

#	ARTICLE	IF	CITATIONS
397	Prohormone Convertase PC5 Is a Candidate Processing Enzyme for Prorenin in the Human Adrenal Cortex. <i>Hypertension</i> , 1996, 28, 840-846.	2.7	34
398	Neural and glial-mediated effects of growth factors acting via tyrosine kinase receptors on luteinizing hormone-releasing hormone neurons. <i>Endocrinology</i> , 1996, 137, 2593-2605.	2.8	22
399	Role of prohormone convertases in the tissue-specific processing of proglucagon. <i>Molecular Endocrinology</i> , 1996, 10, 342-355.	3.7	127
400	[1] Molecular strategies for identifying processing enzymes. <i>Methods in Neurosciences</i> , 1995, 23, 3-15.	0.5	8
401	Linkage mapping of the gene for the LIM-homeoprotein LIM3 (locus Lhx3) to mouse Chromosome 2. <i>Mammalian Genome</i> , 1995, 6, 818-819.	2.2	14
402	Application of the multiple antigenic peptides (MAP) strategy to the production of prohormone convertases antibodies: Synthesis, characterization and use of 8-branched immunogenic peptides. <i>Journal of Peptide Science</i> , 1995, 1, 385-395.	1.4	39
403	The distinct gene expression of the pro-hormone convertases in the rat heart suggests potential substrates. <i>Cell and Tissue Research</i> , 1995, 279, 539-549.	2.9	58
404	Fluorescent Peptidyl Substrates as an Aid in Studying the Substrate Specificity of Human Prohormone Convertase PC1 and Human Furin and Designing a Potent Irreversible Inhibitor. <i>Journal of Biological Chemistry</i> , 1995, 270, 19225-19231.	3.4	80
405	Expression of Candidate Pro-GnRH Processing Enzymes in Rat Hypothalamus and an Immortalized Hypothalamic Neuronal Cell Line. <i>Neuroendocrinology</i> , 1995, 62, 166-177.	2.5	37
406	Parathyroid Hormone Is Preferentially Cleaved to Parathyroid Hormone by the Prohormone Convertase Furin. <i>Journal of Biological Chemistry</i> , 1995, 270, 9517-9525.	3.4	71
407	Processing Specificity and Biosynthesis of the <i>Drosophila melanogaster</i> Convertases dfurin1, dfurin1-CRR, dfurin1-X, and dfurin2. <i>Journal of Biological Chemistry</i> , 1995, 270, 1020-1028.	3.4	49
408	Chromosomal assignment of the genes for proprotein convertases PC4, PC5, and PACE 4 in mouse and human. <i>Genomics</i> , 1995, 26, 123-129.	2.9	32
409	Structural Investigation and Kinetic Characterization of Potential Cleavage Sites of HIV Gp160 by Human Furin and Pc1. <i>Biochemical and Biophysical Research Communications</i> , 1995, 213, 356-361.	2.1	20
410	Processing of secretogranin II by prohormone convertases: Importance of PC1 in generation of secretoneurin. <i>FEBS Letters</i> , 1995, 360, 294-298.	2.8	78
411	Structure-function studies on the biosynthesis and bioactivity of the precursor convertase PC2 and the formation of the PC2/7B2 complex. <i>FEBS Letters</i> , 1995, 362, 151-155.	2.8	21
412	Comparative proteolytic processing of rat prosomatostatin by the convertases PC1, PC2, furin, PACE4 and PC5 in constitutive and regulated secretory pathways. <i>FEBS Letters</i> , 1995, 362, 143-146.	2.8	78
413	Maintained PC1 and PC2 Expression in the AtT-20 Variant Cell Line 6T3 Lacking Regulated Secretion and POMC: Restored POMC Expression and Regulated Secretion after cAMP Treatment. <i>DNA and Cell Biology</i> , 1995, 14, 175-188.	1.9	25
414	7B2 Is a Specific Intracellular Binding Protein of the Prohormone Convertase PC2. <i>Journal of Neurochemistry</i> , 1995, 64, 2303-2311.	3.9	91



#	ARTICLE	IF	CITATIONS
415	Peptidyl substrates containing unnatural amino acid at the P <sup>21</sup> position are potent inhibitors of prohormone convertases. <i>International Journal of Peptide and Protein Research</i> , 1995, 46, 228-237.	0.1	30
416	The distinct gene expression of the pro-hormone convertases in the rat heart suggests potential substrates. <i>Cell and Tissue Research</i> , 1995, 279, 539-549.	2.9	2
417	Mouse insulinoma beta TC3 cells express prodynorphin messenger ribonucleic acid and derived peptides: a unique cellular model for the study of prodynorphin biosynthesis and processing. <i>Endocrinology</i> , 1995, 136, 1187-1196.	2.8	4
418	Age-related alterations in the expression of prohormone convertase messenger ribonucleic acid (mRNA) levels in hypothalamic proopiomelanocortin mRNA neurons in the female C57BL/6J mouse. <i>Endocrinology</i> , 1995, 136, 2721-2729.	2.8	9
419	[13] Pro-protein convertases of subtilisin/kexin family. <i>Methods in Enzymology</i> , 1994, 244, 175-188.	1.0	84
420	Differential expression of the neuroendocrine polypeptide 7B2 in hypothalami of Prader-(Labhart)-Willi syndrome patients. <i>Brain Research</i> , 1994, 657, 281-293.	2.2	31
421	Proteolytic processing of human prorenin in renal and non-renal tissues. <i>Kidney International</i> , 1994, 46, 1522-1524.	5.2	40
422	The family of subtilisin/kexin like pro-protein and pro-hormone convertases: Divergent or shared functions. <i>Biochimie</i> , 1994, 76, 197-209.	2.6	417
423	Co-expression of PC2 and proenkephalin in human tumoral adrenal medullary tissues. <i>Biochimie</i> , 1994, 76, 241-244.	2.6	6
424	Approaches in the study of the posttranslational processing of opioid peptide peptide precursors: Focus on prodynorphin. <i>Regulatory Peptides</i> , 1994, 54, 77-78.	1.9	1
425	Expression of PC2 and PC1/PC3 in human pheochromocytomas. <i>Molecular and Cellular Endocrinology</i> , 1994, 99, 307-314.	3.2	24
426	Structure of the Gene for the Testis-Specific Proprotein Convertase 4 and of Its Alternate Messenger RNA Isoforms. <i>Genomics</i> , 1994, 20, 231-237.	2.9	45
427	Processing of prodynorphin by the prohormone convertase PC1 results in high molecular weight intermediate forms. <i>FEBS Letters</i> , 1994, 337, 60-65.	2.8	75
428	The Mouse Homeoprotein mLIM-3 Is Expressed Early in Cells Derived from the Neuroepithelium and Persists in Adult Pituitary. <i>DNA and Cell Biology</i> , 1994, 13, 1163-1180.	1.9	98
429	Gene Organization of the Mouse Pro-Hormone and Pro-Protein Convertase PC1. <i>DNA and Cell Biology</i> , 1994, 13, 395-407.	1.9	49
430	Large Denseâ€œCore Vesicles in Rat Adrenal After Reserpine: Levels of mRNAs of Soluble and Membraneâ€œBound: Constituents in Chromaffin and Ganglion Cells Indicate a Biosynthesis of Vesicles with Higher Secretory Quanta. <i>Journal of Neurochemistry</i> , 1994, 62, 2448-2456.	3.9	35
431	Design and synthesis of novel inhibitors of prohormone convertases. <i>International Journal of Peptide and Protein Research</i> , 1994, 44, 253-261.	0.1	32
432	Developmental expression of the prohormone convertases PC1 and PC2 in mouse pancreatic islets. <i>Endocrinology</i> , 1994, 135, 1651-1660.	2.8	33

#	ARTICLE	IF	CITATIONS
433	Expression of the prohormone convertase PC2 correlates with the presence of corticotropin-like intermediate lobe peptide in human adrenocorticotropin-secreting tumors. <i>Journal of Clinical Endocrinology and Metabolism</i> , 1994, 79, 1503-1506.	3.6	20
434	Testis-Specific Proprotein Convertase 4: Gene Structure, Optional Exons, and mRNA Isoforms. , 1994, , 388-399.		0
435	Synthesis of the segment (11â€“23) located in the first tandem repeat of plasma kallikrein: comparative binding studies of this and another segment (328â€“343) to high-molecular-mass kininogen. <i>Biomedical Applications</i> , 1993, 615, 251-264.	1.7	1
436	Processing, release and metabolism of cholecystokinin in SK-N-MCIXC cells. <i>Neuropeptides</i> , 1993, 25, 19-30.	2.2	9
437	cDNA structure and in situ localization of the <i>Aplysia californica</i> pro-hormone convertase PC2. <i>FEBS Letters</i> , 1993, 330, 343-346.	2.8	20
438	Region specific expression of furin mRNA in the rat brain. <i>Neuroscience Letters</i> , 1993, 149, 27-30.	2.1	55
439	Molecular cloning and expression of the ovine testicular follicle stimulating hormone receptor. <i>Molecular and Cellular Endocrinology</i> , 1993, 93, 219-226.	3.2	67
440	Protease inhibitors suppress in vitro growth of human small cell lung cancer. <i>Peptides</i> , 1993, 14, 1021-1028.	2.4	21
441	From Proopiomelanocortin to Cancer.. <i>Annals of the New York Academy of Sciences</i> , 1993, 680, 13-19.	3.8	20
442	Mammalian Paired Basic Amino Acid Convertases of Prohormones and Proproteins. <i>Annals of the New York Academy of Sciences</i> , 1993, 680, 135-146.	3.8	82
443	The family of pro-hormone and pro-protein convertases. <i>Biochemical Society Transactions</i> , 1993, 21, 685-691.	3.4	76
444	Ontogeny of the prohormone convertases PC1 and PC2 in the mouse hypophysis and their colocalization with corticotropin and alpha-melanotropin.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1993, 90, 4922-4926.	7.1	125
445	cDNA structure of the mouse and rat subtilisin/kexin-like PC5: a candidate proprotein convertase expressed in endocrine and nonendocrine cells.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1993, 90, 6691-6695.	7.1	235
446	Tissue-Specific and Substrate-Specific Endoproteolytic Cleavage of Monkey Pro-Opiomelanocortin in Heterologous Endocrine Cells: Processing at Lys-Lys Dibasic Pairs. <i>Neuroendocrinology</i> , 1993, 58, 94-105.	2.5	4
447	The cDNA Sequence of the Human Pro-Hormone and Pro-Protein Convertase PC1. <i>DNA and Cell Biology</i> , 1992, 11, 283-289.	1.9	53
448	Regional localization of three convertases, PC1 (Nec-1), PC2 (Nec-2), and furin (Fur), on mouse chromosomes. <i>Genomics</i> , 1992, 13, 1356-1358.	2.9	22
449	Proprotein and prohormone convertases of the subtilisin family. <i>Trends in Endocrinology and Metabolism</i> , 1992, 3, 133-140.	7.1	151
450	Dynamics of 7B2 and galanin expression in solitary magnocellular hypothalamic vasopressin neurons of the homozygous Brattleboro rat. <i>Brain Research</i> , 1992, 585, 275-282.	2.2	14

#	ARTICLE	IF	CITATIONS
451	The cDNA structure of the porcine pro-hormone convertase PC2 and the comparative processing by PC1 and PC2 of the N-terminal glycopeptide segment of porcine POMC. <i>FEBS Letters</i> , 1992, 310, 235-239.	2.8	36
452	Affinity purification of proteinases by a combination of immobilized peptidyl aldehyde and semicarbazone. <i>Biomedical Applications</i> , 1992, 581, 17-29.	1.7	5
453	Testicular expression of PC4 in the rat: molecular diversity of a novel germ cell-specific Kex2/subtilisin-like proprotein convertase. <i>Molecular Endocrinology</i> , 1992, 6, 1559-1570.	3.7	115
454	Distribution and regulation of the prohormone convertases PC1 and PC2 in the rat pituitary. <i>Molecular Endocrinology</i> , 1992, 6, 485-497.	3.7	157
455	The messenger RNA for pituitary protein 7B2 is widely expressed in mouse CNS and castration reduces its pituitary level. <i>Neurochemistry International</i> , 1991, 19, 103-111.	3.8	4
456	Chromosomal assignments of the genes for neuroendocrine convertase PC1 (NEC1) to human 5q15â€²21, neuroendocrine convertase PC2 (NEC2) to human 20p11.1â€²11.2, and furin (mouse 7[D1-E2] region). <i>Genomics</i> , 1991, 11, 103-107.	2.9	47
457	Expression and Sorting of Rat Plasma Kallikrein in POMC-Producing AtT-20 Cells. <i>DNA and Cell Biology</i> , 1991, 10, 259-269.	1.9	6
458	Gene structure and chromosomal localization of plasma kallikrein. <i>Biochemistry</i> , 1991, 30, 1628-1635.	2.5	89
459	The production by alternate splicing of two mRNAs differing by one codon could be an intrinsic property of neuroendocrine protein 7B2 gene expression in man. <i>Biochemical and Biophysical Research Communications</i> , 1991, 174, 156-162.	2.1	9
460	Immunological identification and sequence characterization of a peptide derived from the processing of neuroendocrine protein 7B2. <i>FEBS Letters</i> , 1991, 294, 23-26.	2.8	23
461	Neuroanatomical and Functional Studies of Peptide Precursor-Processing Enzymes. <i>Enzyme</i> , 1991, 45, 285-300.	0.7	23
462	N-linked glycosylation affects the processing of mouse submaxillary gland prorenin in transfected AtT20 cells. <i>FEBS Journal</i> , 1991, 198, 535-540.	0.2	14
463	PC1 and PC2 are proprotein convertases capable of cleaving proopiomelanocortin at distinct pairs of basic residues.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1991, 88, 3564-3568.	7.1	586
464	Primary structure of mouse chromogranin B deduced from cDNA sequence. <i>Nucleic Acids Research</i> , 1990, 18, 1298-1298.	14.5	32
465	cDNA Sequence of Two Distinct Pituitary Proteins Homologous to Kex2 and Furin Gene Products: Tissue-Specific mRNAs Encoding Candidates for Pro-Hormone Processing Proteinases. <i>DNA and Cell Biology</i> , 1990, 9, 415-424.	1.9	529
466	Assignment of the gene for neuroendocrine protein 7B2 (SGNE1 locus) to mouse chromosome region 2[E3â€²F3] and to human chromosome region 15q11-q15. <i>Genomics</i> , 1990, 6, 436-440.	2.9	28
467	Mouse Plasma Kallikrein: cDNA Structure, Enzyme Characterization, and Comparison of Protein and mRNA Levels among Species. <i>DNA and Cell Biology</i> , 1990, 9, 737-748.	1.9	31
468	Fragmentation of bovine chromogranin A by plasma kallikrein. <i>Life Sciences</i> , 1990, 46, 1427-1433.	4.3	24

#	ARTICLE	IF	CITATIONS
469	Syntheses of argininal semicarbazone containing peptides and their applications in the affinity chromatography of serine proteinases. International Journal of Peptide and Protein Research, 1990, 36, 7-17.	0.1	11
470	The cDNA Structure of Rat Plasma Kallikrein. DNA and Cell Biology, 1989, 8, 563-574.	5.2	28
471	Rat plasma kallikrein: purification, NH <sub>2</sub> -terminal sequencing and development of a specific radioimmunoassay. BBA - Proteins and Proteomics, 1989, 999, 103-110.	2.1	9
472	Functional diversity of bioactive peptides in the nervous system itself: "How the brain may understand". Bioscience Reports, 1989, 9, 693-700.	2.4	10
473	Microsequencing evidence for the maturation of human proopiomelanocortin into an 18 amino acid $\beta$ -melanocyte stimulating hormone [h <sup>12</sup> MSH(5 <sup>1-22</sup> )] in nonpituitary tissue. Peptides, 1989, 10, 83-87.	2.4	10
474	Substrate phosphorylation can inhibit proteolysis by trypsin-like enzymes. Archives of Biochemistry and Biophysics, 1989, 272, 274-280.	3.0	52
475	Stable and transient expression of mouse submaxillary gland renin cDNA in AtT20 cells: Proteolytic processing and secretory pathways. FEBS Letters, 1989, 245, 70-74.	2.8	16
476	cDNA sequence of neuroendocrine protein 7B2 expressed in beta cell tumors of transgenic mice. International Journal of Peptide and Protein Research, 1989, 33, 39-45.	0.1	50
477	Distribution of a novel pituitary protein (7B2) in mammalian gastrointestinal tract and pancreas. Digestive Diseases and Sciences, 1988, 33, 718-723.	2.3	17
478	The high-molecular-mass kininogen deficient rat expresses all kininogen mRNA species, but does not export the high-molecular-mass kininogen synthesized. FEBS Letters, 1988, 239, 59-64.	2.8	20
479	Regional processing of the N- and C-terminal domains of proopiomelanocortin in monkey pituitary and brain. Neuropeptides, 1988, 11, 111-118.	2.2	7
480	Localization of the Human Prostatic Secretory Protein PSP <sub>94</sub> and its mRNA in the Epithelial Cells of the Prostate. Journal of Andrology, 1988, 9, 253-260.	2.0	25
481	NH <sub>2</sub> -terminal fragment of rat pro-atrial natriuretic factor in the circulation: Identification, radioimmunoassay and half-life. Peptides, 1988, 9, 47-53.	2.4	113
482	Structural and immunological homology of human and porcine pituitary and plasma IRCM-serine protease 1 to plasma kallikrein: marked selectivity for pairs of basic residues suggests a widespread role in pro-hormone and pro-enzyme processing. Biochimie, 1988, 70, 33-46.	2.6	22
483	Correct processing and secretion of a human prostatic secretory protein (PSP94) in Escherichia coli. Gene, 1988, 73, 479-487.	2.2	10
484	Two-dimensional immunoblotting analysis and immunocytochemical localization of the secretory polypeptide 7B2 in adrenal medulla. Neuroscience Letters, 1988, 95, 81-87.	2.1	27
485	The Possible Role of Plasma Kallikrein in Pro-Hormone and Pro-Enzyme Processing. , 1988, , 179-188.		0
486	Characterization of N-Terminal Fragment of Proopiomelanocortin in Cerebrospinal Fluid*. Journal of Clinical Endocrinology and Metabolism, 1987, 65, 198-202.	3.6	3

#	ARTICLE	IF	CITATIONS
487	Gamma-melanotropin response to ovine corticotropin releasing factor in normal humans. <i>Neuropeptides</i> , 1987, 9, 269-282.	2.2	6
488	IRCM-Serine Protease # 1 from Pituitary and Heart A Common Prohormone Maturation Enzyme. <i>Annals of the New York Academy of Sciences</i> , 1987, 493, 403-405.	3.8	0
489	Chromogranin B (secretogranin I), a putative precursor of two novel pituitary peptides through processing at paired basic residues. <i>FEBS Letters</i> , 1987, 224, 142-148.	2.8	84
490	Chromogranin A can act as a reversible processing enzyme inhibitor. <i>FEBS Letters</i> , 1987, 211, 144-150.	2.8	80
491	Isolation from Human Seminal Plasma of an Abundant 16 kDa Protein Originating from the Prostate, Its Identification with a 94-Residue Peptide Originally Described As I <sup>2</sup> inhibin. <i>Journal of Andrology</i> , 1987, 8, 182-189.	2.0	66
492	Elevation of plasma 7B2 (a novel pituitary protein) in cord blood at obstetrical delivery and the possible correlation with GH. <i>Life Sciences</i> , 1987, 41, 1921-1927.	4.3	6
493	Bovine adrenal chromaffin granules are a site of synthesis of atrial natriuretic factor. <i>Biochemical and Biophysical Research Communications</i> , 1987, 147, 957-963.	2.1	68
494	Evidence for a novel pituitary protein (7B2) in human brain, cerebrospinal fluid and plasma: Brain concentrations in controls and patients with Alzheimer's disease. <i>Peptides</i> , 1987, 8, 593-598.	2.4	38
495	Molecular Cloning and Sequence of the cDNA for a 94-Amino-Acid Seminal Plasma Protein Secreted by the Human Prostate. <i>DNA and Cell Biology</i> , 1987, 6, 23-29.	5.2	71
496	The pituitary polypeptide $\alpha$ -7B2 is associated with LH/FSH and TSH cells and is localized within secretory vesicles. <i>Cell and Tissue Research</i> , 1987, 250, 205-214.	2.9	52
497	Presence of the Novel Pituitary Protein $\gamma$ 7B2 in Bovine Chromaffin Granules: Possible Co-Release of 7B2 and Catecholamine as Induced by Nicotine. <i>Journal of Neurochemistry</i> , 1987, 49, 1810-1814.	3.9	25
498	Developmental changes in immunoreactive content of novel pituitary protein 7B2 in human pancreas and its identification in pancreatic tumors. <i>Diabetes</i> , 1987, 36, 1276-1279.	0.6	8
499	Purification of ovine placental lactogen ( $\alpha$ PL) using high-performance liquid chromatography. <i>FEBS Letters</i> , 1986, 199, 259-264.	2.8	16
500	Binding inhibition by monoclonal antibodies of ovine placental lactogen to growth hormone receptors in human liver. <i>FEBS Letters</i> , 1986, 201, 168-172.	2.8	1
501	Purification and sequence determination of bovine atrial natriuretic factor. <i>Life Sciences</i> , 1986, 38, 1309-1315.	4.3	22
502	Demonstration of the vasopressin associated glycopeptide in the brain and peripheral tissues of the Brattleboro rat. <i>Neuropeptides</i> , 1986, 7, 79-85.	2.2	23
503	CNS distribution of a novel pituitary protein $\alpha$ -7B2 <sup>TM</sup> : localization in secretory and synaptic vesicles. <i>Brain Research</i> , 1986, 380, 349-356.	2.2	53
504	Propressophysin is present in neurones at multiple sites in wistar and homozygous brattleboro rat brain. <i>Brain Research</i> , 1986, 379, 171-175.	2.2	25

#	ARTICLE	IF	CITATIONS
505	Specific Release of a Novel Pituitary Polypeptide, 7B2, from Rat Anterior Pituitary Cells in vitro by Luteinizing Hormone-Releasing Hormone. <i>Neuroendocrinology</i> , 1986, 44, 373-377.	2.5	33
506	Isolation and primary structure of novel neurointermediate pituitary peptides derived from the C-terminal of the rat vasopressin-neurophysin precursor (propressophysin). <i>FEBS Journal</i> , 1986, 156, 137-142.	0.2	15
507	Homologous IRCM-Serine Protease 1 from pituitary, heart atrium and ventricle: A common pro-hormone maturation enzyme?. <i>Bioscience Reports</i> , 1986, 6, 835-844.	2.4	41
508	A novel pituitary protein (7B2)-like immunoreactivity is secreted by a rat pheochromocytoma cell line (PC12). <i>Journal of Endocrinology</i> , 1986, 108, 151-155.	2.6	17
509	Inhibition of release of a novel pituitary polypeptide, 7B2, follicle-stimulating hormone, and luteinizing hormone from rat anterior pituitary cells in vitro by human $\beta^2$ -inhibin. <i>Canadian Journal of Physiology and Pharmacology</i> , 1986, 64, 1259-1262.	1.4	5
510	Light and electron microscopic localization of the N-terminal fragment of human pro-opiomelanocortin in the human pituitary gland and in neoplasms. <i>Virchows Archiv A, Pathological Anatomy and Histopathology</i> , 1985, 408, 281-287.	1.4	5
511	Novel approach to rapid and sensitive localization of protein disulfide bridges by high-performance liquid chromatography and electrochemical detection. <i>Journal of Chromatography A</i> , 1985, 326, 339-348.	3.7	12
512	Ontogeny of a novel pituitary protein (7B2) in the human fetal intestine. <i>Regulatory Peptides</i> , 1985, 12, 289-296.	1.9	12
513	Regional distribution of a novel pituitary protein (7B2) in the rat spinal cord: Effect of neonatal capsaicin treatment and thoracic cord transection. <i>Neuroscience Letters</i> , 1985, 55, 151-156.	2.1	21
514	Immunoreactivity of vasopressin and a novel pituitary protein $\sim$ 7B2 $\sim$ ™ in long-evans and brattleboro rat hypothalamus and hypophysis. <i>Neuroscience Letters</i> , 1985, 60, 7-12.	2.1	13
515	Regional distribution of a novel pituitary protein (7B2) in the rat brain. <i>Brain Research</i> , 1985, 338, 91-96.	2.2	41
516	Pro-opiomelanocortin-related peptides in cerebrospinal fluid: A study of manic-depressive disorder. <i>Psychiatry Research</i> , 1985, 16, 287-302.	3.3	20
517	Isolation and characterization of four proteases from porcine pituitary neurointermediate lobes: Relationship to the maturation enzyme of prohormones. <i>Neuropeptides</i> , 1985, 5, 493-496.	2.2	16
518	Structure-activity relationships of atrial natriuretic factor (ANF). II. Effect of chain-length modifications on vascular reactivity. <i>Biochemical and Biophysical Research Communications</i> , 1985, 126, 178-184.	2.1	69
519	Gawł, a novel human pituitary polypeptide: Isolation, immunocytochemical localization and complete amino acid sequence. <i>Biochemical and Biophysical Research Communications</i> , 1985, 126, 602-609.	2.1	31
520	Structure-activity relationships of atrial natriuretic factor (ANF) III correlation of receptor affinity with relative potency on aldosterone production in zona glomerulosa cells. <i>Biochemical and Biophysical Research Communications</i> , 1985, 132, 360-367.	2.1	38
521	Identification of a biologically active circulating form of rat atrial natriuretic factor. <i>Biochemical and Biophysical Research Communications</i> , 1985, 130, 981-986.	2.1	164
522	ANF (Arg 101 - Tyr 126) is the peptide secreted by rat atrial cardiocytes in culture. <i>Biochemical and Biophysical Research Communications</i> , 1985, 130, 1217-1225.	2.1	11

#	ARTICLE	IF	CITATIONS
523	$\hat{I}^2$ -Inhibin contains the active core of human seminal plasma $\hat{I}^2$ -inhibin: synthesis and bioactivity. FEBS Letters, 1985, 181, 57-63.	2.8	15
524	Effects of several pro-opiomelanocortin derived peptides on steroidogenesis in ovine and bovine adrenal cells. The Journal of Steroid Biochemistry, 1985, 23, 185-190.	1.1	15
525	A NOVEL PITUITARY PROTEIN (7B2) IN THE RAT UROGENITAL TRACT. Biomedical Research, 1985, 6, 139-143.	0.9	8
526	Microsequence of Polypeptide Hormones: Its Usefulness to Monitor the Isolation of Novel Molecules. , 1985, , 195-209.		0
527	Effects of Proopi melanocortin-Derived Peptides, Methionine-Enkephal in and Forskolin on the Maturation of Ovine Fetal Adrenal Cells in Culture. Biology of Reproduction, 1984, 31, 694-704.	2.7	25
528	Isolation, structure, and synthesis of a human seminal plasma peptide with inhibin-like activity. Science, 1984, 223, 1199-1202.	12.6	73
529	Amino acid sequence of rat submaxillary tonin reveals similarities to serine proteases. Nature, 1984, 307, 555-558.	27.8	36
530	Structure of the carboxyl propeptide of chicken type II procollagen determined by DNA and protein sequence analysis. Biochemistry, 1984, 23, 617-624.	2.5	69
531	Primary structure of a high M <sub>r</sub> form of rat atrial natriuretic factor. FEBS Letters, 1984, 167, 352-356.	2.8	118
532	The major androgen-dependent protease in dog prostate belongs to the kallikrein family: confirmation by partial amino acid sequencing. FEBS Letters, 1984, 175, 1-7.	2.8	36
533	Complete amino acid sequence of human seminal plasma $\hat{I}^2$ -inhibin. FEBS Letters, 1984, 175, 349-355.	2.8	105
534	Partial amino acid sequence of a human seminal plasma peptide with inhibin-like activity. FEBS Letters, 1984, 167, 98-102.	2.8	37
535	Atrial pronatriodilatin: a precursor for natriuretic factor and cardiodilatin. FEBS Letters, 1984, 172, 80-86.	2.8	49
536	Structure-activity relationships of atrial natriuretic factor (ANF). I. Natriuretic activity and relaxation of intestinal smooth muscle. Biochemical and Biophysical Research Communications, 1984, 125, 938-946.	2.1	95
537	Tissue Distribution and Molecular Forms of a Novel Pituitary Protein in the Rat. Neuroendocrinology, 1984, 39, 453-458.	2.5	107
538	Amino acid sequence of homologous rat atrial peptides: natriuretic activity of native and synthetic forms.. Proceedings of the National Academy of Sciences of the United States of America, 1984, 81, 2640-2644.	7.1	213
539	Molecular cloning and characterization of DNA sequences encoding rat and human atrial natriuretic factors.. Proceedings of the National Academy of Sciences of the United States of America, 1984, 81, 6325-6329.	7.1	125
540	Characterization of $\hat{I}^2$ -endorphin immunoreactive peptides in rat pituitary and brain by coupled gel and reversed-phase high-performance liquid chromatography. Journal of Chromatography A, 1983, 266, 163-172.	3.7	29

#	ARTICLE	IF	CITATIONS
541	Enzymatic maturation of pro-opiomelanocortin by anterior pituitary granules. Journal of Chromatography A, 1983, 266, 213-224.	3.7	7
542	Peptides of the neurointermediary lobe of rat pituitary stimulate GH secretion in vitro. Molecular and Cellular Endocrinology, 1983, 32, 47-55.	3.2	6
543	Presence of vasopressin, oxytocin and neurophysin in the retina of mammals, effect of light and darkness, comparison with the neuropeptide content of the neurohypophysis and the pineal gland. Peptides, 1983, 4, 509-515.	2.4	27
544	Metabolic clearance rate and half-time disappearance rate of human N-terminal and adrenocorticotropin of pro-opiomelanocortin in the rat: A comparative study. Life Sciences, 1983, 33, 2599-2608.	4.3	16
545	Regional heterogeneity in the processing of pro-opiomelanocortin in rat brain. Life Sciences, 1983, 33, 49-52.	4.3	34
546	Immunocytochemistry of the C-terminal peptide of proressophysin (CPP): Relationship to vasopressin, oxytocin and neurophysin. Neuropeptides, 1983, 3, 321-336.	2.2	12
547	Isolation and NH <sub>2</sub> -terminal sequence of a highly conserved human and porcine pituitary protein belonging to a new superfamily. Archives of Biochemistry and Biophysics, 1983, 225, 525-534.	3.0	140
548	Purification of three rat atrial natriuretic factors and their amino acid composition. FEBS Letters, 1983, 164, 286-290.	2.8	77
549	Enzymatic cleavage of pro-opiomelanocortin by anterior pituitary granules. FEBS Letters, 1983, 159, 68-74.	2.8	14
550	Response of Human Aldosteronoma Cells in Culture to the N-Terminal Glycopeptide of Pro-Opiomelanocortin and <sup>125</sup> I-MSH. Hormone and Metabolic Research, 1983, 15, 181-184.	1.5	21
551	Proteases and posttranslational processing of prohormones: a review. Canadian Journal of Biochemistry and Cell Biology, 1983, 61, 501-515.	1.3	136
552	Primary structure determination of <i>Escherichia coli</i> heat-stable enterotoxin of porcine origin. Canadian Journal of Biochemistry and Cell Biology, 1983, 61, 287-292.	1.3	82
553	The carboxy terminus of the precursor to vasopressin and neurophysin: immunocytochemistry in rat brain. Science, 1982, 217, 853-855.	12.6	35
554	Effects of pro-opiomelanocortin fragments on release of catecholamines from adrenal chromaffin cells. Neuroscience Letters, 1982, 28, 199-204.	2.1	6
555	Peptide mapping of <sup>125</sup> I-labelled membrane protein of influenza viruses by reverse-phase high-performance liquid chromatography. Journal of Virological Methods, 1982, 4, 77-85.	2.1	5
556	Concomitant changes of ACTH, <sup>125</sup> I-endorphin and N-terminal portion of pro-opiomelanocortin in rats. Life Sciences, 1982, 30, 1159-1164.	4.3	21
557	Characterization of multiple forms of porcine anterior pituitary pro-opiomelanocortin N-terminal glycopeptide. Biochemistry, 1982, 21, 5341-5346.	2.5	33
558	Isolation and NH <sub>2</sub> -terminal sequence of a novel porcine anterior pituitary polypeptide. FEBS Letters, 1982, 147, 261-266.	2.8	129



#	ARTICLE	IF	CITATIONS
559	The primary structure of human $\hat{\nu}^2$ -lipotropin. FEBS Letters, 1982, 147, 267-272.	2.8	6
560	CORTICOTROPIN RELEASING FACTOR (CRF): EFFECTS ON THE RELEASE OF PRO-OPIOMELANOCORTIN (POMC)-RELATED PEPTIDES BY HUMAN ANTERIOR PITUITARY CELLS IN VITRO. Endocrinology, 1982, 111, 1388-1390.	2.8	64
561	Purification of radiolabeled and native polypeptides by gel permeation high-performance liquid chromatography. Analytical Biochemistry, 1982, 125, 406-414.	2.4	23
562	Isolation and purification of Escherichia coli heat-stable enterotoxin of porcine origin. Analytical Biochemistry, 1982, 127, 267-275.	2.4	49
563	Distribution pattern in the human pituitary and hypothalamus of a new neuropeptide: The C-terminal glycoprotein-fragment of human pro-pressophysin (CPP). Histochemistry, 1982, 75, 319-326.	1.9	14
564	Expression of variant forms of proopi melanocortin, the common precursor to corticotropin and $\beta$ -lipotropin in the rat pars intermedia. Biochemistry, 1981, 20, 2475-2481.	2.5	23
565	Effect of chronic morphine treatment on $\hat{\nu}^2$ -endorphin biosynthesis by the rat neurointermediate lobe. European Journal of Pharmacology, 1981, 72, 313-321.	3.5	31
566	The missing fragment of the pro-sequence of human pro-opiomelanocortin: Sequence and evidence for C-terminal amidation. Biochemical and Biophysical Research Communications, 1981, 102, 710-716.	2.1	64
567	The complete sequence of a novel human pituitary glycopeptide homologous to pig posterior pituitary glycopeptide. Biochemical and Biophysical Research Communications, 1981, 100, 901-907.	2.1	59
568	Biosynthesis of $\hat{\nu}^2$ -endorphin by the neurointermediate lobes from rats treated with morphine or alcohol. Life Sciences, 1981, 29, 1973-1982.	4.3	41
569	Reinvestigation of the N-terminal amino acid sequence of $\hat{\nu}^2$ -lipotropin from human pituitary glands. Biochemical and Biophysical Research Communications, 1981, 103, 1329-1335.	2.1	19
570	Immunohistochemical localization of $\hat{\nu}^3$ -melanocyte-stimulating hormone in the rat pituitary gland. Regulatory Peptides, 1981, 2, 81-89.	1.9	8
571	Complete amino acid sequence of a human pituitary glycopeptide: an important maturation product of pro-opiomelanocortin. Proceedings of the National Academy of Sciences of the United States of America, 1981, 78, 4236-4240.	7.1	61
572	Evidence for a Signal Sequence at the N Terminus of the Common Precursor to Adrenocorticotrophin and $\beta$ -Lipotropin in Mouse Pituitary Cells. FEBS Journal, 1981, 116, 255-259.	0.2	12
573	Chemistry and biosynthesis of pro-opiomelanocortin. Molecular and Cellular Biochemistry, 1981, 34, 101-127.	3.1	88
574	Sequence homologies between tonin, nerve growth factor $\hat{\nu}^3$ -subunit, epidermal growth factor-binding protein and serine proteases. Nature, 1981, 292, 383-384.	27.8	53
575	Effect of N-Terminal Portion of Pro-opiomelanocortin on Aldosterone Release by Human Adrenal Adenoma In Vitro*. Journal of Clinical Endocrinology and Metabolism, 1981, 52, 1053-1056.	3.6	56
576	Biosynthesis of $\hat{\nu}^2$ -Endorphin from Proopi melanocortin. , 1981, , 65-87.		0

#	ARTICLE	IF	CITATIONS
577	A rapid high-performance liquid chromatography purification method of Iodinated polypeptide hormones. <i>Analytical Biochemistry</i> , 1980, 109, 185-191.	2.4	61
578	Reversed-phase high-performance liquid chromatographic purification and characterization of the adrenocorticotropin/lipotropin precursor and its fragments. <i>Journal of Chromatography A</i> , 1980, 193, 291-299.	3.7	34
579	A novel human pituitary peptide containing the $\hat{I}^3$ -MSH sequence. <i>Nature</i> , 1980, 285, 415-416.	27.8	39
580	Processing of Two Forms of the Common Precursor to alpha-Melanotropin and beta-Endorphin in the Rat Pars Intermedia. Evidence for and Partial Characterization of New Pituitary Peptides. <i>FEBS Journal</i> , 1980, 110, 387-396.	0.2	30
581	Purification and characterization of the N-terminal fragment of pro-opiomelanocortin from human pituitaries: Homology to the bovine sequence. <i>Biochemical and Biophysical Research Communications</i> , 1980, 95, 1417-1424.	2.1	35
582	Two glycosylation sites on the N-terminal segment of porcine pars distalis pro-opiomelanocortin. <i>FEBS Letters</i> , 1980, 122, 279-282.	2.8	14
583	Substrate specificity of the enzyme tonin: cleavage of substance P. <i>FEBS Letters</i> , 1980, 113, 173-176.	2.8	24
584	PRESENCE OF A PRE-SEQUENCE (SIGNAL SEQUENCE) IN THE COMMON PRECURSOR TO ACTH AND ENDORPHIN AND THE ROLE OF GLYCOSYLATION IN PROCESSING OF THE PRECURSOR AND SECRETION OF ACTH AND ENDORPHIN. <i>Annals of the New York Academy of Sciences</i> , 1980, 343, 79-93.	3.8	45
585	CHEMICAL CHARACTERIZATION OF RAT $\hat{I}^2$ -MSH/ $\hat{I}^2$ -ENDORPHIN PRECURSOR FROM PARS INTERMEDIA. <i>Annals of the New York Academy of Sciences</i> , 1980, 343, 443-446.	3.8	8
586	Partial N-terminal amino acid sequence of pro-opio-melanocortin (ACTH/beta-LPH precursor) from rat pars intermedia. <i>Biochemical and Biophysical Research Communications</i> , 1980, 92, 1042-1051.	2.1	36
587	Specific cleavage of beta-LPH and ACTH by tonin: Release of an opiate-like peptide beta-LPH (61-78). <i>Biochemical and Biophysical Research Communications</i> , 1979, 86, 1002-1013.	2.1	30
588	Concomitant synthesis of beta-endorphin and alpha-melanotropin from two forms of pro-opiomelanocortin in the rat pars intermedia.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1979, 76, 5085-5089.	7.1	91
589	Immunological characterization of $\hat{I}^2$ -lipotropin fragments (endorphin, $\hat{I}^2$ -MSH, and N-fragment) from fish pituitaries. <i>General and Comparative Endocrinology</i> , 1978, 34, 163-168.	1.8	29
590	Isolation and partial characterization of a biosynthetic N-terminal methionyl peptide of bovine pars intermedia: Relationship to ubiquitin. <i>Biochemical and Biophysical Research Communications</i> , 1978, 80, 600-608.	2.1	11
591	Biosynthesis of a ubiquitin-related peptide in rat brain and in human and mouse pituitary tumors. <i>Biochemical and Biophysical Research Communications</i> , 1978, 84, 874-885.	2.1	8
592	Biosynthesis of beta-endorphin, beta-lipotropin and the putative ACTH-LPH precursor in the frog pars intermedia. <i>Life Sciences</i> , 1978, 23, 2281-2291.	4.3	34
593	Content of $\hat{I}^2$ -LPH and its fragments (including endorphins) in anterior and intermediate lobes of the bovine pituitary gland. <i>Life Sciences</i> , 1978, 22, 1715-1722.	4.3	61
594	Beta-endorphin and beta-lipotropin secretion by an acth-secreting mouse pituitary tumor. <i>FEBS Letters</i> , 1978, 90, 353-356.	2.8	6

#	ARTICLE	IF	CITATIONS
595	Biosynthesis of beta-endorphin from beta-lipotropin and a larger molecular weight precursor in rat pars intermedia.. Proceedings of the National Academy of Sciences of the United States of America, 1978, 75, 4719-4723.	7.1	157
596	In vitro biosynthesis and chemical characterization of beta-lipotropin, gamma-lipotropin, and beta-endorphin in rat pars intermedia.. Proceedings of the National Academy of Sciences of the United States of America, 1978, 75, 3153-3157.	7.1	67
597	In vitro biosynthesis of beta-endorphin, gamma-lipoprotein, and beta-lipotropin by the pars intermedia of beef pituitary glands.. Proceedings of the National Academy of Sciences of the United States of America, 1977, 74, 4276-4280.	7.1	41
598	In vitro biosynthesis of beta-endorphin in pituitary glands.. Proceedings of the National Academy of Sciences of the United States of America, 1977, 74, 1403-1406.	7.1	36
599	Effects of peptides of pituitary origin on the formation of C21 steroid by fetal calf adrenal cells in culture. Biochemical and Biophysical Research Communications, 1977, 79, 553-560.	2.1	12
600	The complete sequence of sheep beta-endorphin. Biochemical and Biophysical Research Communications, 1977, 74, 1528-1535.	2.1	27
601	Morphine-like activity of sheep $\hat{1}^2$ -lipotropin and of its tryptic fragments. Canadian Journal of Biochemistry, 1977, 55, 35-40.	1.4	14
602	Inhibition by beta-endorphin of dopamine-sensitive anenylate cyclase in rat striatum. Biochemical and Biophysical Research Communications, 1977, 77, 442-447.	2.1	15
603	$\hat{1}^2$ -endorphin induced akinesia in rats: Effect of apomorphine and $\hat{1}^{\pm}$ -methyl-p-tyrosine and related modifications of dopamine turnover in the basal ganglia. Life Sciences, 1977, 20, 1149-1155.	4.3	42
604	A $\hat{1}$ -LPH PRECURSOR MODEL: RECENT DEVELOPMENTS CONCERNING MORPHINE-LIKE SUBSTANCES. Annals of the New York Academy of Sciences, 1977, 297, 84-105.	3.8	27
605	Isolation of peptides with opiate activity from sheep and human pituitaries: Relationship to beta-lipotropin. Biochemical and Biophysical Research Communications, 1976, 72, 472-478.	2.1	141
606	Transport of ions of one kind through thin membranes. Journal of Membrane Biology, 1974, 16, 1-16.	2.1	35
607	Transport of ions of one kind through thin membranes. Journal of Membrane Biology, 1974, 16, 17-42.	2.1	30
608	Transport of ions of one kind through thin membranes. Journal of Membrane Biology, 1972, 10, 171-192.	2.1	57
609	Osteopontin as a novel substrate for proprotein convertase 5/6 (PCSK5) in bone. Bone Abstracts, 0, , .	0.0	0