Elif Bilgin Dogru

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/7218110/publications.pdf

Version: 2024-02-01

840776 996975 36 290 11 15 citations h-index g-index papers 36 36 36 640 docs citations times ranked citing authors all docs

#	Article	IF	CITATIONS
1	Clinical and Prognostic Significance of Coagulation Assays in Advanced Epithelial Ovarian Cancer. International Journal of Gynecological Cancer, 2013, 23, 276-281.	2.5	28
2	Serum IGF-1 and IGFBP-3 levels as clinical markers for patients with lung cancer. Biomedical Reports, 2016, 4, 609-614.	2.0	21
3	EMMPRIN and ADAM12 in prostate cancer: preliminary results of a prospective study. Tumor Biology, 2014, 35, 11647-11653.	1.8	17
4	Clinical and prognostic significance of coagulation assays in melanoma. Melanoma Research, 2012, 22, 368-375.	1.2	16
5	Clinical Significance of Circulating Serum Cellular Heat Shock Protein 90 (HSP90) Level in Patients with Cutaneous Malignant Melanoma. Asian Pacific Journal of Cancer Prevention, 2017, 18, 599-601.	1.2	16
6	Clinical significance of serum insulin-like growth factor-1 (IGF-1) and insulin-like growth factor binding protein-3 (IGFBP-3) in patients with breast cancer. Tumor Biology, 2014, 35, 9303-9309.	1.8	15
7	Clinical significance of serum interleukin-29, interleukin-32, and tumor necrosis factor alpha levels in patients with gastric cancer. Tumor Biology, 2016, 37, 405-412.	1.8	15
8	Serum activated leukocyte cell adhesion molecule and intercellular adhesion molecule-1 in patients with gastric cancer: Can they be used as biomarkers?. Biomedicine and Pharmacotherapy, 2016, 77, 86-91.	5.6	14
9	Androgen Stimulation of PCA3 and miR-141 and Their Release from Prostate Cancer Cells. Cell Journal, 2015, 16, 488-93.	0.2	13
10	Clinical significance of serum epidermal growth factor receptor (EGFR) levels in patients with breast cancer. Cytokine, 2015, 71, 66-70.	3.2	12
11	miR-141 and miR-375 induction and release are different from PSA mRNA and PCA3 upon androgen stimulation of LNCaP cells. Biomedical Reports, 2013, 1, 802-806.	2.0	11
12	Clinical Signifıcance of Serum Ykl-40 (Chitinase-3-Like-1 Protein) as a Biomarker in Melanoma: an Analysis of 112 Turkish Patients. Asian Pacific Journal of Cancer Prevention, 2017, 18, 1383-1387.	1.2	10
13	Clinical significance of serum fibronectin and vitronectin levels in melanoma patients. Melanoma Research, 2014, 24, 475-479.	1.2	9
14	Clinical Significance of Coagulation Assays in Metastatic Pancreatic Adenocarcinoma. Journal of Gastrointestinal Cancer, 2013, 44, 404-409.	1.3	8
15	Serum M65 as a Biomarker for Metastatic Renal Cell Carcinoma. Clinical Genitourinary Cancer, 2013, 11, 290-296.	1.9	8
16	Serum levels of vascular cell adhesion molecule-1 (VCAM-1) may have diagnostic, predictive, and prognostic roles in patients with lung cancer treated with platinum-based chemotherapy. Tumor Biology, 2014, 35, 7871-7875.	1.8	8
17	Levels of serum fibronectin as a biomarker in gastric cancer patients: Correlation with clinical diagnosis and outcome. Molecular and Clinical Oncology, 2016, 4, 655-659.	1.0	8
18	Clinical Significance of Serum Galectin-3 Levels in Gastric Cancer Patients. Journal of Gastrointestinal Cancer, 2016, 47, 182-186.	1.3	8

#	Article	IF	Citations
19	Markers of Bone Metastases in Breast and Lung Cancers. Asian Pacific Journal of Cancer Prevention, 2012, 13, 4331-4334.	1.2	8
20	Clinical significance of serum M30 and M65 levels in metastatic pancreatic adenocarcinoma. Tumor Biology, 2013, 34, 3529-3536.	1.8	7
21	Clinical Significance of Serum Membrane-Bound Mucin-2 Levels in Breast Cancer. Biomolecules, 2019, 9, 40.	4.0	7
22	For Which Cancer Types can Neuron-Specific Enolase be Clinically Helpful in Turkish Patients?. Asian Pacific Journal of Cancer Prevention, 2013, 14, 2541-2544.	1.2	7
23	Clinical significance of serum claudin-1 levels in melanoma patients. Melanoma Research, 2016, 26, 377-381.	1.2	5
24	Circulating serum levels of angiopoietin-1 and angiopoietin-2 in nasopharynx and larynx carcinoma patients. Tumor Biology, 2016, 37, 8979-8983.	1.8	4
25	Clinical significance of serum Protease-Activated Receptor-1 (PAR-1) levels in patients with cutaneous melanoma. BBA Clinical, 2016, 5, 166-169.	4.1	3
26	Clinical significance of serum laminin levels in patients with lung cancer. Biomedical Reports, 2016, 4, 485-488.	2.0	3
27	Significance of serum neural precursor cell‑expressed developmentally downregulated protein 9 in melanoma. Molecular and Clinical Oncology, 2017, 8, 204-208.	1.0	3
28	Clinical significance of serum laminin and type-IV collagen levels in cutaneous melanoma patients. Molecular and Clinical Oncology, 2016, 5, 195-200.	1.0	2
29	Evaluation of epidermal growth factor receptor serum levels and their association with clinicopathological characteristics in patients with colorectal cancer. Molecular and Clinical Oncology, 2017, 7, 787-797.	1.0	2
30	Serum leptin levels may have diagnostic and predictive roles in patients with pancreatic adenocarcinoma treated with gemcitabine-based chemotherapy. Journal of B U on, 2016, 21, 895-902.	0.4	2
31	Angiogenic factors in small vell lung cancer (VEGF, VEGFR-1, VEGFR-2). Turk Onkoloji Dergisi, 2012, 27, 55-61.	0.0	O
32	Clinical significance of serum protease-activated receptor 1 (PAR1) level in patients with breast cancer. Journal of Oncological Science, 2016, 2, 7-11.	0.1	0
33	Clinical significance of caveolin-1 and fibronectin in pancreatic cancer. European Journal of Cancer, 2017, 72, S78-S79.	2.8	0
34	Serum caveolin-1 and fibronectin levels as clinical markers for patients with malignant melanoma. European Journal of Cancer, 2017, 72, S125-S126.	2.8	0
35	The biological role of AKT serine/threonine kinase 2 in lung cancer. European Journal of Cancer, 2017, 72, S138.	2.8	0
36	Circulating interleukin-18 (IL-18) as a predictor of response to gemcitabine based chemotherapy in patients with pancreatic adenocarcinoma Journal of Clinical Oncology, 2016, 34, e15678-e15678.	1.6	0