

Vladimir Semiglazov

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7218205/publications.pdf>

Version: 2024-02-01

71
papers

28,308
citations

53794

45
h-index

102487

66
g-index

73
all docs

73
docs citations

73
times ranked

21658
citing authors

#	ARTICLE	IF	CITATIONS
1	The TRAR gene classifier to predict response to neoadjuvant therapy in HER2-positive and ER-positive breast cancer patients: an explorative analysis from the NeoSphere trial. <i>Molecular Oncology</i> , 2022, 16, 2355-2366.	4.6	3
2	Customizing local and systemic therapies for women with early breast cancer: the St. Gallen International Consensus Guidelines for treatment of early breast cancer 2021. <i>Annals of Oncology</i> , 2021, 32, 1216-1235.	1.2	354
3	Derived Neutrophil-to-Lymphocyte Ratio Predicts Pathological Complete Response to Neoadjuvant Chemotherapy in Breast Cancer. <i>Frontiers in Oncology</i> , 2021, 11, 827625.	2.8	7
4	Pertuzumab, trastuzumab, and docetaxel for HER2-positive metastatic breast cancer (CLEOPATRA): end-of-study results from a double-blind, randomised, placebo-controlled, phase 3 study. <i>Lancet Oncology</i> , 2020, 21, 519-530.	10.7	441
5	The role of digital mammography, scintimammography with ^{99m}Tc -methoxyisobutylisonitrile (MIBI) and ultrasound in the diagnosis of multicentric breast cancer. <i>Opuholi Zenskoj Reproktivnoj Sistemy</i> , 2020, 15, 12-22.	0.4	4
6	EXPERIENCE OF SENTINEL LYMPH NODE BIOPSY AFTER NEOADJUVANT CHEMOTHERAPY. <i>Voprosy Onkologii</i> , 2020, 66, 370-375.	0.2	0
7	A phase 1b/2, open-label, dose-escalation, and dose-confirmation study of eribulin mesilate in combination with capecitabine. <i>British Journal of Cancer</i> , 2019, 120, 579-586.	6.4	2
8	OPTIMIZATION IN DIEP FLAP BREAST RECONSTRUCTION (DEEP INFERIOR EPIGASTRIC ARTERY PERFORATOR) Tj ETQo 0 0 rgBT /Overlo	0.2	0
9	Comparing Neoadjuvant Nab-paclitaxel vs Paclitaxel Both Followed by Anthracycline Regimens in Women With ERBB2/HER2 -Negative Breast Cancerâ€”The Evaluating Treatment With Neoadjuvant Abraxane (ETNA) Trial. <i>JAMA Oncology</i> , 2018, 4, 302.	7.1	115
10	Phase III, Randomized, Double-Blind Study Comparing the Efficacy, Safety, and Immunogenicity of SB3 (Trastuzumab Biosimilar) and Reference Trastuzumab in Patients Treated With Neoadjuvant Therapy for Human Epidermal Growth Factor Receptor 2-Positive Early Breast Cancer. <i>Journal of Clinical Oncology</i> , 2018, 36, 968-974.	1.6	78
11	CLINICAL AND BIOLOGICAL MODEL FOR EVALUATION OF THE EFFECTIVENESS OF SYSTEMIC THERAPY FOR BREAST CANCER. <i>Voprosy Onkologii</i> , 2018, 64, 289-297.	0.2	0
12	SENTINEL LYMPH NODE BIOPSY IN EARLY BREAST CANCER: THE EXPERIENCE OF THE N.N. PETROV RESEARCH INSTITUTE OF ONCOLOGY. <i>Voprosy Onkologii</i> , 2017, 63, 267-273.	0.2	3
13	5-year analysis of neoadjuvant pertuzumab and trastuzumab in patients with locally advanced, inflammatory, or early-stage HER2-positive breast cancer (NeoSphere): a multicentre, open-label, phase 2 randomised trial. <i>Lancet Oncology</i> , 2016, 17, 791-800.	10.7	623
14	Pertuzumab, Trastuzumab, and Docetaxel in HER2-Positive Metastatic Breast Cancer. <i>New England Journal of Medicine</i> , 2015, 372, 724-734.	27.0	1,658
15	RECIST for Response (Clinical and Imaging) in Neoadjuvant Clinical Trials in Operable Breast Cancer. <i>Journal of the National Cancer Institute Monographs</i> , 2015, 2015, 21-23.	2.1	32
16	Tailoring therapiesâ€”improving the management of early breast cancer: St Gallen International Expert Consensus on the Primary Therapy of Early Breast Cancer 2015. <i>Annals of Oncology</i> , 2015, 26, 1533-1546.	1.2	1,449
17	Patients' preferences for subcutaneous trastuzumab versus conventional intravenous infusion for the adjuvant treatment of HER2-positive early breast cancer: final analysis of 488 patients in the international, randomized, two-cohort PrefHer study. <i>Annals of Oncology</i> , 2014, 25, 1979-1987.	1.2	122
18	Pathological complete response and long-term clinical benefit in breast cancer: the CTNeoBC pooled analysis. <i>Lancet</i> , 2014, 384, 164-172.	13.7	3,224

#	ARTICLE	IF	CITATIONS
19	Challenges to effective cancer control in China, India, and Russia. <i>Lancet Oncology</i> , The, 2014, 15, 489-538.	10.7	411
20	Neoadjuvant and adjuvant trastuzumab in patients with HER2-positive locally advanced breast cancer (NOAH): follow-up of a randomised controlled superiority trial with a parallel HER2-negative cohort. <i>Lancet Oncology</i> , The, 2014, 15, 640-647.	10.7	406
21	Phase III trial of nonpegylated liposomal doxorubicin in combination with trastuzumab and paclitaxel in HER2-positive metastatic breast cancer. <i>Annals of Oncology</i> , 2014, 25, 592-598.	1.2	82
22	Research-Based PAM50 Subtype Predictor Identifies Higher Responses and Improved Survival Outcomes in HER2-Positive Breast Cancer in the NOAH Study. <i>Clinical Cancer Research</i> , 2014, 20, 511-521.	7.0	191
23	Eniluracil Plus 5-Fluorouracil and Leucovorin: Treatment for Metastatic Breast Cancer Patients in Whom Capecitabine Treatment Rapidly Failed. <i>Clinical Breast Cancer</i> , 2014, 14, 26-30.	2.4	6
24	A Comparison of Proposed Biosimilar and Originator Filgrastim for the Prevention of Neutropenia in Patients with Breast Cancer Receiving Myelosuppressive Adjuvant or Neoadjuvant Chemotherapy: Phase III, Randomized, Double-Blind Trial (The PIONEER study). <i>Blood</i> , 2014, 124, 5133-5133.	1.4	4
25	Personalizing the treatment of women with early breast cancer: highlights of the St Gallen International Expert Consensus on the Primary Therapy of Early Breast Cancer 2013. <i>Annals of Oncology</i> , 2013, 24, 2206-2223.	1.2	2,805
26	Treatment of older patients with HER2-positive metastatic breast cancer with pertuzumab, trastuzumab, and docetaxel: subgroup analyses from a randomized, double-blind, placebo-controlled phase III trial (CLEOPATRA). <i>Breast Cancer Research and Treatment</i> , 2013, 142, 89-99.	2.5	62
27	Preference for subcutaneous or intravenous administration of trastuzumab in patients with HER2-positive early breast cancer (PrefHer): an open-label randomised study. <i>Lancet Oncology</i> , The, 2013, 14, 962-970.	10.7	173
28	Pertuzumab, trastuzumab, and docetaxel for HER2-positive metastatic breast cancer (CLEOPATRA) <i>Lancet Oncology</i> , The, 2013, 14, 461-471.	10.7	849
29	AVEREL: A Randomized Phase III Trial Evaluating Bevacizumab in Combination With Docetaxel and Trastuzumab As First-Line Therapy for HER2-Positive Locally Recurrent/Metastatic Breast Cancer. <i>Journal of Clinical Oncology</i> , 2013, 31, 1719-1725.	1.6	247
30	Phase II Study of Bevacizumab in Combination with Trastuzumab and Capecitabine as First-Line Treatment for HER-2-positive Locally Recurrent or Metastatic Breast Cancer. <i>Oncologist</i> , 2012, 17, 469-475.	3.7	48
31	Randomized phase II non-inferiority study (NO16853) of two different doses of capecitabine in combination with docetaxel for locally advanced/metastatic breast cancer. <i>Annals of Oncology</i> , 2012, 23, 589-597.	1.2	13
32	Lapatinib with trastuzumab for HER2-positive early breast cancer (NeoALTTO): a randomised, open-label, multicentre, phase 3 trial. <i>Lancet</i> , The, 2012, 379, 633-640.	13.7	1,165
33	Efficacy and safety of neoadjuvant pertuzumab and trastuzumab in women with locally advanced, inflammatory, or early HER2-positive breast cancer (NeoSphere): a randomised multicentre, open-label, phase 2 trial. <i>Lancet Oncology</i> , The, 2012, 13, 25-32.	10.7	1,879
34	First-line bevacizumab-containing therapy for breast cancer: results in patients aged ≥70 years treated in the ATHENA study. <i>Annals of Oncology</i> , 2012, 23, 111-118.	1.2	49
35	Subcutaneous versus intravenous administration of (neo)adjuvant trastuzumab in patients with HER2-positive, clinical stage III breast cancer (HannaH study): a phase 3, open-label, multicentre, randomised trial. <i>Lancet Oncology</i> , The, 2012, 13, 869-878.	10.7	303
36	High prevalence and breast cancer predisposing role of the BLM c.1642 C>T (Q548X) mutation in Russia. <i>International Journal of Cancer</i> , 2012, 130, 2867-2873.	5.1	58

#	ARTICLE	IF	CITATIONS
37	Endocrine Metabolic Disorders in Patients with Breast Cancer, Carriers of BRCA1 Gene Mutations. <i>Bulletin of Experimental Biology and Medicine</i> , 2012, 152, 610-612.	0.8	2
38	Pathological complete response rates following different neoadjuvant chemotherapy regimens for operable breast cancer according to ER status, in two parallel, randomized phase II trials with an adaptive study design (ECTO II). <i>Breast Cancer Research and Treatment</i> , 2012, 132, 843-851.	2.5	43
39	Recommendations from an International Consensus Conference on the Current Status and Future of Neoadjuvant Systemic Therapy in Primary Breast Cancer. <i>Annals of Surgical Oncology</i> , 2012, 19, 1508-1516.	1.5	401
40	Treatment with trastuzumab for 1 year after adjuvant chemotherapy in patients with HER2-positive early breast cancer: a 4-year follow-up of a randomised controlled trial. <i>Lancet Oncology</i> , The, 2011, 12, 236-244.	10.7	575
41	More favorable progesterone receptor phenotype of breast cancer in diabetics treated with metformin. <i>Medical Oncology</i> , 2011, 28, 1260-1263.	2.5	34
42	Phase III Study of Doxorubicin/Cyclophosphamide With Concomitant Versus Sequential Docetaxel As Adjuvant Treatment in Patients With Human Epidermal Growth Factor Receptor 2â€œNormal, Node-Positive Breast Cancer: BCIRG-005 Trial. <i>Journal of Clinical Oncology</i> , 2011, 29, 3877-3884.	1.6	135
43	Randomized Phase II Trial of Letrozole plus Anti-MUC1 Antibody AS1402 in Hormone Receptorâ€œPositive Locally Advanced or Metastatic Breast Cancer. <i>Clinical Cancer Research</i> , 2011, 17, 6822-6830.	7.0	63
44	Phosphatidylinositol-3-kinase alpha catalytic subunit mutation and response to neoadjuvant endocrine therapy for estrogen receptor positive breast cancer. <i>Breast Cancer Research and Treatment</i> , 2010, 119, 379-390.	2.5	122
45	Neoadjuvant chemotherapy with trastuzumab followed by adjuvant trastuzumab versus neoadjuvant chemotherapy alone, in patients with HER2-positive locally advanced breast cancer (the NOAH trial): a randomised controlled superiority trial with a parallel HER2-negative cohort. <i>Lancet</i> , The, 2010, 375, 377-384.	13.7	1,061
46	A Phase III Randomized Equivalence Study of Biosimilar Filgrastim versus Amgen Filgrastim in Patients Receiving Myelosuppressive Chemotherapy for Breast Cancer. <i>Oncology Research and Treatment</i> , 2010, 33, 504-511.	1.2	45
47	Phase II Randomized Study of Neoadjuvant Everolimus Plus Letrozole Compared With Placebo Plus Letrozole in Patients With Estrogen Receptorâ€œPositive Breast Cancer. <i>Journal of Clinical Oncology</i> , 2009, 27, 2630-2637.	1.6	582
48	Phase III Trial Evaluating the Addition of Paclitaxel to Doxorubicin Followed by Cyclophosphamide, Methotrexate, and Fluorouracil, As Adjuvant or Primary Systemic Therapy: European Cooperative Trial in Operable Breast Cancer. <i>Journal of Clinical Oncology</i> , 2009, 27, 2474-2481.	1.6	194
49	Adjuvant endocrine therapy for perimenopausal women with early breast cancer. <i>Breast</i> , 2009, 18, 2-7.	2.2	14
50	Large family with both parents affected by distinct BRCA1 mutations: implications for genetic testing. <i>Hereditary Cancer in Clinical Practice</i> , 2009, 7, 2.	1.5	3
51	A phase II, randomized, blinded study of the farnesyltransferase inhibitor tipifarnib combined with letrozole in the treatment of advanced breast cancer after antiestrogen therapy. <i>Breast Cancer Research and Treatment</i> , 2008, 110, 327-335.	2.5	60
52	Need for global action for cancer control. <i>Annals of Oncology</i> , 2008, 19, 1519-1521.	1.2	29
53	Recommendations from an international expert panel on the use of neoadjuvant (primary) systemic treatment of operable breast cancer: new perspectives 2006. <i>Annals of Oncology</i> , 2007, 18, 1927-1934.	1.2	324
54	Signs of proinflammatory/genotoxic switch (adipogenotoxicosis) in mammary fat of breast cancer patients: Role of menopausal status, estrogens and hyperglycemia. <i>International Journal of Cancer</i> , 2007, 121, 514-519.	5.1	22

#	ARTICLE	IF	CITATIONS
55	Phase 2 randomized trial of primary endocrine therapy versus chemotherapy in postmenopausal patients with estrogen receptor-positive breast cancer. <i>Cancer</i> , 2007, 110, 244-254.	4.1	273
56	Founder mutations in early-onset, familial and bilateral breast cancer patients from Russia. <i>Familial Cancer</i> , 2007, 6, 281-286.	1.9	67
57	CHEK2 1100delC mutation is frequent among Russian breast cancer patients. <i>Breast Cancer Research and Treatment</i> , 2006, 100, 99-102.	2.5	32
58	Phase III Trial Comparing Three Doses of Docetaxel for Second-Line Treatment of Advanced Breast Cancer. <i>Journal of Clinical Oncology</i> , 2006, 24, 4963-4970.	1.6	147
59	Recommendations From an International Expert Panel on the Use of Neoadjuvant (Primary) Systemic Treatment of Operable Breast Cancer: An Update. <i>Journal of Clinical Oncology</i> , 2006, 24, 1940-1949.	1.6	579
60	NBS1 657del5 mutation may contribute only to a limited fraction of breast cancer cases in Russia. <i>International Journal of Cancer</i> , 2005, 114, 585-589.	5.1	59
61	Maintaining Normal Hemoglobin Levels With Epoetin Alfa in Mainly Nonanemic Patients With Metastatic Breast Cancer Receiving First-Line Chemotherapy: A Survival Study. <i>Journal of Clinical Oncology</i> , 2005, 23, 5960-5972.	1.6	622
62	Feasibility and Tolerability of Sequential Doxorubicin/Paclitaxel Followed by Cyclophosphamide, Methotrexate, and Fluorouracil and Its Effects on Tumor Response as Preoperative Therapy. <i>Clinical Cancer Research</i> , 2005, 11, 8715-8721.	7.0	146
63	Effects of radiotherapy and of differences in the extent of surgery for early breast cancer on local recurrence and 15-year survival: an overview of the randomised trials. <i>Lancet</i> , The, 2005, 366, 2087-2106.	13.7	4,596
64	Phase II Study of Temozolomide (CC-779), a Novel Inhibitor of mTOR, in Heavily Pretreated Patients With Locally Advanced or Metastatic Breast Cancer. <i>Journal of Clinical Oncology</i> , 2005, 23, 5314-5322.	1.6	461
65	Expression of estrogen receptors- α and β in primary breast neoplasms and tumors exposed to neoadjuvant hormonal therapy. <i>Bulletin of Experimental Biology and Medicine</i> , 2004, 138, 494-496.	0.8	4
66	Expression of Estrogen Receptors- α and Aromatase Activity in Primary Mammary Gland Tumors. <i>Bulletin of Experimental Biology and Medicine</i> , 2003, 136, 487-489.	0.8	0
67	Androgen administration in middle-aged and ageing men: effects of oral testosterone undecanoate on dihydrotestosterone, oestradiol and prostate volume. <i>Journal of Developmental and Physical Disabilities</i> , 2002, 25, 119-125.	3.6	56
68	Preoperative treatment of postmenopausal breast cancer patients with letrozole: A randomized double-blind multicenter study. <i>Annals of Oncology</i> , 2001, 12, 1527-1532.	1.2	617
69	N3 category in the TNM classification of breast cancer needs to be defined more precisely. <i>Breast</i> , 1997, 6, 217-220.	2.2	0
70	Aromatase in breast cancer tissue ? localization and relationship with reproductive status of patients. <i>Journal of Cancer Research and Clinical Oncology</i> , 1996, 122, 495-498.	2.5	26
71	Activity of aromatase in breast cancer tissue: Role of the cell substrate. <i>Bulletin of Experimental Biology and Medicine</i> , 1995, 120, 1042-1045.	0.8	0