

Margreet Ten Have

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7222148/publications.pdf>

Version: 2024-02-01

55
papers

4,287
citations

147801

31
h-index

149698

56
g-index

56
all docs

56
docs citations

56
times ranked

6232
citing authors

#	ARTICLE	IF	CITATIONS
1	Anxiety Disorders and Risk for Suicidal Ideation and Suicide Attempts. Archives of General Psychiatry, 2005, 62, 1249.	12.3	664
2	Prevalence of mental disorders and trends from 1996 to 2009. Results from the Netherlands Mental Health Survey and Incidence Study-2. Social Psychiatry and Psychiatric Epidemiology, 2012, 47, 203-213.	3.1	325
3	Screening for mood and anxiety disorders with the five-item, the three-item, and the two-item Mental Health Inventory. Psychiatry Research, 2009, 168, 250-255.	3.3	264
4	The cross-national epidemiology of social anxiety disorder: Data from the World Mental Health Survey Initiative. BMC Medicine, 2017, 15, 143.	5.5	258
5	Are attitudes towards mental health help-seeking associated with service use? Results from the European Study of Epidemiology of Mental Disorders. Social Psychiatry and Psychiatric Epidemiology, 2010, 45, 153-163.	3.1	212
6	Incidence and Course of Suicidal Ideation and Suicide Attempts in the General Population. Canadian Journal of Psychiatry, 2009, 54, 824-833.	1.9	198
7	Bipolar disorder in the general population in The Netherlands (prevalence, consequences and care) Tj ETQq1 1 0.784314 rgBT /Overlo Journal of Affective Disorders, 2002, 68, 203-213.	4.1	175
8	Childhood adversities and risk for suicidal ideation and attempts: a longitudinal population-based study. Psychological Medicine, 2006, 36, 1769-1778.	4.5	168
9	The Netherlands Mental Health Survey and Incidence Study (NEMESIS): design and methods. International Journal of Methods in Psychiatric Research, 2010, 19, 125-141.	2.1	165
10	Childhood adversities and post-traumatic stress disorder: evidence for stress sensitisation in the World Mental Health Surveys. British Journal of Psychiatry, 2017, 211, 280-288.	2.8	118
11	The disease burden of childhood adversities in adults: A population-based study. Child Abuse and Neglect, 2011, 35, 937-945.	2.6	101
12	Physical exercise in adults and mental health status. Journal of Psychosomatic Research, 2011, 71, 342-348.	2.6	86
13	Phenotypically Continuous With Clinical Psychosis, Discontinuous in Need for Care: Evidence for an Extended Psychosis Phenotype. Schizophrenia Bulletin, 2012, 38, 231-238.	4.3	85
14	Sociodemographic and psychiatric predictors of attrition in a prospective psychiatric epidemiological study among the general population. Result of the Netherlands Mental Health Survey and Incidence Study-2. Comprehensive Psychiatry, 2013, 54, 1131-1139.	3.1	79
15	Predictors of suicidality in depressive spectrum disorders in the general population: results of the Netherlands Mental Health Survey and Incidence Study. Social Psychiatry and Psychiatric Epidemiology, 2010, 45, 513-521.	3.1	78
16	Prevalence of bipolar disorder in the general population: a Reappraisal Study of the Netherlands Mental Health Survey and Incidence Study. Acta Psychiatrica Scandinavica, 2004, 110, 374-382.	4.5	77
17	The association between type and number of adverse working conditions and mental health during a time of economic crisis (2010-2012). Social Psychiatry and Psychiatric Epidemiology, 2015, 50, 899-907.	3.1	72
18	Recurrence and chronicity of major depressive disorder and their risk indicators in a population cohort. Acta Psychiatrica Scandinavica, 2018, 137, 503-515.	4.5	71

#	ARTICLE	IF	CITATIONS
19	Prevalence rates of borderline personality disorder symptoms: a study based on the Netherlands Mental Health Survey and Incidence Study-2. <i>BMC Psychiatry</i> , 2016, 16, 249.	2.6	66
20	Does neuroticism explain variations in care service use for mental health problems in the general population?. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2005, 40, 425-431.	3.1	61
21	Does educational background explain inequalities in care service use for mental health problems in the Dutch general population?. <i>Acta Psychiatrica Scandinavica</i> , 2003, 107, 178-187.	4.5	59
22	Course of subthreshold depression into a depressive disorder and its risk factors. <i>Journal of Affective Disorders</i> , 2018, 241, 206-215.	4.1	55
23	Combined effect of mental disorder and low social support on care service use for mental health problems in the Dutch general population. <i>Psychological Medicine</i> , 2002, 32, 311-323.	4.5	52
24	Common mental disorder severity and its association with treatment contact and treatment intensity for mental health problems. <i>Psychological Medicine</i> , 2013, 43, 2203-2213.	4.5	52
25	Duration of major and minor depressive episodes and associated risk indicators in a psychiatric epidemiological cohort study of the general population. <i>Acta Psychiatrica Scandinavica</i> , 2017, 136, 300-312.	4.5	52
26	Prevalence and course of subthreshold anxiety disorder in the general population: A three-year follow-up study. <i>Journal of Affective Disorders</i> , 2019, 247, 105-113.	4.1	52
27	The identification of symptom-based subtypes of depression: A nationally representative cohort study. <i>Journal of Affective Disorders</i> , 2016, 190, 395-406.	4.1	50
28	Prevalence and risk factors for first onset of suicidal behaviors in the Netherlands Mental Health Survey and Incidence Study-2. <i>Journal of Affective Disorders</i> , 2013, 147, 205-211.	4.1	49
29	What depressive symptoms are associated with the use of care services?. <i>Journal of Affective Disorders</i> , 2004, 80, 239-248.	4.1	46
30	The bidirectional relationship between loneliness and common mental disorders in adults: findings from a longitudinal population-based cohort study. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2020, 55, 1297-1310.	3.1	46
31	Recovery from mood and anxiety disorders: The influence of positive mental health. <i>Journal of Affective Disorders</i> , 2019, 252, 107-113.	4.1	39
32	The interplay between emotional exhaustion, common mental disorders, functioning and health care use in the working population. <i>Journal of Psychosomatic Research</i> , 2017, 100, 8-14.	2.6	32
33	Predictors of incident care service utilisation for mental health problems in the Dutch general population. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2001, 36, 141-149.	3.1	30
34	The longitudinal association between lifetime mental disorders and first onset or recurrent suicide ideation. <i>BMC Psychiatry</i> , 2019, 19, 345.	2.6	30
35	Psychotic experiences and incident suicidal ideation and behaviour: Disentangling the longitudinal associations from connected psychopathology. <i>Psychiatry Research</i> , 2016, 245, 267-275.	3.3	25
36	Differences in alcohol use between younger and older people: Results from a general population study. <i>Drug and Alcohol Dependence</i> , 2019, 202, 18-23.	3.2	25

#	ARTICLE	IF	CITATIONS
37	Demographic and need factors of early, delayed and no mental health care use in major depression: a prospective study. <i>BMC Psychiatry</i> , 2017, 17, 367.	2.6	24
38	Borderline personality symptoms and work performance: a population-based survey. <i>BMC Psychiatry</i> , 2018, 18, 202.	2.6	23
39	Inequalities in mental health care and social services utilisation by immigrant women. <i>European Journal of Public Health</i> , 1999, 9, 45-51.	0.3	22
40	Emotional disorders among informal caregivers in the general population: target groups for prevention. <i>BMC Psychiatry</i> , 2015, 15, 23.	2.6	22
41	Insomnia among current and remitted common mental disorders and the association with role functioning: results from a general population study. <i>Sleep Medicine</i> , 2016, 25, 34-41.	1.6	21
42	The Relationship Between Excessive Alcohol Consumption and Alcohol Use Disorders According to DSM-IV and DSM-5. <i>Alcoholism: Clinical and Experimental Research</i> , 2014, 38, 249-256.	2.4	19
43	Distribution of ADHD symptoms, and associated comorbidity, exposure to risk factors and disability: Results from a general population study. <i>Psychiatry Research</i> , 2018, 267, 256-265.	3.3	18
44	Low self-recognition and awareness of past hypomanic and manic episodes in the general population. <i>International Journal of Bipolar Disorders</i> , 2015, 3, 22.	2.2	15
45	Childhood maltreatment, vulnerability characteristics and adult incident common mental disorders: 3-year longitudinal data among >10,000 adults in the general population. <i>Journal of Psychiatric Research</i> , 2019, 113, 199-207.	3.1	13
46	Recurrence of anxiety disorders and its predictors in the general population. <i>Psychological Medicine</i> , 2023, 53, 1334-1342.	4.5	13
47	Associations of work and health-related characteristics with intention to continue working after the age of 65 years. <i>European Journal of Public Health</i> , 2015, 25, 122-124.	0.3	12
48	Explaining service use for mental health problems in the Dutch general population: the role of resources, emotional disorder and functional impairment. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2006, 41, 285-293.	3.1	11
49	Duration of anxiety disorder and its associated risk indicators: Results of a longitudinal study of the general population. <i>Depression and Anxiety</i> , 2021, 38, 328-336.	4.1	11
50	The Bidirectional Relationship Between Debts and Common Mental Disorders: Results of a longitudinal Population-Based Study. <i>Administration and Policy in Mental Health and Mental Health Services Research</i> , 2021, 48, 810-820.	2.1	11
51	The long-lasting impact of childhood trauma on adult chronic physical disorders. <i>Journal of Psychiatric Research</i> , 2021, 136, 87-94.	3.1	9
52	Upward trends in the use of community mental health and social work services in the Netherlands between 1979 and 1995: are particular sociodemographic groups responsible?. <i>Acta Psychiatrica Scandinavica</i> , 2003, 108, 447-454.	4.5	6
53	Associations between DSM-IV mental disorders and subsequent onset of arthritis. <i>Journal of Psychosomatic Research</i> , 2016, 82, 11-16.	2.6	6
54	Demand for Mental Health Care and Changes in Service Use Patterns in the Netherlands, 1979 to 1995. <i>Psychiatric Services</i> , 2005, 56, 1409-1415.	2.0	5

#	ARTICLE	IF	CITATIONS
55	Impact of borderline personality disorder traits on the association between age and health-related quality of life: A cohort study in the general population. <i>European Psychiatry</i> , 2021, 64, e33.	0.2	5