

# Margreet Ten Have

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7222148/publications.pdf>

Version: 2024-02-01

55  
papers

4,287  
citations

147801

31  
h-index

149698

56  
g-index

56  
all docs

56  
docs citations

56  
times ranked

6232  
citing authors

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 1  | Recurrence of anxiety disorders and its predictors in the general population. <i>Psychological Medicine</i> , 2023, 53, 1334-1342.  | 4.5 | 13        |
| 2  | Duration of anxiety disorder and its associated risk indicators: Results of a longitudinal study of the general population. <i>Depression and Anxiety</i> , 2021, 38, 328-336.  | 4.1 | 11        |
| 3  | The Bidirectional Relationship Between Debts and Common Mental Disorders: Results of a longitudinal Population-Based Study. <i>Administration and Policy in Mental Health and Mental Health Services Research</i> , 2021, 48, 810-820.  | 2.1 | 11        |
| 4  | The long-lasting impact of childhood trauma on adult chronic physical disorders. <i>Journal of Psychiatric Research</i> , 2021, 136, 87-94.   | 3.1 | 9         |
| 5  | Impact of borderline personality disorder traits on the association between age and health-related quality of life: A cohort study in the general population. <i>European Psychiatry</i> , 2021, 64, e33.                               | 0.2 | 5         |
| 6  | The bidirectional relationship between loneliness and common mental disorders in adults: findings from a longitudinal population-based cohort study. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2020, 55, 1297-1310.       | 3.1 | 46        |
| 7  | Differences in alcohol use between younger and older people: Results from a general population study. <i>Drug and Alcohol Dependence</i> , 2019, 202, 18-23.  | 3.2 | 25        |
| 8  | The longitudinal association between lifetime mental disorders and first onset or recurrent suicide ideation. <i>BMC Psychiatry</i> , 2019, 19, 345.  | 2.6 | 30        |
| 9  | Recovery from mood and anxiety disorders: The influence of positive mental health. <i>Journal of Affective Disorders</i> , 2019, 252, 107-113.  | 4.1 | 39        |
| 10 | Childhood maltreatment, vulnerability characteristics and adult incident common mental disorders: 3-year longitudinal data among >10,000 adults in the general population. <i>Journal of Psychiatric Research</i> , 2019, 113, 199-207. | 3.1 | 13        |
| 11 | Prevalence and course of subthreshold anxiety disorder in the general population: A three-year follow-up study. <i>Journal of Affective Disorders</i> , 2019, 247, 105-113.   | 4.1 | 52        |
| 12 | Recurrence and chronicity of major depressive disorder and their risk indicators in a population cohort. <i>Acta Psychiatrica Scandinavica</i> , 2018, 137, 503-515.  | 4.5 | 71        |
| 13 | Borderline personality symptoms and work performance: a population-based survey. <i>BMC Psychiatry</i> , 2018, 18, 202.   | 2.6 | 23        |
| 14 | Course of subthreshold depression into a depressive disorder and its risk factors. <i>Journal of Affective Disorders</i> , 2018, 241, 206-215.  | 4.1 | 55        |
| 15 | Distribution of ADHD symptoms, and associated comorbidity, exposure to risk factors and disability: Results from a general population study. <i>Psychiatry Research</i> , 2018, 267, 256-265.   | 3.3 | 18        |
| 16 | Duration of major and minor depressive episodes and associated risk indicators in a psychiatric epidemiological cohort study of the general population. <i>Acta Psychiatrica Scandinavica</i> , 2017, 136, 300-312.                     | 4.5 | 52        |
| 17 | Childhood adversities and post-traumatic stress disorder: evidence for stress sensitisation in the World Mental Health Surveys. <i>British Journal of Psychiatry</i> , 2017, 211, 280-288.  | 2.8 | 118       |
| 18 | The interplay between emotional exhaustion, common mental disorders, functioning and health care use in the working population. <i>Journal of Psychosomatic Research</i> , 2017, 100, 8-14.   | 2.6 | 32        |

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 19 | The cross-national epidemiology of social anxiety disorder: Data from the World Mental Health Survey Initiative. <i>BMC Medicine</i> , 2017, 15, 143.  | 5.5 | 258       |
| 20 | Demographic and need factors of early, delayed and no mental health care use in major depression: a prospective study. <i>BMC Psychiatry</i> , 2017, 17, 367.  | 2.6 | 24        |
| 21 | Insomnia among current and remitted common mental disorders and the association with role functioning: results from a general population study. <i>Sleep Medicine</i> , 2016, 25, 34-41.   | 1.6 | 21        |
| 22 | Psychotic experiences and incident suicidal ideation and behaviour: Disentangling the longitudinal associations from connected psychopathology. <i>Psychiatry Research</i> , 2016, 245, 267-275.   | 3.3 | 25        |
| 23 | Prevalence rates of borderline personality disorder symptoms: a study based on the Netherlands Mental Health Survey and Incidence Study-2. <i>BMC Psychiatry</i> , 2016, 16, 249.  | 2.6 | 66        |
| 24 | Associations between DSM-IV mental disorders and subsequent onset of arthritis. <i>Journal of Psychosomatic Research</i> , 2016, 82, 11-16.  | 2.6 | 6         |
| 25 | The identification of symptom-based subtypes of depression: A nationally representative cohort study. <i>Journal of Affective Disorders</i> , 2016, 190, 395-406.  | 4.1 | 50        |
| 26 | Low self-recognition and awareness of past hypomanic and manic episodes in the general population. <i>International Journal of Bipolar Disorders</i> , 2015, 3, 22.  | 2.2 | 15        |
| 27 | Associations of work and health-related characteristics with intention to continue working after the age of 65 years. <i>European Journal of Public Health</i> , 2015, 25, 122-124.  | 0.3 | 12        |
| 28 | The association between type and number of adverse working conditions and mental health during a time of economic crisis (2010-2012). <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2015, 50, 899-907.   | 3.1 | 72        |
| 29 | Emotional disorders among informal caregivers in the general population: target groups for prevention. <i>BMC Psychiatry</i> , 2015, 15, 23.   | 2.6 | 22        |
| 30 | The Relationship Between Excessive Alcohol Consumption and Alcohol Use Disorders According to DSM-IV and DSM-5. <i>Alcoholism: Clinical and Experimental Research</i> , 2014, 38, 249-256.   | 2.4 | 19        |
| 31 | Prevalence and risk factors for first onset of suicidal behaviors in the Netherlands Mental Health Survey and Incidence Study-2. <i>Journal of Affective Disorders</i> , 2013, 147, 205-211.   | 4.1 | 49        |
| 32 | Sociodemographic and psychiatric predictors of attrition in a prospective psychiatric epidemiological study among the general population. Result of the Netherlands Mental Health Survey and Incidence Study-2. <i>Comprehensive Psychiatry</i> , 2013, 54, 1131-1139. | 3.1 | 79        |
| 33 | Common mental disorder severity and its association with treatment contact and treatment intensity for mental health problems. <i>Psychological Medicine</i> , 2013, 43, 2203-2213.  | 4.5 | 52        |
| 34 | Phenotypically Continuous With Clinical Psychosis, Discontinuous in Need for Care: Evidence for an Extended Psychosis Phenotype. <i>Schizophrenia Bulletin</i> , 2012, 38, 231-238.  | 4.3 | 85        |
| 35 | Prevalence of mental disorders and trends from 1996 to 2009. Results from the Netherlands Mental Health Survey and Incidence Study-2. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2012, 47, 203-213.   | 3.1 | 325       |
| 36 | Physical exercise in adults and mental health status. <i>Journal of Psychosomatic Research</i> , 2011, 71, 342-348.  | 2.6 | 86        |

| #  | ARTICLE   | IF   | CITATIONS |
|----|---|------|-----------|
| 37 | The disease burden of childhood adversities in adults: A population-based study. <i>Child Abuse and Neglect</i> , 2011, 35, 937-945.  | 2.6  | 101       |
| 38 | Are attitudes towards mental health help-seeking associated with service use? Results from the European Study of Epidemiology of Mental Disorders. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2010, 45, 153-163.           | 3.1  | 212       |
| 39 | Predictors of suicidality in depressive spectrum disorders in the general population: results of the Netherlands Mental Health Survey and Incidence Study. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2010, 45, 513-521.   | 3.1  | 78        |
| 40 | The Netherlands Mental Health Survey and Incidence Study (NEMESIS): design and methods. <i>International Journal of Methods in Psychiatric Research</i> , 2010, 19, 125-141.  | 2.1  | 165       |
| 41 | Screening for mood and anxiety disorders with the five-item, the three-item, and the two-item Mental Health Inventory. <i>Psychiatry Research</i> , 2009, 168, 250-255.   | 3.3  | 264       |
| 42 | Incidence and Course of Suicidal Ideation and Suicide Attempts in the General Population. <i>Canadian Journal of Psychiatry</i> , 2009, 54, 824-833.  | 1.9  | 198       |
| 43 | Childhood adversities and risk for suicidal ideation and attempts: a longitudinal population-based study. <i>Psychological Medicine</i> , 2006, 36, 1769-1778.  | 4.5  | 168       |
| 44 | Explaining service use for mental health problems in the Dutch general population: the role of resources, emotional disorder and functional impairment. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2006, 41, 285-293.      | 3.1  | 11        |
| 45 | Does neuroticism explain variations in care service use for mental health problems in the general population?. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2005, 40, 425-431.   | 3.1  | 61        |
| 46 | Anxiety Disorders and Risk for Suicidal Ideation and Suicide Attempts. <i>Archives of General Psychiatry</i> , 2005, 62, 1249.  | 12.3 | 664       |
| 47 | Demand for Mental Health Care and Changes in Service Use Patterns in the Netherlands, 1979 to 1995. <i>Psychiatric Services</i> , 2005, 56, 1409-1415.  | 2.0  | 5         |
| 48 | Prevalence of bipolar disorder in the general population: a Reappraisal Study of the Netherlands Mental Health Survey and Incidence Study. <i>Acta Psychiatrica Scandinavica</i> , 2004, 110, 374-382.                                  | 4.5  | 77        |
| 49 | What depressive symptoms are associated with the use of care services?. <i>Journal of Affective Disorders</i> , 2004, 80, 239-248.  | 4.1  | 46        |
| 50 | Does educational background explain inequalities in care service use for mental health problems in the Dutch general population?. <i>Acta Psychiatrica Scandinavica</i> , 2003, 107, 178-187.   | 4.5  | 59        |
| 51 | Upward trends in the use of community mental health and social work services in the Netherlands between 1979 and 1995: are particular sociodemographic groups responsible?. <i>Acta Psychiatrica Scandinavica</i> , 2003, 108, 447-454. | 4.5  | 6         |
| 52 | Combined effect of mental disorder and low social support on care service use for mental health problems in the Dutch general population. <i>Psychological Medicine</i> , 2002, 32, 311-323.  | 4.5  | 52        |
| 53 | Bipolar disorder in the general population in The Netherlands (prevalence, consequences and care) Tj ETQq1 1 0.784314 rgBT /Overlo<br><i>Journal of Affective Disorders</i> , 2002, 68, 203-213.  | 4.1  | 175       |
| 54 | Predictors of incident care service utilisation for mental health problems in the Dutch general population. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2001, 36, 141-149.  | 3.1  | 30        |

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 55 | Inequalities in mental health care and social services utilisation by immigrant women. <i>European Journal of Public Health</i> , 1999, 9, 45-51. | 0.3 | 22        |