

Sho-Ichi Yamagishi

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7229750/publications.pdf>

Version: 2024-02-01

379
papers

23,881
citations

6613

79
h-index

10445

139
g-index

379
all docs

379
docs citations

379
times ranked

20070
citing authors

#	ARTICLE	IF	CITATIONS
1	Predictive ability of serum advanced glycation end products at 11 to 13 weeks of gestation for early-onset preeclampsia. <i>AJOG Global Reports</i> , 2022, 2, 100052.	1.0	1
2	Luseogliflozin inhibits high glucose-induced TGF- β 2 expression in mouse cardiomyocytes by suppressing NHE-1 activity. <i>Journal of International Medical Research</i> , 2022, 50, 030006052210974.	1.0	8
3	Casein Hydrolysate Containing Milk-Derived Peptides Reduces Facial Pigmentation Partly by Decreasing Advanced Glycation End Products in the Skin: A Randomized Double-Blind Placebo-Controlled Trial. <i>Rejuvenation Research</i> , 2021, 24, 97-103.	1.8	3
4	DNA aptamer raised against receptor for advanced glycation end products suppresses renal tubular damage and improves insulin resistance in diabetic mice. <i>Diabetes and Vascular Disease Research</i> , 2021, 18, 147916412199053.	2.0	5
5	Glucose Variability is Independently Correlated with Serum Level of Pigment Epithelium-Derived Factor in Type 2 Diabetes. <i>Diabetes Therapy</i> , 2021, 12, 827-842.	2.5	2
6	Pigment epithelium-derived factor inhibits advanced glycation end product-induced proliferation, VEGF and MMP-9 expression in breast cancer cells via interaction with laminin receptor. <i>Oncology Letters</i> , 2021, 22, 629.	1.8	9
7	Glucose-Dependent Insulinotropic Polypeptide Suppresses Foam Cell Formation of Macrophages through Inhibition of the Cyclin-Dependent Kinase 5-CD36 Pathway. <i>Biomedicines</i> , 2021, 9, 832.	3.2	7
8	DNA-Aptamer Raised against Receptor for Advanced Glycation End Products Improves Survival Rate in Septic Mice. <i>Oxidative Medicine and Cellular Longevity</i> , 2021, 2021, 1-20.	4.0	3
9	Effects of omarigliptin on glucose variability and oxidative stress in type 2 diabetes patients: A prospective study. <i>Diabetes Research and Clinical Practice</i> , 2021, 179, 108999.	2.8	4
10	Increased Urinary Levels of Pentosidine Measured by a Newly Developed Enzyme-Linked Immunosorbent Assay Are Independently Correlated with Fracture After Fall. <i>Rejuvenation Research</i> , 2021, 24, 449-455.	1.8	0
11	Glucose-dependent insulinotropic polypeptide inhibits cardiac hypertrophy and fibrosis in diabetic mice via suppression of TGF- β 2. <i>Diabetes and Vascular Disease Research</i> , 2021, 18, 147916412199903.	2.0	4
12	Two-hour postload plasma glucose and pigment epithelium-derived factor levels are markers of coronary artery inflammation in type 2 diabetic patients. <i>Journal of Nuclear Cardiology</i> , 2020, 27, 1352-1364.	2.1	5
13	Oral administration of spa-derived green alga improves insulin resistance in overweight subjects: Mechanistic insights from fructose-fed rats. <i>Pharmacological Research</i> , 2020, 152, 104633.	7.1	2
14	Association of advanced glycation end products with sarcopenia and frailty in chronic kidney disease. <i>Scientific Reports</i> , 2020, 10, 17647.	3.3	37
15	A Dipeptidyl Peptidase-4 Inhibitor Inhibits Foam Cell Formation of Macrophages in Type 1 Diabetes via Suppression of CD36 and ACAT-1 Expression. <i>International Journal of Molecular Sciences</i> , 2020, 21, 4811.	4.1	20
16	AGE-RAGE Axis Stimulates Oxidized LDL Uptake into Macrophages through Cyclin-Dependent Kinase 5-CD36 Pathway via Oxidative Stress Generation. <i>International Journal of Molecular Sciences</i> , 2020, 21, 9263.	4.1	11
17	Butanolic Extract of Noni Inhibits Proliferation, Inflammation, and Proprotein Convertase Subtilisin Kexin Type 9 (PCSK9) Expression in Cultured Smooth Muscle Cells. <i>Natural Product Communications</i> , 2020, 15, 1934578X2093203.	0.5	0
18	Pathological Role of Receptor for Advanced Glycation End Products in Calcified Aortic Valve Stenosis. <i>Journal of the American Heart Association</i> , 2020, 9, e015261.	3.7	12

#	ARTICLE	IF	CITATIONS
19	GIP as a Potential Therapeutic Target for Atherosclerotic Cardiovascular Disease—A Systematic Review. <i>International Journal of Molecular Sciences</i> , 2020, 21, 1509.	4.1	29
20	Glyceraldehyde-Derived Pyridinium Evokes Renal Tubular Cell Damage via RAGE Interaction. <i>International Journal of Molecular Sciences</i> , 2020, 21, 2604.	4.1	5
21	Brown adipose tissue activation in severe heart failure. <i>European Heart Journal</i> , 2020, 41, 2415-2415.	2.2	3
22	Albuminuria—Lowering effect of sodium—glucose cotransporter 2 inhibitors could be partly attributable to the attenuation of tubular damage in type 2 diabetic patients. <i>Diabetes/Metabolism Research and Reviews</i> , 2020, 36, e3327.	4.0	2
23	Fructose causes endothelial cell damage via activation of advanced glycation end products—receptor system. <i>Diabetes and Vascular Disease Research</i> , 2019, 16, 556-561.	2.0	11
24	Role of Advanced Glycation Endproduct (AGE)-Receptor for Advanced Glycation Endproduct (RAGE) Axis in Cardiovascular Disease and Its Therapeutic Intervention. <i>Circulation Journal</i> , 2019, 83, 1822-1828.	1.6	39
25	Glycolaldehyde-modified advanced glycation end-products inhibit differentiation of human monocytes into osteoclasts via upregulation of IL-10. <i>Bone</i> , 2019, 128, 115034.	2.9	26
26	Relationship between glucose variability evaluated by continuous glucose monitoring and clinical factors, including glucagon-stimulated insulin secretion in patients with type 2 diabetes. <i>Diabetes Research and Clinical Practice</i> , 2019, 158, 107904.	2.8	9
27	Long-Term Local Injection of RAGE-Aptamer Suppresses the Growth of Malignant Melanoma in Nude Mice. <i>Journal of Oncology</i> , 2019, 2019, 1-10.	1.3	11
28	Concerns about clinical efficacy and safety of warfarin in diabetic patients with atrial fibrillation. <i>Cardiovascular Diabetology</i> , 2019, 18, 12.	6.8	16
29	Serum Levels of Protein-Bound Methylglyoxal-Derived Hydroimidazolone-1 are Independently Correlated with Asymmetric Dimethylarginine. <i>Rejuvenation Research</i> , 2019, 22, 431-438.	1.8	6
30	Therapeutic Potential of Pigment Epithelium-derived Factor in Cancer. <i>Current Pharmaceutical Design</i> , 2019, 25, 313-324.	1.9	8
31	Pathological Role of Advanced Glycation End Products (AGEs) and their Receptor Axis in Atrial Fibrillation. <i>Mini-Reviews in Medicinal Chemistry</i> , 2019, 19, 1040-1048.	2.4	9
32	Switching Dipeptidyl Peptidase-4 Inhibitors to Tofogliflozin, a Selective Inhibitor of Sodium-Glucose Cotransporter 2 Improve Arterial Stiffness Evaluated by Cardio-Ankle Vascular Index in Patients with Type 2 Diabetes: A Pilot Study. <i>Current Vascular Pharmacology</i> , 2019, 17, 411-420.	1.7	22
33	Endothelial dysfunction as a common soil of lower urinary tract symptoms and cardiovascular disease. <i>International Journal of Cardiology</i> , 2018, 261, 209-210.	1.7	1
34	Role of Ligands of Receptor for Advanced Glycation End Products (RAGE) in Peripheral Artery Disease. <i>Rejuvenation Research</i> , 2018, 21, 456-463.	1.8	20
35	RAGE-aptamer attenuates deoxycorticosterone acetate/salt-induced renal injury in mice. <i>Scientific Reports</i> , 2018, 8, 2686.	3.3	24
36	Association of advanced glycation end products, evaluated by skin autofluorescence, with lifestyle habits in a general Japanese population. <i>Journal of International Medical Research</i> , 2018, 46, 1043-1051.	1.0	34

#	ARTICLE	IF	CITATIONS
37	Diabetes and Female Sterility/Infertility. , 2018, , 177-183.		3
38	Diabetes and Advanced Glycation End Products. , 2018, , 201-212.		6
39	Role of Hyperglycemia-Induced Advanced Glycation End Product (AGE) Accumulation in Atherosclerosis. <i>Annals of Vascular Diseases</i> , 2018, 11, 253-258.	0.5	48
40	Are Finger Skin Fluorophores Other Than Advanced Glycation End Products (AGEs) Associated With Impaired Musculoskeletal Properties?. <i>Journals of Gerontology - Series A Biological Sciences and Medical Sciences</i> , 2018, 75, 401-402.	3.6	1
41	Aqueous Extract of Glucoraphanin-Rich Broccoli Sprouts Inhibits Formation of Advanced Glycation End Products and Attenuates Inflammatory Reactions in Endothelial Cells. <i>Evidence-based Complementary and Alternative Medicine</i> , 2018, 2018, 1-6.	1.2	15
42	Clinical markers associated with glycaemic response to dipeptidyl peptidase-4 inhibitor therapy. <i>Diabetes/Metabolism Research and Reviews</i> , 2018, 34, e3024.	4.0	2
43	An Overview on Diabetic Nephropathy. , 2018, , 125-137.		0
44	Advanced glycation end products evoke inflammatory reactions in proximal tubular cells via autocrine production of dipeptidyl peptidase-4. <i>Microvascular Research</i> , 2018, 120, 90-93.	2.5	18
45	Association of skin autofluorescence with plaque vulnerability evaluated by optical coherence tomography in patients with cardiovascular disease. <i>Atherosclerosis</i> , 2018, 274, 47-53.	0.8	12
46	Sex disparity in cardiovascular mortality rates associated with diabetes. <i>Diabetes/Metabolism Research and Reviews</i> , 2018, 34, e3059.	4.0	3
47	Therapeutic Potential of DNA-aptamers Raised Against AGE-RAGE Axis in Diabetes-related Complications. <i>Current Pharmaceutical Design</i> , 2018, 24, 2802-2809.	1.9	25
48	Ratio of serum levels of AGEs to soluble RAGE is correlated with trimethylamine-N-oxide in non-diabetic subjects. <i>International Journal of Food Sciences and Nutrition</i> , 2017, 68, 1013-1020.	2.8	9
49	Pigment Epithelium-Derived Factor (PEDF) Prevents Hepatic Fat Storage, Inflammation, and Fibrosis in Dietary Steatohepatitis of Mice. <i>Digestive Diseases and Sciences</i> , 2017, 62, 1527-1536.	2.3	21
50	RAGE-Aptamer Blocks the Development and Progression of Experimental Diabetic Nephropathy. <i>Diabetes</i> , 2017, 66, 1683-1695.	0.6	91
51	High serum soluble tumor necrosis factor receptor 1 predicts poor treatment response in acute-stage schizophrenia. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2017, 76, 145-154.	4.8	11
52	N-butanol extracts of <i>Morinda citrifolia</i> suppress advanced glycation end products (AGE)-induced inflammatory reactions in endothelial cells through its anti-oxidative properties. <i>BMC Complementary and Alternative Medicine</i> , 2017, 17, 137.	3.7	15
53	Methylglyoxal-derived hydroimidazolone-1 evokes inflammatory reactions in endothelial cells via an interaction with receptor for advanced glycation end products. <i>Diabetes and Vascular Disease Research</i> , 2017, 14, 450-453.	2.0	27
54	Glycation and cardiovascular disease in diabetes: A perspective on the concept of metabolic memory. <i>Journal of Diabetes</i> , 2017, 9, 141-148.	1.8	68

#	ARTICLE	IF	CITATIONS
55	RAGE-aptamer Attenuates the Growth and Liver Metastasis of Malignant Melanoma in Nude Mice. <i>Molecular Medicine</i> , 2017, 23, 295-306.	4.4	27
56	Phytochemicals Against Advanced Glycation End Products (AGEs) and the Receptor System. <i>Current Pharmaceutical Design</i> , 2017, 23, 1135-1141.	1.9	19
57	Circulating level of pigment epithelium-derived factor is associated with vascular function and structure: A cross-sectional study. <i>International Journal of Cardiology</i> , 2016, 225, 91-95.	1.7	10
58	Tofogliflozin, a selective inhibitor of sodium-glucose cotransporter 2, suppresses renal damage in KKAY/Ta mice, obese and type 2 diabetic animals. <i>Diabetes and Vascular Disease Research</i> , 2016, 13, 438-441.	2.0	21
59	Ranirestat has a stronger inhibitory activity on aldose reductase and suppresses inflammatory reactions in high glucose-exposed endothelial cells. <i>Diabetes and Vascular Disease Research</i> , 2016, 13, 312-315.	2.0	7
60	Iridoids are natural glycation inhibitors. <i>Glycoconjugate Journal</i> , 2016, 33, 671-681.	2.7	19
61	DNA-aptamers raised against AGEs as a blocker of various aging-related disorders. <i>Glycoconjugate Journal</i> , 2016, 33, 683-690.	2.7	19
62	Serum Levels of Growth Differentiation Factor 11 Are Independently Associated with Low Hemoglobin Values in Hemodialysis Patients. <i>BioResearch Open Access</i> , 2016, 5, 155-158.	2.6	8
63	Mechanism for the Development of Bone Disease in Diabetes: Increased Oxidative Stress and Advanced Glycation End Products. , 2016, , 63-79.		3
64	Protective role of sulphoraphane against vascular complications in diabetes. <i>Pharmaceutical Biology</i> , 2016, 54, 2329-2339.	2.9	28
65	Protective Role of Sodium-Glucose Co-Transporter 2 Inhibition Against Vascular Complications in Diabetes. <i>Rejuvenation Research</i> , 2016, 19, 107-114.	1.8	10
66	Pathologic role of dietary advanced glycation end products in cardiometabolic disorders, and therapeutic intervention. <i>Nutrition</i> , 2016, 32, 157-165.	2.4	54
67	Anagliptin, A Dipeptidyl Peptidase-4 Inhibitor Ameliorates Arterial Stiffness in Association with Reduction of Remnant-Like Particle Cholesterol and Alanine Transaminase Levels in Type 2 Diabetic Patients. <i>Current Vascular Pharmacology</i> , 2016, 14, 552-562.	1.7	21
68	Pigment epithelium-derived factor inhibits caveolin-induced interleukin-8 gene expression and proliferation of human prostate cancer cells. <i>Oncology Letters</i> , 2015, 10, 2644-2648.	1.8	6
69	DNA Aptamer Raised against Advanced Glycation End Products Prevents Abnormalities in Electroretinograms of Experimental Diabetic Retinopathy. <i>Ophthalmic Research</i> , 2015, 54, 175-180.	1.9	11
70	Insulin stimulates SGLT2-mediated tubular glucose absorption via oxidative stress generation. <i>Diabetology and Metabolic Syndrome</i> , 2015, 7, 48.	2.7	58
71	Elevation of Serum Levels of Advanced Glycation End Products in Patients With Non-alcoholic or Alcoholic Hepatocellular Carcinoma. <i>Journal of Clinical Laboratory Analysis</i> , 2015, 29, 480-484.	2.1	28
72	Assessment of the Concentrations of Various Advanced Glycation End-Products in Beverages and Foods That Are Commonly Consumed in Japan. <i>PLoS ONE</i> , 2015, 10, e0118652.	2.5	64

#	ARTICLE	IF	CITATIONS
73	Advanced Glycation End Products: A Molecular Target for Vascular Complications in Diabetes. <i>Molecular Medicine</i> , 2015, 21, S32-S40.	4.4	126
74	Receptor for advanced glycation endproducts and progressive kidney disease. <i>Current Opinion in Nephrology and Hypertension</i> , 2015, 24, 54-60.	2.0	38
75	Ratio of Serum Levels of AGEs to Soluble Form of RAGE Is a Predictor of Endothelial Function. <i>Diabetes Care</i> , 2015, 38, 119-125.	8.6	95
76	Serum levels of pigment epithelium-derived factor (PEDF) are inversely associated with circulating levels of dipeptidyl peptidase-4 (DPP-4) in humans. <i>International Journal of Cardiology</i> , 2015, 184, 14-16.	1.7	4
77	Rivaroxaban inhibits oxidative and inflammatory reactions in advanced glycation end product-exposed tubular cells by blocking thrombin/protease-activated receptor-2 system. <i>Thrombosis Research</i> , 2015, 135, 770-773.	1.7	17
78	Crosstalk between advanced glycation end products (AGEs)-receptor RAGE axis and dipeptidyl peptidase-4-incretin system in diabetic vascular complications. <i>Cardiovascular Diabetology</i> , 2015, 14, 2.	6.8	95
79	Dipeptidyl peptidase-4 deficiency protects against experimental diabetic nephropathy partly by blocking the advanced glycation end products-receptor axis. <i>Laboratory Investigation</i> , 2015, 95, 525-533.	3.7	43
80	Evaluation of tissue accumulation levels of advanced glycation end products by skin autofluorescence: A novel marker of vascular complications in high-risk patients for cardiovascular disease. <i>International Journal of Cardiology</i> , 2015, 185, 263-268.	1.7	85
81	Glyceraldehyde-derived pyridinium (GLAP) evokes oxidative stress and inflammatory and thrombogenic reactions in endothelial cells via the interaction with RAGE. <i>Cardiovascular Diabetology</i> , 2015, 14, 1.	6.8	87
82	Role of receptor for advanced glycation end products (RAGE) in liver disease. <i>European Journal of Medical Research</i> , 2015, 20, 15.	2.2	49
83	Altered serum glyceraldehyde-derived advanced glycation end product (AGE) and soluble AGE receptor levels indicate carbonyl stress in patients with schizophrenia. <i>Neuroscience Letters</i> , 2015, 593, 51-55.	2.1	19
84	Pigment epithelium-derived factor is associated with necrotic core progression during statin therapy. <i>Coronary Artery Disease</i> , 2015, 26, 107-113.	0.7	3
85	Oral L-Carnitine Supplementation Increases Trimethylamine-N-oxide but Reduces Markers of Vascular Injury in Hemodialysis Patients. <i>Journal of Cardiovascular Pharmacology</i> , 2015, 65, 289-295.	1.9	65
86	Clinical and Biochemical Factors Associated With Area and Metabolic Activity in the Visceral and Subcutaneous Adipose Tissues by FDG-PET/CT. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2015, 100, E739-E747.	3.6	40
87	Development of a monoclonal antibody-based ELISA system for glyceraldehyde-derived advanced glycation end products. <i>Immunology Letters</i> , 2015, 167, 141-146.	2.5	17
88	Role of Receptor for Advanced Glycation End Products (RAGE) and Its Ligands in Cancer Risk. <i>Rejuvenation Research</i> , 2015, 18, 48-56.	1.8	60
89	Oral consumption of sulforaphane precursor-rich broccoli supersprouts decreases serum levels of advanced glycation end products in humans. <i>Diabetes Frontier Online</i> , 2015, 2, 011-011.	0.0	2
90	Involvement of Iron-Evoked Oxidative Stress in Smoking-Related Endothelial Dysfunction in Healthy Young Men. <i>PLoS ONE</i> , 2014, 9, e89433.	2.5	7

#	ARTICLE	IF	CITATIONS
91	Switching to multiple daily injection therapy with glulisine improves glycaemic control, vascular damage and treatment satisfaction in basal insulin glargine-injected diabetic patients. <i>Diabetes/Metabolism Research and Reviews</i> , 2014, 30, 693-700.	4.0	19
92	Apixaban exerts anti-inflammatory effects in mesangial cells by blocking thrombin/protease-activated receptor-1 system. <i>Thrombosis Research</i> , 2014, 134, 1365-1367.	1.7	19
93	Possible effects of glimepiride beyond glycemic control in patients with type 2 diabetes: a preliminary report. <i>Cardiovascular Diabetology</i> , 2014, 13, 15.	6.8	28
94	Involvement of the TAGE-RAGE system in non-alcoholic steatohepatitis: Novel treatment strategies. <i>World Journal of Hepatology</i> , 2014, 6, 880.	2.0	28
95	Amelioration of experimental autoimmune uveoretinitis by inhibition of glyceraldehyde-derived advanced glycation end-product formation. <i>Journal of Leukocyte Biology</i> , 2014, 96, 1077-1085.	3.3	12
96	Advanced glycation end products potentiate citrated plasma-evoked oxidative and inflammatory reactions in endothelial cells by up-regulating protease-activated receptor-1 expression. <i>Cardiovascular Diabetology</i> , 2014, 13, 60.	6.8	50
97	Change in serum PEDF level after pioglitazone treatment is independently correlated with that in HOMA-IR. <i>International Journal of Cardiology</i> , 2014, 172, 244-246.	1.7	4
98	Serum asymmetric dimethylarginine levels are independently associated with procollagen III N-terminal peptide in nonalcoholic fatty liver disease patients. <i>Clinical and Experimental Medicine</i> , 2014, 14, 45-51.	3.6	2
99	DNA aptamer raised against advanced glycation end products inhibits neointimal hyperplasia in balloon-injured rat carotid arteries. <i>International Journal of Cardiology</i> , 2014, 171, 443-446.	1.7	17
100	Linagliptin, a xanthine-based dipeptidyl peptidase-4 inhibitor, decreases serum uric acid levels in type 2 diabetic patients partly by suppressing xanthine oxidase activity. <i>International Journal of Cardiology</i> , 2014, 176, 550-552.	1.7	33
101	DNA aptamer raised against advanced glycation end products inhibits melanoma growth in nude mice. <i>Laboratory Investigation</i> , 2014, 94, 422-429.	3.7	39
102	Dialysate Vascular Endothelial Growth Factor Is an Independent Determinant of Serum Albumin Levels and Predicts Future Withdrawal From Peritoneal Dialysis in Uremic Patients. <i>Therapeutic Apheresis and Dialysis</i> , 2014, 18, 391-397.	0.9	9
103	Elevation of soluble form of receptor for advanced glycation end products (sRAGE) in recurrent pregnancy losses (RPL): possible participation of RAGE in RPL. <i>Fertility and Sterility</i> , 2014, 102, 782-789.	1.0	20
104	Irbesartan inhibits advanced glycation end product-induced increase in asymmetric dimethylarginine level in mesangial cells through its anti-oxidative properties. <i>International Journal of Cardiology</i> , 2014, 176, 1120-1122.	1.7	16
105	Ramipril inhibits AGE-RAGE-induced matrix metalloproteinase-2 activation in experimental diabetic nephropathy. <i>Diabetology and Metabolic Syndrome</i> , 2014, 6, 86.	2.7	29
106	Sulforaphane inhibits advanced glycation end product-induced pericyte damage by reducing expression of receptor for advanced glycation end products. <i>Nutrition Research</i> , 2014, 34, 807-813.	2.9	26
107	Laminin receptor mediates anti-inflammatory and anti-thrombogenic effects of pigment epithelium-derived factor in myeloma cells. <i>Biochemical and Biophysical Research Communications</i> , 2014, 443, 847-851.	2.1	18
108	Pigment Epithelium-Derived Factor Improves Metabolic Derangements and Ameliorates Dysregulation of Adipocytokines in Obese Type 2 Diabetic Rats. <i>American Journal of Pathology</i> , 2014, 184, 1094-1103.	3.8	22

#	ARTICLE	IF	CITATIONS
109	Vascular injury is improved by pre-meal glulisine-based bolus insulin therapy in type 2 diabetic patients. <i>IJC Metabolic & Endocrine</i> , 2014, 4, 70-72.	0.5	0
110	Effect of statins on the serum soluble form of receptor for advanced glycation end-products and its association with coronary atherosclerosis in patients with angina pectoris. <i>IJC Metabolic & Endocrine</i> , 2014, 4, 47-52.	0.5	8
111	Sulforaphane inhibits formation of advanced glycation end products in vitro. <i>Diabetes Frontier Online</i> , 2014, 1, 001-001.	0.0	2
112	Pigment Epithelium-derived Factor (PEDF) and Cardiometabolic Disorders. <i>Current Pharmaceutical Design</i> , 2014, 20, 2377-2386.	1.9	29
113	Role of AGEs-RAGE System in Cardiovascular Disease. <i>Current Pharmaceutical Design</i> , 2014, 20, 2395-2402.	1.9	143
114	Nonalcoholic Fatty Liver Disease and Cardiovascular Disease. <i>Current Pharmaceutical Design</i> , 2014, 20, 2403-2411.	1.9	16
115	Molecular Imaging of Vascular Inflammation. <i>Current Pharmaceutical Design</i> , 2014, 20, 2439-2447.	1.9	13
116	Role of Asymmetric Dimethylarginine in Cardiorenal Syndrome. <i>Current Pharmaceutical Design</i> , 2014, 20, 2448-2455.	1.9	20
117	Minodronate. , 2014, , 2861-2864.		0
118	Minodronate. , 2014, , 1-4.		0
119	DNA Aptamer Raised Against AGEs Blocks the Progression of Experimental Diabetic Nephropathy. <i>Diabetes</i> , 2013, 62, 3241-3250.	0.6	72
120	Relationship between Advanced Glycation End Products and Plaque Progression in Patients with Acute Coronary Syndrome: The JAPAN-ACS Sub-study. <i>Cardiovascular Diabetology</i> , 2013, 12, 5.	6.8	55
121	Potential Inhibitory Effects of L-Carnitine Supplementation on Tissue Advanced Glycation End Products in Patients with Hemodialysis. <i>Rejuvenation Research</i> , 2013, 16, 460-466.	1.8	27
122	Efficacy of alogliptin, a dipeptidyl peptidase-4 inhibitor, on glucose parameters, the activity of the advanced glycation end product (AGE) receptor for AGE (RAGE) axis and albuminuria in Japanese type 2 diabetes. <i>Diabetes/Metabolism Research and Reviews</i> , 2013, 29, 624-630.	4.0	59
123	Asymmetrical dimethylarginine level is independently associated with circulating levels of RAGE and PEDF. <i>International Journal of Cardiology</i> , 2013, 167, 3072-3074.	1.7	2
124	Evidence for a Positive Association Between Serum Carnitine and Free Testosterone Levels in Uremic Men with Hemodialysis. <i>Rejuvenation Research</i> , 2013, 16, 200-205.	1.8	3
125	Blockade by phosphorothioate aptamers of advanced glycation end products-induced damage in cultured pericytes and endothelial cells. <i>Microvascular Research</i> , 2013, 90, 64-70.	2.5	37
126	Advanced glycation end products evoke endothelial cell damage by stimulating soluble dipeptidyl peptidase-4 production and its interaction with mannose 6-phosphate/insulin-like growth factor II receptor. <i>Cardiovascular Diabetology</i> , 2013, 12, 125.	6.8	142

#	ARTICLE	IF	CITATIONS
127	Effects of Pioglitazone on Visceral Fat Metabolic Activity in Impaired Glucose Tolerance or Type 2 Diabetes Mellitus. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2013, 98, 4438-4445.	3.6	39
128	sRAGE is associated with low waist circumference and Hb levels in NAFLD. <i>Open Medicine (Poland)</i> , 2013, 8, 830-834.	1.3	0
129	Pigment epithelium-derived factor (PEDF) binds to caveolin-1 and inhibits the pro-inflammatory effects of caveolin-1 in endothelial cells. <i>Biochemical and Biophysical Research Communications</i> , 2013, 441, 405-410.	2.1	18
130	Rosuvastatin restores advanced glycation end product-induced decrease in sirtuin1 (SIRT1) mRNA levels in THP-1 monocytic cells through its anti-oxidative properties. <i>International Journal of Cardiology</i> , 2013, 169, e102-e103.	1.7	4
131	Pigment epithelium-derived factor (PEDF) inhibits survival and proliferation of VEGF-exposed multiple myeloma cells through its anti-oxidative properties. <i>Biochemical and Biophysical Research Communications</i> , 2013, 431, 693-697.	2.1	21
132	Pigment epithelium-derived factor (PEDF) blocks high glucose-induced inflammatory reactions in endothelial cells through its anti-oxidative properties. <i>International Journal of Cardiology</i> , 2013, 168, 3004-3006.	1.7	8
133	Serum levels of advanced glycation end products (AGEs) are independently correlated with circulating levels of dipeptidyl peptidase-4 (DPP-4) in humans. <i>Clinical Biochemistry</i> , 2013, 46, 300-303.	1.9	35
134	Pioglitazone Decreases Asymmetric Dimethylarginine Levels in Patients with Impaired Glucose Tolerance or Type 2 Diabetes. <i>Rejuvenation Research</i> , 2013, 16, 344-351.	1.8	12
135	Telmisartan inhibits AGE-induced podocyte damage and detachment. <i>Microvascular Research</i> , 2013, 88, 79-83.	2.5	21
136	Author reply: Comment to "Atorvastatin improves disease activity of nonalcoholic steatohepatitis partly through its tumour necrosis factor- α -lowering property". <i>Digestive and Liver Disease</i> , 2013, 45, 82.	0.9	0
137	Author reply: Comment to "Atorvastatin improves disease activity of nonalcoholic steatohepatitis partly through its tumour necrosis factor- α -lowering property". <i>Digestive and Liver Disease</i> , 2013, 45, 83-84.	0.9	0
138	Glucagon-Like Peptide-1 Receptor Agonist Inhibits Asymmetric Dimethylarginine Generation in the Kidney of Streptozotocin-Induced Diabetic Rats by Blocking Advanced Glycation End Product-Induced Protein Arginine Methyltransferase-1 Expression. <i>American Journal of Pathology</i> , 2013, 182, 132-141.	3.8	125
139	PEDF inhibits AGE-induced podocyte apoptosis via PPAR-gamma activation. <i>Microvascular Research</i> , 2013, 85, 54-58.	2.5	48
140	Comment on: Selvin et al. sRAGE and Risk of Diabetes, Cardiovascular Disease, and Death. <i>Diabetes</i> 2013;62:2116-2121. <i>Diabetes</i> , 2013, 62, e26-e26.	0.6	6
141	Beneficial Effects of Vildagliptin on Retinal Injury in Obese Type 2 Diabetic Rats. <i>Ophthalmic Research</i> , 2013, 50, 221-226.	1.9	21
142	Involvement of advanced glycation end product-induced asymmetric dimethylarginine generation in endothelial dysfunction. <i>Diabetes and Vascular Disease Research</i> , 2013, 10, 436-441.	2.0	55
143	Sodium-glucose cotransporter 2-mediated oxidative stress augments advanced glycation end products-induced tubular cell apoptosis. <i>Diabetes/Metabolism Research and Reviews</i> , 2013, 29, 406-412.	4.0	73
144	Pigment Epithelium-Derived Factor as a New Predictor of Mortality Among Chronic Kidney Disease Patients Treated With Hemodialysis. <i>Therapeutic Apheresis and Dialysis</i> , 2013, 17, 625-630.	0.9	4

#	ARTICLE	IF	CITATIONS
145	A Diabetes Treatment Strategy to Reduce the Risk of Cardiovascular Events: Clinical Benefits and Potential of Linagliptin. <i>Immunology, Endocrine and Metabolic Agents in Medicinal Chemistry</i> , 2013, 13, 81-88.	0.5	2
146	Comment on: Monami et al. Dipeptidyl Peptidase-4 Inhibitors and Bone Fractures: A Meta-analysis of Randomized Clinical Trials. <i>Diabetes Care</i> 2011;34:2474-2476. <i>Diabetes Care</i> , 2012, 35, e33-e33.	8.6	3
147	Positive Association Between Serum Level of Glycereraldehyde-Derived Advanced Glycation End Products and Vascular Inflammation Evaluated by [18F]Fluorodeoxyglucose Positron Emission Tomography. <i>Diabetes Care</i> , 2012, 35, 2618-2625.	8.6	78
148	Addition of aliskiren to olmesartan ameliorates tubular injury in chronic kidney disease patients partly by reducing proteinuria. <i>JRAAS - Journal of the Renin-Angiotensin-Aldosterone System</i> , 2012, 13, 122-127.	1.7	16
149	An Overview of Diabetic Nephropathy. , 2012, , 145-157.		2
150	Serum Levels of Advanced Glycation End Products (AGEs) are Independent Correlates of Insulin Resistance in Nondiabetic Subjects. <i>Cardiovascular Therapeutics</i> , 2012, 30, 42-48.	2.5	96
151	Serum Levels of Advanced Glycation End Products (AGEs) are Inversely Associated with the Number and Migratory Activity of Circulating Endothelial Progenitor Cells in Apparently Healthy Subjects. <i>Cardiovascular Therapeutics</i> , 2012, 30, 249-254.	2.5	42
152	Decreased serum carnitine is independently correlated with increased tissue accumulation levels of advanced glycation end products in haemodialysis patients. <i>Nephrology</i> , 2012, 17, 689-694.	1.6	20
153	Potential Clinical Utility of Advanced Glycation End Product Cross-Link Breakers in Age- and Diabetes-Associated Disorders. <i>Rejuvenation Research</i> , 2012, 15, 564-572.	1.8	57
154	Serum levels of pigment epithelium-derived factor, a novel marker of insulin resistance, are independently associated with fasting apolipoprotein B48 levels in humans. <i>Clinical Biochemistry</i> , 2012, 45, 1404-1408.	1.9	7
155	Serum levels of pigment epithelium-derived factor (PEDF) are independently associated with procollagen III N-terminal peptide levels in patients with nonalcoholic fatty liver disease. <i>Clinical Biochemistry</i> , 2012, 45, 1554-1557.	1.9	4
156	Pravastatin inhibits advanced glycation end products (AGEs)-induced proximal tubular cell apoptosis and injury by reducing receptor for AGEs (RAGE) level. <i>Metabolism: Clinical and Experimental</i> , 2012, 61, 1067-1072.	3.4	66
157	Advanced Glycation End Products Increase Permeability of Brain Microvascular Endothelial Cells through Reactive Oxygen Species-Induced Vascular Endothelial Growth Factor Expression. <i>Journal of Stroke and Cerebrovascular Diseases</i> , 2012, 21, 293-298.	1.6	31
158	PEDF-derived peptide inhibits corneal angiogenesis by suppressing VEGF expression. <i>Microvascular Research</i> , 2012, 84, 105-108.	2.5	39
159	Addition of aliskiren to angiotensin II receptor blockers ameliorates renal tubular injury and reduces intima media thickness of carotid artery in patients with diabetic nephropathy. <i>International Journal of Cardiology</i> , 2012, 155, 294-296.	1.7	6
160	Vildagliptin inhibits oxidative stress and vascular damage in streptozotocin-induced diabetic rats. <i>International Journal of Cardiology</i> , 2012, 158, 171-173.	1.7	29
161	Serum pigment epithelium-derived factor levels are independently associated with decreased number of circulating endothelial progenitor cells in healthy non-smokers. <i>International Journal of Cardiology</i> , 2012, 158, 310-312.	1.7	2
162	Adiponectin is inversely associated with ratio of serum levels of AGEs to sRAGE and vascular inflammation. <i>International Journal of Cardiology</i> , 2012, 158, 461-462.	1.7	32

#	ARTICLE	IF	CITATIONS
163	Role of advanced glycation end products (AGEs) and oxidative stress in vascular complications in diabetes. <i>Biochimica Et Biophysica Acta - General Subjects</i> , 2012, 1820, 663-671.	2.4	226
164	Atorvastatin improves disease activity of nonalcoholic steatohepatitis partly through its tumour necrosis factor- α -lowering property. <i>Digestive and Liver Disease</i> , 2012, 44, 492-496.	0.9	50
165	Increased insulinogenic index is an independent determinant of nonalcoholic fatty liver disease activity score in patients with normal glucose tolerance. <i>Digestive and Liver Disease</i> , 2012, 44, 935-939.	0.9	4
166	Beneficial effects of metformin and irbesartan on advanced glycation end products (AGEs)-RAGE-induced proximal tubular cell injury. <i>Pharmacological Research</i> , 2012, 65, 297-302.	7.1	39
167	Proteinuria elevates asymmetric dimethylarginine levels via protein arginine methyltransferase-1 overexpression in a rat model of nephrotic syndrome. <i>Life Sciences</i> , 2012, 91, 301-305.	4.3	13
168	Glucagon-like peptide-1 inhibits angiotensin II-induced mesangial cell damage via protein kinase A. <i>Microvascular Research</i> , 2012, 84, 395-398.	2.5	38
169	Statins and hemoperfusion improve 28-day survival in septic shock patients. <i>Open Medicine (Poland)</i> , 2012, 7, 475-480.	1.3	0
170	Pioglitazone Attenuates Atherosclerotic Plaque Inflammation in Patients With Impaired Glucose Tolerance or Diabetes. <i>JACC: Cardiovascular Imaging</i> , 2011, 4, 1110-1118.	5.3	126
171	Pigment epithelium-derived factor (PEDF) inhibits proximal tubular cell injury in early diabetic nephropathy by suppressing advanced glycation end products (AGEs)-receptor (RAGE) axis. <i>Pharmacological Research</i> , 2011, 63, 241-248.	7.1	50
172	Vildagliptin blocks vascular injury in thoracic aorta of diabetic rats by suppressing advanced glycation end product-RAGE receptor axis. <i>Pharmacological Research</i> , 2011, 63, 383-388.	7.1	88
173	Administration of Pigment Epithelium-Derived Factor Inhibits Left Ventricular Remodeling and Improves Cardiac Function in Rats with Acute Myocardial Infarction. <i>American Journal of Pathology</i> , 2011, 178, 591-598.	3.8	41
174	Irbesartan inhibits advanced glycation end product (AGE)-induced up-regulation of vascular cell adhesion molecule-1 (VCAM-1) mRNA levels in glomerular endothelial cells. <i>Microvascular Research</i> , 2011, 81, 269-273.	2.5	20
175	Osteogenic differentiation of mesenchymal stem cells is preserved by partial, but not full peroxisome proliferator-activated receptor-gamma agonist. <i>International Journal of Cardiology</i> , 2011, 146, 109-110.	1.7	1
176	Co-treatment with azelidipine and olmesartan inhibits advanced glycation end products (AGEs) elicited down-regulation of adiponectin mRNA levels in cultured adipocytes partly via its anti-oxidative property. <i>International Journal of Cardiology</i> , 2011, 146, 264-266.	1.7	7
177	Ezetimibe, an inhibitor of intestinal cholesterol absorption, decreases serum level of malondialdehyde-modified low-density lipoprotein in patients with hypercholesterolemia. <i>International Journal of Cardiology</i> , 2011, 146, 420-421.	1.7	9
178	Pigment epithelium-derived factor (PEDF) blocks advanced glycation end products (AGEs)-RAGE-induced suppression of adiponectin mRNA level in adipocytes by inhibiting NADPH oxidase-mediated oxidative stress generation. <i>International Journal of Cardiology</i> , 2011, 152, 408-410.	1.7	29
179	Nitric oxide, a janus-faced therapeutic target for diabetic microangiopathy-Friend or foe?. <i>Pharmacological Research</i> , 2011, 64, 187-194.	7.1	100
180	Serum level of pigment epithelium-derived factor is a marker of atherosclerosis in humans. <i>Atherosclerosis</i> , 2011, 219, 311-315.	0.8	48

#	ARTICLE	IF	CITATIONS
181	Activation of Receptor for Advanced Glycation End Products Induces Osteogenic Differentiation of Vascular Smooth Muscle Cells. <i>Journal of Atherosclerosis and Thrombosis</i> , 2011, 18, 670-683.	2.0	72
182	Pleiotropic Effects of Glucagon-like Peptide-1 (GLP-1)-Based Therapies on Vascular Complications in Diabetes. <i>Current Pharmaceutical Design</i> , 2011, 17, 4379-4385.	1.9	35
183	Association between Plasma Pigment Epithelium-derived Factor Levels and Tissue Characteristics of Coronary Plaque using Integrated Backscatter Intravascular Ultrasound. <i>Internal Medicine</i> , 2011, 50, 1889-1894.	0.7	5
184	Role of Advanced Glycation End Products (AGEs) in Osteoporosis in Diabetes. <i>Current Drug Targets</i> , 2011, 12, 2096-2102.	2.1	127
185	Tissue level of advanced glycation end products is an independent determinant of high-sensitivity C-reactive protein levels in haemodialysis patients. <i>Nephrology</i> , 2011, 16, 299-303.	1.6	21
186	Oral adsorbent AST-120 ameliorates tubular injury in chronic renal failure patients by reducing proteinuria and oxidative stress generation. <i>Metabolism: Clinical and Experimental</i> , 2011, 60, 260-264.	3.4	44
187	Glucagon-like peptide-1 suppresses advanced glycation end product-induced monocyte chemoattractant protein-1 expression in mesangial cells by reducing advanced glycation end product receptor level. <i>Metabolism: Clinical and Experimental</i> , 2011, 60, 1271-1277.	3.4	86
188	Suppression of high-mobility group box-1 and receptor for advanced glycation end-product axis by polymyxin B-immobilized fiber hemoperfusion in septic shock patients. <i>Journal of Critical Care</i> , 2011, 26, 546-549.	2.2	13
189	Role of advanced glycation end products (AGEs) and receptor for AGEs (RAGE) in vascular damage in diabetes. <i>Experimental Gerontology</i> , 2011, 46, 217-224.	2.8	231
190	High White Blood Cell Count and Low Estimated Glomerular Filtration Rate Are Independently Associated With Serum Level of Monocyte Chemoattractant Protein-1 in a General Population. <i>Clinical Cardiology</i> , 2011, 34, 189-194.	1.8	15
191	Beneficial Cardiometabolic Actions of Telmisartan Plus Amlodipine Therapy in Elderly Patients With Poorly Controlled Hypertension. <i>Clinical Cardiology</i> , 2011, 34, 261-265.	1.8	7
192	Calcium Channel Blocker Inhibition of AGE and RAGE Axis Limits Renal Injury in Nondiabetic Patients With Stage I or II Chronic Kidney Disease. <i>Clinical Cardiology</i> , 2011, 34, 372-377.	1.8	23
193	Vardenafil, an inhibitor of phosphodiesterase-5, blocks advanced glycation end product (AGE)-induced up-regulation of monocyte chemoattractant protein-1 mRNA levels in endothelial cells by suppressing AGE receptor (RAGE) expression via elevation of cGMP. <i>Clinical and Experimental Medicine</i> , 2011, 11, 131-135.	3.6	19
194	Involvement of membrane type 1 matrix metalloproteinase (MT1-MMP) in RAGE activation signaling pathways. <i>Journal of Cellular Physiology</i> , 2011, 226, 1554-1563.	4.1	22
195	Increased levels of soluble receptor for advanced glycation end products (sRAGE) and high mobility group box-1 (HMGB1) are associated with death in patients with acute respiratory distress syndrome. <i>Clinical Biochemistry</i> , 2011, 44, 601-604.	1.9	70
196	Significance of pigment epithelium-derived factor levels with angiotensin II type 1 receptor blockers in patients with successful coronary stent implantation. <i>JRAAS - Journal of the Renin-Angiotensin-Aldosterone System</i> , 2011, 12, 320-325.	1.7	1
197	Advanced Glycation End Products (AGEs), Oxidative Stress and Diabetic Retinopathy. <i>Current Pharmaceutical Biotechnology</i> , 2011, 12, 362-368.	1.6	66
198	Minodronate. , 2011, , 2325-2327.		0

#	ARTICLE	IF	CITATIONS
199	Low Serum Level of the Endogenous Secretory Receptor for Advanced Glycation End Products (esRAGE) Is a Risk Factor for Prevalent Vertebral Fractures Independent of Bone Mineral Density in Patients With Type 2 Diabetes. <i>Diabetes Care</i> , 2010, 33, e58-e58.	8.6	2
200	The p66 ^{shc} Gene Expression in Peripheral Blood Monocytes Is Increased in Patients With Coronary Artery Disease. <i>Clinical Cardiology</i> , 2010, 33, 548-552.	1.8	16
201	An α -glucosidase inhibitor, acarbose treatment decreases serum levels of glyceraldehyde-derived advanced glycation end products (AGEs) in patients with type 2 diabetes. <i>Clinical and Experimental Medicine</i> , 2010, 10, 139-141.	3.6	29
202	Atorvastatin decreases serum levels of advanced glycation endproducts (AGEs) in nonalcoholic steatohepatitis (NASH) patients with dyslipidemia: clinical usefulness of AGEs as a biomarker for the attenuation of NASH. <i>Journal of Gastroenterology</i> , 2010, 45, 750-757.	5.1	141
203	Immunological detection of fructose-derived advanced glycation end-products. <i>Laboratory Investigation</i> , 2010, 90, 1117-1127.	3.7	49
204	Atorvastatin Reduces Proteinuria in Non-Diabetic Chronic Kidney Disease Patients Partly via Lowering Serum Levels of Advanced Glycation End Products (AGEs). <i>Oxidative Medicine and Cellular Longevity</i> , 2010, 3, 304-307.	4.0	26
205	Advanced Glycation end Products, Oxidative Stress and Diabetic Nephropathy. <i>Oxidative Medicine and Cellular Longevity</i> , 2010, 3, 101-108.	4.0	298
206	Efficacy of Combination Therapy with Telmisartan Plus Amlodipine in Patients with Poorly Controlled Hypertension. <i>Oxidative Medicine and Cellular Longevity</i> , 2010, 3, 342-346.	4.0	6
207	Soluble form of a receptor for advanced glycation end products sRAGE as a biomarker. <i>Frontiers in Bioscience - Elite</i> , 2010, E2, 1184-1195.	1.8	114
208	Involvement of TAGE-RAGE System in the Pathogenesis of Diabetic Retinopathy. <i>Journal of Ophthalmology</i> , 2010, 2010, 1-12.	1.3	18
209	Cancer Malignancy Is Enhanced by Glyceraldehyde-Derived Advanced Glycation End-Products. <i>Journal of Oncology</i> , 2010, 2010, 1-8.	1.3	51
210	Involvement of the Toxic AGEs (TAGE)-RAGE System in the Pathogenesis of Diabetic Vascular Complications: A Novel Therapeutic Strategy. <i>Current Drug Targets</i> , 2010, 11, 1468-1482.	2.1	81
211	Insulin Resistance is an Independent Correlate of High Serum Levels of Advanced Glycation End Products (AGEs) and Low Testosterone in Non-Diabetic Men. <i>Oxidative Medicine and Cellular Longevity</i> , 2010, 3, 262-265.	4.0	18
212	An improved anion-exchange high-performance liquid chromatography method for measuring oxidized form of LDLs in human plasma. <i>Annals of Clinical Biochemistry</i> , 2010, 47, 460-466.	1.6	2
213	Irbesartan Inhibits Albumin-Elicited Proximal Tubular Cell Apoptosis and Injury In Vitro. <i>Protein and Peptide Letters</i> , 2010, 17, 74-77.	0.9	12
214	Smooth Muscle Cell Pathophysiology and Advanced Glycation End Products (AGEs). <i>Current Drug Targets</i> , 2010, 11, 875-881.	2.1	24
215	Pyridoxamine, an Inhibitor of Advanced Glycation End Product (AGE) Formation Ameliorates Insulin Resistance in Obese, Type 2 Diabetic Mice. <i>Protein and Peptide Letters</i> , 2010, 17, 1177-1181.	0.9	45
216	Topical application of anti-angiogenic peptides based on pigment epithelium-derived factor can improve psoriasis. <i>Journal of Dermatological Science</i> , 2010, 57, 183-191.	1.9	33

#	ARTICLE	IF	CITATIONS
217	Positive association of circulating levels of advanced glycation end products (AGEs) with pigment epithelium-derived factor (PEDF) in a general population. <i>Pharmacological Research</i> , 2010, 61, 103-107.	7.1	31
218	Irbesartan inhibits advanced glycation end product (AGE)-induced proximal tubular cell injury in vitro by suppressing receptor for AGEs (RAGE) expression. <i>Pharmacological Research</i> , 2010, 61, 34-39.	7.1	62
219	Co-administration of ezetimibe enhances proteinuria-lowering effects of pitavastatin in chronic kidney disease patients partly via a cholesterol-independent manner. <i>Pharmacological Research</i> , 2010, 61, 58-61.	7.1	42
220	Involvement of angiotensin II in intestinal cholesterol absorption. <i>Pharmacological Research</i> , 2010, 61, 460-465.	7.1	8
221	Pigment epithelium-derived factor inhibits advanced glycation end product-elicited mesangial cell damage by blocking NF- κ B activation. <i>Microvascular Research</i> , 2010, 80, 227-232.	2.5	51
222	Pigment epithelium-derived factor (PEDF) inhibits collagen-induced platelet activation by reducing intraplatelet nitrotyrosine levels. <i>International Journal of Cardiology</i> , 2010, 140, 121-122.	1.7	9
223	Serum levels of pigment epithelium-derived factor (PEDF) are an independent determinant of insulin resistance in patients with essential hypertension. <i>International Journal of Cardiology</i> , 2010, 143, 96-98.	1.7	30
224	Pigment epithelium-derived factor (PEDF) inhibits diabetes- or advanced glycation end product (AGE)-induced platelet CD40 ligand overexpression in rats. <i>International Journal of Cardiology</i> , 2010, 144, 283-285.	1.7	4
225	Involvement of aldosterone-mineralocorticoid receptor system in advanced glycation end product (AGE)-elicited plasminogen activator inhibitor-1 (PAI-1) expression in diabetes. <i>International Journal of Cardiology</i> , 2010, 145, 566-567.	1.7	3
226	Glucagon-like peptide-1 (GLP-1) inhibits advanced glycation end product (AGE)-induced up-regulation of VCAM-1 mRNA levels in endothelial cells by suppressing AGE receptor (RAGE) expression. <i>Biochemical and Biophysical Research Communications</i> , 2010, 391, 1405-1408.	2.1	136
227	Nifedipine, a calcium channel blocker, inhibits inflammatory and fibrogenic gene expressions in advanced glycation end product (AGE)-exposed fibroblasts via mineralocorticoid receptor antagonistic activity. <i>Biochemical and Biophysical Research Communications</i> , 2010, 396, 566-570.	2.1	27
228	Nifedipine inhibits advanced glycation end products (AGEs) and their receptor (RAGE) interaction-mediated proximal tubular cell injury via peroxisome proliferator-activated receptor-gamma activation. <i>Biochemical and Biophysical Research Communications</i> , 2010, 398, 326-330.	2.1	33
229	Pigment Epithelium-Derived Factor Inhibits Lysosomal Degradation of Bcl-xL and Apoptosis in HepG2 cells. <i>American Journal of Pathology</i> , 2010, 176, 168-176.	3.8	21
230	Administration of pigment epithelium-derived factor (PEDF) reduces proteinuria by suppressing decreased nephrin and increased VEGF expression in the glomeruli of adriamycin-injected rats. <i>Nephrology Dialysis Transplantation</i> , 2009, 24, 1397-1406.	0.7	33
231	Independent determinants of soluble form of receptor for advanced glycation end products in elderly hypertensive patients. <i>Metabolism: Clinical and Experimental</i> , 2009, 58, 421-425.	3.4	19
232	Factors associated with serum high mobility group box 1 (HMGB1) levels in a general population. <i>Metabolism: Clinical and Experimental</i> , 2009, 58, 1688-1693.	3.4	43
233	Protective role of pigment epithelium-derived factor (PEDF) in early phase of experimental diabetic retinopathy. <i>Diabetes/Metabolism Research and Reviews</i> , 2009, 25, 678-686.	4.0	103
234	Serum levels of pigment epithelium-derived factor (PEDF) are positively associated with visceral adiposity in Japanese patients with type 2 diabetes. <i>Diabetes/Metabolism Research and Reviews</i> , 2009, 25, 52-56.	4.0	61

#	ARTICLE	IF	CITATIONS
235	Pigment epithelium-derived factor (PEDF) prevents platelet activation and aggregation in diabetic rats by blocking deleterious effects of advanced glycation end products (AGEs). <i>Diabetes/Metabolism Research and Reviews</i> , 2009, 25, 266-271.	4.0	41
236	Effects of high-AGE beverage on RAGE and VEGF expressions in the liver and kidneys. <i>European Journal of Nutrition</i> , 2009, 48, 6-11.	3.9	33
237	Advanced Glycation End Products and Receptor-Induced Oxidative Stress System in Diabetic Vascular Complications. <i>Therapeutic Apheresis and Dialysis</i> , 2009, 13, 534-539.	0.9	53
238	Decreased High-Density Lipoprotein Cholesterol Level is an Independent Correlate of Circulating Tumor Necrosis Factor- α in a General Population. <i>Clinical Cardiology</i> , 2009, 32, E29-32.	1.8	15
239	Low-Density Lipoprotein Levels are One of the Independent Determinants of Circulating Levels of Advanced Glycation End Products in Nondiabetic Subjects. <i>Clinical Cardiology</i> , 2009, 32, E12-5.	1.8	8
240	Nifedipine, a calcium channel blocker, inhibits advanced glycation end product (AGE)-elicited mesangial cell damage by suppressing AGE receptor (RAGE) expression via peroxisome proliferator-activated receptor-gamma activation. <i>Biochemical and Biophysical Research Communications</i> , 2009, 385, 269-272.	2.1	48
241	Serum level of pigment epithelium derived factor (PEDF) is an independent determinant of resting heart rate in Japanese subjects. <i>International Journal of Cardiology</i> , 2009, 136, 245-247.	1.7	13
242	Regulation of advanced glycation end product (AGE)-receptor (RAGE) system by PPAR-gamma agonists and its implication in cardiovascular disease. <i>Pharmacological Research</i> , 2009, 60, 174-178.	7.1	67
243	Ezetimibe decreases serum levels of asymmetric dimethylarginine (ADMA) and ameliorates renal injury in non-diabetic chronic kidney disease patients in a cholesterol-independent manner. <i>Pharmacological Research</i> , 2009, 60, 525-528.	7.1	44
244	Circulating levels of advanced glycation end products (AGE) and interleukin-6 (IL-6) are independent determinants of serum asymmetric dimethylarginine (ADMA) levels in patients with septic shock. <i>Pharmacological Research</i> , 2009, 60, 515-518.	7.1	41
245	Positive association of serum levels of advanced glycation end products and high mobility group box-1 with asymmetric dimethylarginine in nondiabetic chronic kidney disease patients. <i>Metabolism: Clinical and Experimental</i> , 2009, 58, 1624-1628.	3.4	48
246	Involvement of Toxic AGEs (TAGE) in the Pathogenesis of Diabetic Vascular Complications and Alzheimer's Disease. <i>Journal of Alzheimer's Disease</i> , 2009, 16, 845-858.	2.6	104
247	Clinical Utility of Acarbose, an α -Glucosidase Inhibitor in Cardiometabolic Disorders. <i>Current Drug Metabolism</i> , 2009, 10, 159-163.	1.2	41
248	Atheroprotective Properties of Pigment Epithelium-Derived Factor (PEDF) in Cardiometabolic Disorders. <i>Current Pharmaceutical Design</i> , 2009, 15, 1027-1033.	1.9	47
249	Advanced glycation end products enhance the proliferation and activation of hepatic stellate cells. <i>Journal of Gastroenterology</i> , 2008, 43, 298-304.	5.1	93
250	Circulating advanced glycation end products (AGEs) and soluble form of receptor for AGEs (sRAGE) are independent determinants of serum monocyte chemoattractant protein-1 (MCP-1) levels in patients with type 2 diabetes. <i>Diabetes/Metabolism Research and Reviews</i> , 2008, 24, 109-114.	4.0	80
251	Hyperglycemia induces oxidative and nitrosative stress and increases renal functional impairment in Nrf2-deficient mice. <i>Genes To Cells</i> , 2008, 13, 1159-1170.	1.2	175
252	Inhibitors of Advanced Glycation End Products (AGEs): Potential Utility for the Treatment of Cardiovascular Disease. <i>Cardiovascular Drug Reviews</i> , 2008, 26, 50-58.	4.1	23

#	ARTICLE	IF	CITATIONS
253	Olmesartan blocks advanced glycation end products (AGEs)-induced angiogenesis in vitro by suppressing receptor for AGEs (RAGE) expression. <i>Microvascular Research</i> , 2008, 75, 130-134.	2.5	56
254	Serum levels of soluble form of receptor for advanced glycation end products (sRAGE) are positively associated with circulating AGEs and soluble form of VCAM-1 in patients with type 2 diabetes. <i>Microvascular Research</i> , 2008, 76, 52-56.	2.5	87
255	Pigment epithelium-derived factor (PEDF) administration inhibits occlusive thrombus formation in rats: A possible participation of reduced intraplatelet PEDF in thrombosis of acute coronary syndromes. <i>Atherosclerosis</i> , 2008, 197, 25-33.	0.8	49
256	Role of AGEs in Diabetic Nephropathy. <i>Current Pharmaceutical Design</i> , 2008, 14, 946-952.	1.9	127
257	Role of Asymmetric Dimethylarginine (ADMA) in Diabetic Vascular Complications. <i>Current Pharmaceutical Design</i> , 2008, 14, 2613-2618.	1.9	38
258	Agents that block advanced glycation end product (AGE)-RAGE (receptor for AGEs)-oxidative stress system: a novel therapeutic strategy for diabetic vascular complications. <i>Expert Opinion on Investigational Drugs</i> , 2008, 17, 983-996.	4.1	121
259	Olmesartan Blocks Inflammatory Reactions in Endothelial Cells Evoked by Advanced Glycation End Products by Suppressing Generation of Reactive Oxygen Species. <i>Ophthalmic Research</i> , 2008, 40, 10-15.	1.9	59
260	Role of Oxidative Stress in the Development of Vascular Injury and its Therapeutic Intervention by Nifedipine. <i>Current Medicinal Chemistry</i> , 2008, 15, 172-177.	2.4	50
261	Receptor for Advanced Glycation End Products (RAGE): A Novel Therapeutic Target for Diabetic Vascular Complication. <i>Current Pharmaceutical Design</i> , 2008, 14, 487-495.	1.9	95
262	AGE-RAGE System and Carcinogenesis. <i>Current Pharmaceutical Design</i> , 2008, 14, 940-945.	1.9	149
263	Advanced Glycation End Products and Insulin Resistance. <i>Current Pharmaceutical Design</i> , 2008, 14, 987-989.	1.9	142
264	Role of Advanced Glycation End Products (AGEs) and Oxidative Stress in Diabetic Retinopathy. <i>Current Pharmaceutical Design</i> , 2008, 14, 962-968.	1.9	89
265	Advanced Glycation End Products (AGEs) and their Involvement in Liver Disease. <i>Current Pharmaceutical Design</i> , 2008, 14, 969-972.	1.9	123
266	Possible Involvement of Advanced Glycation End-Products (AGEs) in the Pathogenesis of Alzheimers Disease. <i>Current Pharmaceutical Design</i> , 2008, 14, 973-978.	1.9	200
267	Possible involvement of advanced glycation end products in carry-over benefits of atorvastatin in ASCOT-BPLA. <i>European Heart Journal</i> , 2008, 29, 1922-1922.	2.2	3
268	Involvement of asymmetric dimethylarginine (ADMA) in tubulointerstitial ischaemia in the early phase of diabetic nephropathy. <i>Nephrology Dialysis Transplantation</i> , 2008, 24, 1162-1169.	0.7	48
269	Comment on: Thallas-Bonke et al. (2008) Inhibition of NADPH Oxidase Prevents Advanced Glycation End Product-Mediated Damage in Diabetic Nephropathy Through a Protein Kinase C- α -Dependent Pathway: <i>Diabetes</i> 57:460-469, 2008. <i>Diabetes</i> , 2008, 57, e13-e13.	0.6	5
270	Telmisartan Inhibits Advanced Glycation End Products (AGEs)-Elicited Endothelial Cell Injury by Suppressing AGE Receptor (RAGE) Expression Via Peroxisome Proliferator-Activated Receptor- γ ; Activation. <i>Protein and Peptide Letters</i> , 2008, 15, 850-853.	0.9	43

#	ARTICLE	IF	CITATIONS
271	Pigment Epithelium-Derived Factor Prevents Melanoma Growth via Angiogenesis Inhibition. <i>Current Pharmaceutical Design</i> , 2008, 14, 3802-3809.	1.9	17
272	Pigment Epithelium-Derived Factor (PEDF): Its Potential Therapeutic Implication in Diabetic Vascular Complications. <i>Current Drug Targets</i> , 2008, 9, 1025-1029.	2.1	40
273	Telmisartan Blocks Advanced Glycation End Product (AGE)-Induced Plasminogen Activator Inhibitor-1 (PAI-1) Gene Expression in Endothelial Cells via Activation of Peroxisome Proliferator-Activated Receptor- α ; (PPAR- α);). <i>Letters in Drug Design and Discovery</i> , 2008, 5, 477-480.	0.7	6
274	Positive association of pigment epithelium-derived factor with total antioxidant capacity in the vitreous fluid of patients with proliferative diabetic retinopathy. <i>British Journal of Ophthalmology</i> , 2007, 91, 885-887.	3.9	38
275	Angiogenesis and Metastasis Inhibitors for the Treatment of Malignant Melanoma. <i>Mini-Reviews in Medicinal Chemistry</i> , 2007, 7, 649-661.	2.4	25
276	Pigment Epithelium-Derived Factor (PEDF) Prevents Advanced Glycation End Products (AGEs)-Elicited Endothelial Nitric Oxide Synthase (eNOS) Reduction Through Its Anti-Oxidative Properties. <i>Protein and Peptide Letters</i> , 2007, 14, 832-835.	0.9	35
277	Potential Utility of Telmisartan, an Angiotensin II Type 1 Receptor Blocker with Peroxisome Proliferator-Activated Receptor- β (PPAR- β)-Modulating Activity for the Treatment of Cardiometabolic Disorders. <i>Current Molecular Medicine</i> , 2007, 7, 463-469.	1.3	72
278	Food-Derived Advanced Glycation end Products (AGEs): A Novel Therapeutic Target for Various Disorders. <i>Current Pharmaceutical Design</i> , 2007, 13, 2832-2836.	1.9	114
279	Positive correlation of pigment epithelium-derived factor and total antioxidant capacity in aqueous humour of patients with uveitis and proliferative diabetic retinopathy. <i>British Journal of Ophthalmology</i> , 2007, 91, 1133-1134.	3.9	22
280	Positive correlation between pigment epithelium-derived factor and monocyte chemoattractant protein-1 levels in the aqueous humour of patients with uveitis. <i>British Journal of Ophthalmology</i> , 2007, 91, 737-738.	3.9	11
281	Pigment-Epithelium-Derived Factor Suppresses Expression of Receptor for Advanced Glycation End Products in the Eye of Diabetic Rats. <i>Ophthalmic Research</i> , 2007, 39, 92-97.	1.9	53
282	Molecular Mechanisms of Diabetic Nephropathy and Its Therapeutic Intervention. <i>Current Drug Targets</i> , 2007, 8, 952-959.	2.1	137
283	Serum Levels of Soluble Form of Receptor for Advanced Glycation End Products (sRAGE) May Reflect Tissue RAGE Expression In Diabetes. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2007, 27, e32; author reply e33-4.	2.4	30
284	Dimethylarginine Dimethylaminohydrolase Prevents Progression of Renal Dysfunction by Inhibiting Loss of Peritubular Capillaries and Tubulointerstitial Fibrosis in a Rat Model of Chronic Kidney Disease. <i>Journal of the American Society of Nephrology: JASN</i> , 2007, 18, 1525-1533.	6.1	106
285	Positive correlation between vitreous levels of advanced glycation end products and vascular endothelial growth factor in patients with diabetic retinopathy sufficiently treated with photocoagulation. <i>British Journal of Ophthalmology</i> , 2007, 91, 397-398.	3.9	14
286	Fluvastatin Alters Platelet Aggregability in Patients With Hypercholesterolemia. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2007, 27, 1471-1477.	2.4	65
287	Kinetics, Role and Therapeutic Implications of Endogenous Soluble form of Receptor for Advanced Glycation end Products (sRAGE) in Diabetes. <i>Current Drug Targets</i> , 2007, 8, 1138-1143.	2.1	71
288	Advanced Glycation End Products (AGEs) and Cardiovascular Disease (CVD) in Diabetes. <i>Cardiovascular and Hematological Agents in Medicinal Chemistry</i> , 2007, 5, 236-240.	1.0	28

#	ARTICLE	IF	CITATIONS
289	Pigment Epithelium-Derived Factor (PEDF) Inhibits Angiotensin II-Induced Smooth Muscle Cell Proliferation Through Its Anti-Oxidative Properties. <i>Protein and Peptide Letters</i> , 2007, 14, 615-617.	0.9	10
290	Signal Transduction Therapy of Diabetic Vascular Complication. <i>Current Signal Transduction Therapy</i> , 2007, 2, 91-100.	0.5	5
291	Upregulation of the ligand-RAGE pathway via the angiotensin II type I receptor is essential in the pathogenesis of diabetic atherosclerosis. <i>Journal of Molecular and Cellular Cardiology</i> , 2007, 43, 455-464.	1.9	37
292	Advanced glycation end products attenuate cellular insulin sensitivity by increasing the generation of intracellular reactive oxygen species in adipocytes. <i>Diabetes Research and Clinical Practice</i> , 2007, 76, 236-244.	2.8	135
293	Diagnostic utility of serum or cerebrospinal fluid levels of toxic advanced glycation end-products (TAGE) in early detection of Alzheimer's disease. <i>Medical Hypotheses</i> , 2007, 69, 1358-1366.	1.5	39
294	In vitro selection of DNA aptamers that block toxic effects of AGE on cultured retinal pericytes. <i>Microvascular Research</i> , 2007, 74, 65-69.	2.5	25
295	Elevated circulating oxidized LDL levels in Japanese subjects with the metabolic syndrome. <i>International Journal of Cardiology</i> , 2007, 118, 270-272.	1.7	21
296	Pigment Epithelium-Derived Factor Inhibits Neointimal Hyperplasia after Vascular Injury by Blocking NADPH Oxidase-Mediated Reactive Oxygen Species Generation. <i>American Journal of Pathology</i> , 2007, 170, 2159-2170.	3.8	66
297	Vascular Inflammation Evaluated by [18F]-Fluorodeoxyglucose Positron Emission Tomography Is Associated With the Metabolic Syndrome. <i>Journal of the American College of Cardiology</i> , 2007, 49, 1533-1539.	2.8	226
298	Serum Levels of sRAGE, the Soluble Form of Receptor for Advanced Glycation End Products, Are Associated with Inflammatory Markers in Patients with Type 2 Diabetes. <i>Molecular Medicine</i> , 2007, 13, 185-189.	4.4	125
299	Elevation of soluble form of receptor for advanced glycation end products (sRAGE) in diabetic subjects with coronary artery disease. <i>Diabetes/Metabolism Research and Reviews</i> , 2007, 23, 368-371.	4.0	124
300	Elevated levels of serum advanced glycation end products in patients with non-alcoholic steatohepatitis. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2007, 22, 1112-1119.	2.8	164
301	Asymmetric dimethylarginine may be a missing link between cardiovascular disease and chronic kidney disease (Review Article). <i>Nephrology</i> , 2007, 12, 582-590.	1.6	61
302	Administration of pigment epithelium-derived factor (PEDF) inhibits cold injury-induced brain edema in mice. <i>Brain Research</i> , 2007, 1167, 92-100.	2.2	25
303	Pitavastatin inhibits lysophosphatidic acid-induced proliferation and monocyte chemoattractant protein-1 expression in aortic smooth muscle cells by suppressing Rac-1-mediated reactive oxygen species generation. <i>Vascular Pharmacology</i> , 2007, 46, 286-292.	2.1	53
304	Asymmetric dimethylarginine (ADMA) is a novel emerging risk factor for cardiovascular disease and the development of renal injury in chronic kidney disease. <i>Clinical and Experimental Nephrology</i> , 2007, 11, 115-121.	1.6	51
305	Increased levels of pigment epithelium-derived factor in aqueous humor of patients with uveitis. <i>British Journal of Ophthalmology</i> , 2007, 91, 149-150.	3.9	13
306	Atorvastatin Inhibits Advanced Glycation End Products (AGE)-Induced C-Reactive Expression in Hepatoma Cells by Suppressing Reactive Oxygen Species Generation. <i>Vascular Disease Prevention</i> , 2007, 4, 213-216.	0.2	2

#	ARTICLE	IF	CITATIONS
307	Elevated Serum Levels of Pigment Epithelium-Derived Factor in the Metabolic Syndrome. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2006, 91, 2447-2450.	3.6	182
308	Positive association of serum levels of advanced glycation end products with thrombogenic markers in humans. <i>Metabolism: Clinical and Experimental</i> , 2006, 55, 912-917.	3.4	40
309	Positive association between serum levels of advanced glycation end products and the soluble form of receptor for advanced glycation end products in nondiabetic subjects. <i>Metabolism: Clinical and Experimental</i> , 2006, 55, 1227-1231.	3.4	137
310	Pigment epithelium-derived factor (PEDF) inhibits advanced glycation end product (AGE)-induced C-reactive protein expression in hepatoma cells by suppressing Rac-1 activation. <i>FEBS Letters</i> , 2006, 580, 2788-2796.	2.8	33
311	Pigment epithelium-derived factor (PEDF) blocks the interleukin-6 signaling to C-reactive protein expression in Hep3B cells by suppressing Rac-1 activation. <i>Life Sciences</i> , 2006, 79, 1981-1987.	4.3	12
312	Pigment-epithelium-derived factor (PEDF) inhibits angiotensin-II-induced vascular endothelial growth factor (VEGF) expression in MOLT-3 T cells through anti-oxidative properties. <i>Microvascular Research</i> , 2006, 71, 222-226.	2.5	23
313	Pigment epithelium-derived factor (PEDF) prevents diabetes- or advanced glycation end products (AGE)-elicited retinal leukostasis. <i>Microvascular Research</i> , 2006, 72, 86-90.	2.5	77
314	Oral Adsorbent AST-120 Decreases Serum Levels of AGEs in Patients with Chronic Renal Failure. <i>Molecular Medicine</i> , 2006, 12, 180-184.	4.4	54
315	Pigment Epithelium-Derived Factor (PEDF) Blocks Angiotensin III-induced T Cell Proliferation by Suppressing Autocrine Production of Interleukin-2. <i>Medicinal Chemistry</i> , 2006, 2, 265-269.	1.5	13
316	Low-Density Lipoprotein Subfractions and the Prevalence of Silent Lacunar Infarction in Subjects with Essential Hypertension. <i>Hypertension Research</i> , 2006, 29, 303-307.	2.7	15
317	Susceptibility of brain microvascular endothelial cells to advanced glycation end products-induced tissue factor upregulation is associated with intracellular reactive oxygen species. <i>Brain Research</i> , 2006, 1108, 179-187.	2.2	40
318	Advanced glycation end-products downregulating intervertebral disc cell production of proteoglycans in vitro. <i>Journal of Neurosurgery: Spine</i> , 2006, 5, 324-329.	1.7	26
319	Telmisartan, its Potential Therapeutic Implications in Cardiometabolic Disorders. <i>Recent Patents on Cardiovascular Drug Discovery</i> , 2006, 1, 79-83.	1.5	2
320	Cardiovascular Disease in Diabetes. <i>Mini-Reviews in Medicinal Chemistry</i> , 2006, 6, 313-318.	2.4	19
321	Advanced Glycation End Products (AGEs) and their Receptor (RAGE) System in Diabetic Retinopathy. <i>Current Drug Discovery Technologies</i> , 2006, 3, 83-88.	1.2	38
322	Circulating Oxidized Low-density Lipoprotein (LDL) Induces Apoptosis and Stimulates Monocyte Chemoattractant Protein-1 Expression in Cultured Endothelial Cells. <i>Letters in Drug Design and Discovery</i> , 2006, 3, 582-585.	0.7	1
323	Atorvastatin and Diabetic Vascular Complications. <i>Current Pharmaceutical Design</i> , 2006, 12, 1549-1554.	1.9	32
324	Pleiotropic Effects of Nifedipine on Atherosclerosis. <i>Current Pharmaceutical Design</i> , 2006, 12, 1543-1547.	1.9	27

#	ARTICLE	IF	CITATIONS
325	TAGE (Toxic AGEs) Theory in Diabetic Complications. <i>Current Molecular Medicine</i> , 2006, 6, 351-358.	1.3	174
326	Molecular Mechanism for Elevation of Asymmetric Dimethylarginine and Its Role for Hypertension in Chronic Kidney Disease. <i>Journal of the American Society of Nephrology: JASN</i> , 2006, 17, 2176-2183.	6.1	153
327	Increased Serum Concentrations of Pigment Epithelium-Derived Factor in Patients with End-Stage Renal Disease. <i>Clinical Chemistry</i> , 2006, 52, 1970-1971.	3.2	31
328	Role of Advanced Glycation End Products (AGEs) in Thrombogenic Abnormalities in Diabetes. <i>Current Neurovascular Research</i> , 2006, 3, 73-77.	1.1	74
329	Toxic Advanced Glycation End Products (TAGE) Theory in Alzheimer's Disease. <i>American Journal of Alzheimer's Disease and Other Dementias</i> , 2006, 21, 197-208.	1.9	115
330	Pigment Epithelium-derived Factor Inhibits Advanced Glycation End Product-induced Retinal Vascular Hyperpermeability by Blocking Reactive Oxygen Species-mediated Vascular Endothelial Growth Factor Expression. <i>Journal of Biological Chemistry</i> , 2006, 281, 20213-20220.	3.4	194
331	Advanced Glycation End-Products Attenuate Human Mesenchymal Stem Cells and Prevent Cognate Differentiation Into Adipose Tissue, Cartilage, and Bone. <i>Journal of Bone and Mineral Research</i> , 2005, 20, 1647-1658.	2.8	245
332	Pigment epithelium-derived factor (PEDF) blocks angiotensin II signaling in endothelial cells via suppression of NADPH oxidase: a novel anti-oxidative mechanism of PEDF. <i>Cell and Tissue Research</i> , 2005, 320, 437-445.	2.9	108
333	Diabetic Vascular Complications: Pathophysiology, Biochemical Basis and Potential Therapeutic Strategy. <i>Current Pharmaceutical Design</i> , 2005, 11, 2279-2299.	1.9	432
334	Advanced Glycation End Products (AGEs) and Diabetic Vascular Complications. <i>Current Diabetes Reviews</i> , 2005, 1, 93-106.	1.3	141
335	Pigment epithelium-derived factor inhibits oxidative stress-induced apoptosis and dysfunction of cultured retinal pericytes. <i>Microvascular Research</i> , 2005, 69, 45-55.	2.5	75
336	Pigment epithelium-derived factor (PEDF) promotes growth of pericytes through autocrine production of platelet-derived growth factor-B. <i>Microvascular Research</i> , 2005, 69, 128-134.	2.5	18
337	Telmisartan inhibits expression of a receptor for advanced glycation end products (RAGE) in angiotensin-II-exposed endothelial cells and decreases serum levels of soluble RAGE in patients with essential hypertension. <i>Microvascular Research</i> , 2005, 70, 137-141.	2.5	107
338	Pigment epithelium-derived factor (PEDF)-induced apoptosis and inhibition of vascular endothelial growth factor (VEGF) expression in MG63 human osteosarcoma cells. <i>Life Sciences</i> , 2005, 77, 3231-3241.	4.3	91
339	Angiotensin II augments advanced glycation end product-induced pericyte apoptosis through RAGE overexpression. <i>FEBS Letters</i> , 2005, 579, 4265-4270.	2.8	66
340	Therapeutic Implications of Blockers of Advanced Glycation End Products (AGEs)-their Receptor (RAGE) System. <i>International Journal of Pharmacology</i> , 2005, 1, 203-209.	0.3	3
341	Involvement of Advanced Glycation End-products (AGEs) in Alzheimers Disease. <i>Current Alzheimer Research</i> , 2004, 1, 39-46.	1.4	116
342	Azelinidipine, A Newly Developed Long-Acting Calcium Antagonist, Inhibits Tumor Necrosis Factor- α -Induced Interleukin-8 Expression in Endothelial Cells through its Anti-Oxidative Properties. <i>Journal of Cardiovascular Pharmacology</i> , 2004, 43, 724-730.	1.9	68

#	ARTICLE	IF	CITATIONS
343	AGEs activate mesangial TGF- β 2 Smad signaling via an angiotensin II type I receptor interaction. <i>Kidney International</i> , 2004, 66, 2137-2147.	5.2	198
344	AGE down-regulation of monocyte RAGE expression and its association with diabetic complications in type 1 diabetes. <i>Journal of Diabetes and Its Complications</i> , 2004, 18, 53-59.	2.3	22
345	Regulation of Human Melanoma Growth and Metastasis by AGE-AGE Receptor Interactions. <i>Journal of Investigative Dermatology</i> , 2004, 122, 461-467.	0.7	130
346	Pigment epithelium-derived factor inhibits TNF- α -induced interleukin-6 expression in endothelial cells by suppressing NADPH oxidase-mediated reactive oxygen species generation. <i>Journal of Molecular and Cellular Cardiology</i> , 2004, 37, 497-506.	1.9	128
347	Overexpression of Pigment Epithelium-Derived Factor Decreases Angiogenesis and Inhibits the Growth of Human Malignant Melanoma Cells in Vivo. <i>American Journal of Pathology</i> , 2004, 164, 1225-1232.	3.8	119
348	Minodronate, a Newly Developed Nitrogen-Containing Bisphosphonate, Suppresses Melanoma Growth and Improves Survival in Nude Mice by Blocking Vascular Endothelial Growth Factor Signaling. <i>American Journal of Pathology</i> , 2004, 165, 1865-1874.	3.8	82
349	Serum levels of non-carboxymethyllysine advanced glycation endproducts are correlated to severity of microvascular complications in patients with Type 1 diabetes. <i>Journal of Diabetes and Its Complications</i> , 2003, 17, 16-21.	2.3	125
350	Advanced glycation end products inhibit de novo protein synthesis and induce TGF- β 2 overexpression in proximal tubular cells. <i>Kidney International</i> , 2003, 63, 464-473.	5.2	140
351	Pigment epithelium-derived factor inhibits leptin-induced angiogenesis by suppressing vascular endothelial growth factor gene expression through anti-oxidative properties. <i>Microvascular Research</i> , 2003, 65, 186-190.	2.5	97
352	Neurotoxicity of Acetaldehyde-Derived Advanced Glycation End Products for Cultured Cortical Neurons. <i>Journal of Neuropathology and Experimental Neurology</i> , 2003, 62, 486-496.	1.7	32
353	Angiogenesis induced by advanced glycation end products and its prevention by cerivastatin. <i>FASEB Journal</i> , 2002, 16, 1928-1930.	0.5	247
354	Advanced Glycation End Product-induced Apoptosis and Overexpression of Vascular Endothelial Growth Factor and Monocyte Chemoattractant Protein-1 in Human-cultured Mesangial Cells. <i>Journal of Biological Chemistry</i> , 2002, 277, 20309-20315.	3.4	275
355	Advanced Glycation End Products-Induced Apoptosis and Overexpression of Vascular Endothelial Growth Factor in Bovine Retinal Pericytes. <i>Biochemical and Biophysical Research Communications</i> , 2002, 290, 973-978.	2.1	204
356	Pigment epithelium-derived factor protects cultured retinal pericytes from advanced glycation end product-induced injury through its antioxidative properties. <i>Biochemical and Biophysical Research Communications</i> , 2002, 296, 877-882.	2.1	178
357	Incadronate disodium inhibits advanced glycation end products-induced angiogenesis in vitro. <i>Biochemical and Biophysical Research Communications</i> , 2002, 297, 419-424.	2.1	55
358	Sorbitol dehydrogenase overexpression potentiates glucose toxicity to cultured retinal pericytes. <i>Biochemical and Biophysical Research Communications</i> , 2002, 299, 183-188.	2.1	56
359	Interferon- β -induced apoptosis and activation of THP-1 macrophages. <i>Life Sciences</i> , 2002, 71, 2499-2508.	4.3	35
360	Advanced glycation end products (AGE) and their receptor (RAGE) in the brain of patients with Creutzfeldt-Jakob disease with prion plaques.. <i>Neuroscience Letters</i> , 2002, 326, 117-120.	2.1	97

#	ARTICLE	IF	CITATIONS
361	Palmitate-Induced Apoptosis of Microvascular Endothelial Cells and Pericytes. <i>Molecular Medicine</i> , 2002, 8, 179-184.	4.4	71
362	CS-866, a New Angiotensin II Type 1 Receptor Antagonist, Ameliorates Glomerular Anionic Site Loss and Prevents Progression of Diabetic Nephropathy in Otsuka Long-Evans Tokushima Fatty Rats. <i>Molecular Medicine</i> , 2002, 8, 591-599.	4.4	23
363	Beraprost Sodium, a Prostaglandin I2 Analogue, Protects Against Advanced Glycation End Products-induced Injury in Cultured Retinal Pericytes. <i>Molecular Medicine</i> , 2002, 8, 546-550.	4.4	52
364	Palmitate-induced apoptosis of microvascular endothelial cells and pericytes. <i>Molecular Medicine</i> , 2002, 8, 179-84.	4.4	24
365	Beraprost sodium, a prostaglandin I2 analogue, protects against advanced glycation end products-induced injury in cultured retinal pericytes. <i>Molecular Medicine</i> , 2002, 8, 546-50.	4.4	13
366	Immunological Detection of a Novel Advanced Glycation End-Product. <i>Molecular Medicine</i> , 2001, 7, 783-791.	4.4	85
367	Leptin Induces Mitochondrial Superoxide Production and Monocyte Chemoattractant Protein-1 Expression in Aortic Endothelial Cells by Increasing Fatty Acid Oxidation via Protein Kinase A. <i>Journal of Biological Chemistry</i> , 2001, 276, 25096-25100.	3.4	530
368	Hyperglycemia Potentiates Collagen-Induced Platelet Activation Through Mitochondrial Superoxide Overproduction. <i>Diabetes</i> , 2001, 50, 1491-1494.	0.6	199
369	Development and prevention of advanced diabetic nephropathy in RAGE-overexpressing mice. <i>Journal of Clinical Investigation</i> , 2001, 108, 261-268.	8.2	430
370	Normalizing mitochondrial superoxide production blocks three pathways of hyperglycaemic damage. <i>Nature</i> , 2000, 404, 787-790.	27.8	3,895
371	Deficient cotinine formation from nicotine is attributed to the whole deletion of the CYP2A6 gene in humans. <i>Clinical Pharmacology and Therapeutics</i> , 2000, 67, 57-69.	4.7	101
372	The Receptor for Advanced Glycation End Products Is Induced by the Glycation Products Themselves and Tumor Necrosis Factor- α through Nuclear Factor- κ B, and by 17 β -Estradiol through Sp-1 in Human Vascular Endothelial Cells. <i>Journal of Biological Chemistry</i> , 2000, 275, 25781-25790.	3.4	383
373	Insulin Stimulates the Growth and Tube Formation of Human Microvascular Endothelial Cells through Autocrine Vascular Endothelial Growth Factor. <i>Microvascular Research</i> , 1999, 57, 329-339.	2.5	61
374	Advanced Glycation Endproducts Accelerate Calcification in Microvascular Pericytes. <i>Biochemical and Biophysical Research Communications</i> , 1999, 258, 353-357.	2.1	103
375	Upregulation of Retinal Vascular Endothelial Growth Factor mRNAs in Spontaneously Diabetic Rats without Ophthalmoscopic Retinopathy. <i>Ophthalmic Research</i> , 1998, 30, 333-339.	1.9	63
376	Advanced Glycation End Products-driven Angiogenesis in Vitro. <i>Journal of Biological Chemistry</i> , 1997, 272, 8723-8730.	3.4	253
377	Advanced glycosylation end products stimulate the growth but inhibit the prostacyclin-producing ability of endothelial cells through interactions with their receptors. <i>FEBS Letters</i> , 1996, 384, 103-106.	2.8	48
378	Advanced Glycation Endproducts Receptor Interactions Stimulate the Growth of Human Pancreatic Cancer Cells through the Induction of Platelet-Derived Growth Factor-B. <i>Biochemical and Biophysical Research Communications</i> , 1996, 222, 700-705.	2.1	46

#	ARTICLE	IF	CITATIONS
379	Possible Participation of Autocrine and Paracrine Vascular Endothelial Growth Factors in Hypoxia-induced Proliferation of Endothelial Cells and Pericytes. <i>Journal of Biological Chemistry</i> , 1995, 270, 28316-28324.	3.4	259