

Dong-Won Kang

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7249945/publications.pdf>

Version: 2024-02-01

14
papers

824
citations

759233

12
h-index

1125743

13
g-index

14
all docs

14
docs citations

14
times ranked

1262
citing authors

#	ARTICLE	IF	CITATIONS
1	Depressive behaviors and decreased expression of serotonin reuptake transporter in rats that experienced neonatal maternal separation. <i>Neuroscience Research</i> , 2007, 58, 32-39.	1.9	198
2	The Inhibitory Effects of Eckol and Dieckol from <i>Ecklonia stolonifera</i> on the Expression of Matrix Metalloproteinase-1 in Human Dermal Fibroblasts. <i>Biological and Pharmaceutical Bulletin</i> , 2006, 29, 1735-1739.	1.4	135
3	Chronic food restriction in young rats results in depression- and anxiety-like behaviors with decreased expression of serotonin reuptake transporter. <i>Brain Research</i> , 2007, 1150, 100-107.	2.2	91
4	Post-weaning isolation promotes food intake and body weight gain in rats that experienced neonatal maternal separation. <i>Brain Research</i> , 2009, 1295, 127-134.	2.2	71
5	Dexamethasone reduces food intake, weight gain and the hypothalamic 5-HT concentration and increases plasma leptin in rats. <i>European Journal of Pharmacology</i> , 2008, 581, 64-70.	3.5	69
6	Reversine stimulates adipocyte differentiation and downregulates Akt and p70s6k signaling pathways in 3T3-L1 cells. <i>Biochemical and Biophysical Research Communications</i> , 2007, 358, 553-558.	2.1	51
7	The 1p-Encoded Protein Stathmin and Resistance of Malignant Gliomas to Nitrosoureas. <i>Journal of the National Cancer Institute</i> , 2007, 99, 639-652.	6.3	49
8	Adolescence fluoxetine increases serotonergic activity in the raphe-hippocampus axis and improves depression-like behaviors in female rats that experienced neonatal maternal separation. <i>Psychoneuroendocrinology</i> , 2013, 38, 777-788.	2.7	46
9	Involvement of HDAC1 and the PI3K/PKC signaling pathways in NF- κ B activation by the HDAC inhibitor apicidin. <i>Biochemical and Biophysical Research Communications</i> , 2006, 347, 1088-1093.	2.1	30
10	Reversal of multidrug resistance by 4-chloro-N-(3-((E)-3-(4-hydroxy-3-methoxyphenyl)acryloyl)phenyl)benzamide through the reversible inhibition of P-glycoprotein. <i>Biochemical and Biophysical Research Communications</i> , 2007, 355, 136-142.	2.1	27
11	Apoptotic potential of sesquiterpene lactone ergolide through the inhibition of NF- κ B signaling pathway. <i>Journal of Pharmacy and Pharmacology</i> , 2010, 57, 1591-1597.	2.4	26
12	The Ketogenic Diet Suppresses the Cathepsin E Expression Induced by Kainic Acid in the Rat Brain. <i>Yonsei Medical Journal</i> , 2010, 51, 653.	2.2	19
13	Regulation of SLC26A3 activity by NHERF4 PDZ-mediated interaction. <i>Cellular Signalling</i> , 2012, 24, 1821-1830.	3.6	11
14	Building up partnership with members of medical society of Zimbabwe to increase their potential to help vulnerable people in healthcare. <i>Journal of Global Health Science</i> , 2020, 2, .	0.3	1