

Hai-Qiang Mai

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7283139/publications.pdf>

Version: 2024-02-01

144
papers

5,831
citations

109321

35
h-index

91884

69
g-index

149
all docs

149
docs citations

149
times ranked

4293
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|--|------|-----------|
| 1 | Chemotherapy and radiotherapy in nasopharyngeal carcinoma: an update of the MAC-NPC meta-analysis. <i>Lancet Oncology</i> , The, 2015, 16, 645-655. | 10.7 | 593 |
| 2 | Concurrent Chemoradiotherapy vs Radiotherapy Alone in Stage II Nasopharyngeal Carcinoma: Phase III Randomized Trial. <i>Journal of the National Cancer Institute</i> , 2011, 103, 1761-1770. | 6.3 | 286 |
| 3 | Establishment and Validation of Prognostic Nomograms for Endemic Nasopharyngeal Carcinoma. <i>Journal of the National Cancer Institute</i> , 2016, 108, djv291. | 6.3 | 281 |
| 4 | What Is the Best Treatment of Locally Advanced Nasopharyngeal Carcinoma? An Individual Patient Data Network Meta-Analysis. <i>Journal of Clinical Oncology</i> , 2017, 35, 498-505. | 1.6 | 263 |
| 5 | Long-Term Survival After Cisplatin-Based Induction Chemotherapy and Radiotherapy for Nasopharyngeal Carcinoma: A Pooled Data Analysis of Two Phase III Trials. <i>Journal of Clinical Oncology</i> , 2005, 23, 1118-1124. | 1.6 | 227 |
| 6 | Neoadjuvant chemotherapy followed by concurrent chemoradiotherapy versus concurrent chemoradiotherapy alone in locoregionally advanced nasopharyngeal carcinoma: A phase III multicentre randomised controlled trial. <i>European Journal of Cancer</i> , 2017, 75, 14-23. | 2.8 | 226 |
| 7 | Toripalimab or placebo plus chemotherapy as first-line treatment in advanced nasopharyngeal carcinoma: a multicenter randomized phase 3 trial. <i>Nature Medicine</i> , 2021, 27, 1536-1543. | 30.7 | 197 |
| 8 | Prospective Study of Tailoring Whole-Body Dual-Modality [¹⁸ F]Fluorodeoxyglucose Positron Emission Tomography/Computed Tomography With Plasma Epstein-Barr Virus DNA for Detecting Distant Metastasis in Endemic Nasopharyngeal Carcinoma at Initial Staging. <i>Journal of Clinical Oncology</i> , 2013, 31, 2861-2869. | 1.6 | 171 |
| 9 | Induction chemotherapy followed by concurrent chemoradiotherapy versus concurrent chemoradiotherapy alone in locoregionally advanced nasopharyngeal carcinoma: long-term results of a phase III multicentre randomised controlled trial. <i>European Journal of Cancer</i> , 2019, 119, 87-96. | 2.8 | 150 |
| 10 | Efficacy and Safety of Locoregional Radiotherapy With Chemotherapy vs Chemotherapy Alone in De Novo Metastatic Nasopharyngeal Carcinoma. <i>JAMA Oncology</i> , 2020, 6, 1345. | 7.1 | 137 |
| 11 | The Chinese Society of Clinical Oncology (CSCO) clinical guidelines for the diagnosis and treatment of nasopharyngeal carcinoma. <i>Cancer Communications</i> , 2021, 41, 1195-1227. | 9.2 | 128 |
| 12 | Concurrent chemoradiotherapy with nedaplatin versus cisplatin in stage II-IVB nasopharyngeal carcinoma: an open-label, non-inferiority, randomised phase 3 trial. <i>Lancet Oncology</i> , The, 2018, 19, 461-473. | 10.7 | 118 |
| 13 | Single-cell transcriptomic analysis defines the interplay between tumor cells, viral infection, and the microenvironment in nasopharyngeal carcinoma. <i>Cell Research</i> , 2020, 30, 950-965. | 12.0 | 111 |
| 14 | The Prognostic Value of Plasma Epstein-Barr Viral DNA and Tumor Response to Neoadjuvant Chemotherapy in Advanced-Stage Nasopharyngeal Carcinoma. <i>International Journal of Radiation Oncology Biology Physics</i> , 2015, 93, 862-869. | 0.8 | 110 |
| 15 | Locoregional radiotherapy in patients with distant metastases of nasopharyngeal carcinoma at diagnosis. <i>Chinese Journal of Cancer</i> , 2013, 32, 604-613. | 4.9 | 109 |
| 16 | Tumour heterogeneity and intercellular networks of nasopharyngeal carcinoma at single cell resolution. <i>Nature Communications</i> , 2021, 12, 741. | 12.8 | 104 |
| 17 | A new prognostic histopathologic classification of nasopharyngeal carcinoma. <i>Chinese Journal of Cancer</i> , 2016, 35, 41. | 4.9 | 83 |
| 18 | STING signaling remodels the tumor microenvironment by antagonizing myeloid-derived suppressor cell expansion. <i>Cell Death and Differentiation</i> , 2019, 26, 2314-2328. | 11.2 | 81 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 19 | Genome-Wide Identification of a Methylation Gene Panel as a Prognostic Biomarker in Nasopharyngeal Carcinoma. <i>Molecular Cancer Therapeutics</i> , 2015, 14, 2864-2873. | 4.1 | 80 |
| 20 | Establishment and validation of M1 stage subdivisions for de novo metastatic nasopharyngeal carcinoma to better predict prognosis and guide treatment. <i>European Journal of Cancer</i> , 2017, 77, 117-126. | 2.8 | 80 |
| 21 | The Association Between the Development of Radiation Therapy, Image Technology, and Chemotherapy, and the Survival of Patients With Nasopharyngeal Carcinoma: A Cohort Study From 1990 to 2012. <i>International Journal of Radiation Oncology Biology Physics</i> , 2019, 105, 581-590. | 0.8 | 80 |
| 22 | Tumor CTLA-4 overexpression predicts poor survival in patients with nasopharyngeal carcinoma. <i>Oncotarget</i> , 2016, 7, 13060-13068. | 1.8 | 80 |
| 23 | A randomized trial of induction chemotherapy plus concurrent chemoradiotherapy versus induction chemotherapy plus radiotherapy for locoregionally advanced nasopharyngeal carcinoma. <i>Oral Oncology</i> , 2012, 48, 1038-1044. | 1.5 | 65 |
| 24 | Phase I trial of adoptively transferred tumor-infiltrating lymphocyte immunotherapy following concurrent chemoradiotherapy in patients with locoregionally advanced nasopharyngeal carcinoma. <i>Oncolmmunology</i> , 2015, 4, e976507. | 4.6 | 61 |
| 25 | Salvage endoscopic nasopharyngectomy and intensity-modulated radiotherapy versus conventional radiotherapy in treating locally recurrent nasopharyngeal carcinoma. <i>Head and Neck</i> , 2015, 37, 1108-1115. | 2.0 | 59 |
| 26 | The Prognostic Value of Treatment-Related Lymphopenia in Nasopharyngeal Carcinoma Patients. <i>Cancer Research and Treatment</i> , 2018, 50, 19-29. | 3.0 | 56 |
| 27 | Galectin-9 promotes a suppressive microenvironment in human cancer by enhancing STING degradation. <i>Oncogenesis</i> , 2020, 9, 65. | 4.9 | 52 |
| 28 | SPINK6 Promotes Metastasis of Nasopharyngeal Carcinoma via Binding and Activation of Epithelial Growth Factor Receptor. <i>Cancer Research</i> , 2017, 77, 579-589. | 0.9 | 47 |
| 29 | The <i>RARS</i> – <i>MAD1L1</i> Fusion Gene Induces Cancer Stem Cell-like Properties and Therapeutic Resistance in Nasopharyngeal Carcinoma. <i>Clinical Cancer Research</i> , 2018, 24, 659-673. | 7.0 | 47 |
| 30 | Elevated peripheral blood lymphocyte-to-monocyte ratio predicts a favorable prognosis in the patients with metastatic nasopharyngeal carcinoma. <i>Chinese Journal of Cancer</i> , 2015, 34, 237-46. | 4.9 | 44 |
| 31 | Optimal cumulative cisplatin dose in nasopharyngeal carcinoma patients based on induction chemotherapy response. <i>Radiotherapy and Oncology</i> , 2019, 137, 83-94. | 0.6 | 44 |
| 32 | Ten-year outcomes of a randomised trial for locoregionally advanced nasopharyngeal carcinoma: A single-institution experience from an endemic area. <i>European Journal of Cancer</i> , 2015, 51, 1760-1770. | 2.8 | 43 |
| 33 | High-Sensitivity C-Reactive Protein Complements Plasma Epstein-Barr Virus Deoxyribonucleic Acid Prognostication in Nasopharyngeal Carcinoma: A Large-Scale Retrospective and Prospective Cohort Study. <i>International Journal of Radiation Oncology Biology Physics</i> , 2015, 91, 325-336. | 0.8 | 41 |
| 34 | Ten-year outcomes of survival and toxicity for a phase III randomised trial of concurrent chemoradiotherapy versus radiotherapy alone in stage II nasopharyngeal carcinoma. <i>European Journal of Cancer</i> , 2019, 110, 24-31. | 2.8 | 40 |
| 35 | Combination of Tumor Volume and Epstein-Barr Virus DNA Improved Prognostic Stratification of Stage II Nasopharyngeal Carcinoma in the Intensity Modulated Radiotherapy Era: A Large-Scale Cohort Study. <i>Cancer Research and Treatment</i> , 2018, 50, 861-871. | 3.0 | 38 |
| 36 | Plasma Epstein-Barr viral DNA complements TNM classification of nasopharyngeal carcinoma in the era of intensity-modulated radiotherapy. <i>Oncotarget</i> , 2016, 7, 6221-6230. | 1.8 | 37 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 37 | New surgical staging system for patients with recurrent nasopharyngeal carcinoma based on the AJCC/UICC rTNM classification system. <i>European Journal of Cancer</i> , 2015, 51, 1771-1779. | 2.8 | 36 |
| 38 | Identifying optimal candidates for local treatment of the primary tumor among patients with de novo metastatic nasopharyngeal carcinoma: a retrospective cohort study based on Epstein-Barr virus DNA level and tumor response to palliative chemotherapy. <i>BMC Cancer</i> , 2019, 19, 92. | 2.6 | 33 |
| 39 | High-density lipoprotein cholesterol as a predictor of poor survival in patients with nasopharyngeal carcinoma. <i>Oncotarget</i> , 2016, 7, 42978-42987. | 1.8 | 32 |
| 40 | Elevated plasma big ET-1 is associated with distant failure in patients with advanced-stage nasopharyngeal carcinoma. <i>Cancer</i> , 2006, 106, 1548-1553. | 4.1 | 31 |
| 41 | Comparison of Long-Term Survival and Toxicity of Cisplatin Delivered Weekly versus Every Three Weeks Concurrently with Intensity-Modulated Radiotherapy in Nasopharyngeal Carcinoma. <i>PLoS ONE</i> , 2014, 9, e110765. | 2.5 | 31 |
| 42 | Elevated levels of plasma D-dimer predict a worse outcome in patients with nasopharyngeal carcinoma. <i>BMC Cancer</i> , 2014, 14, 583. | 2.6 | 29 |
| 43 | Prognostic Value of Plasma Epstein-Barr Virus DNA for Local and Regionally Advanced Nasopharyngeal Carcinoma Treated With Cisplatin-Based Concurrent Chemoradiotherapy in Intensity-Modulated Radiotherapy Era. <i>Medicine (United States)</i> , 2016, 95, e2642. | 1.0 | 29 |
| 44 | Concurrent chemoradiotherapy with or without cetuximab for stage II to IVb nasopharyngeal carcinoma: a case-control study. <i>BMC Cancer</i> , 2017, 17, 567. | 2.6 | 29 |
| 45 | Is Hemoglobin Level in Patients with Nasopharyngeal Carcinoma Still a Significant Prognostic Factor in the Era of Intensity-Modulated Radiotherapy Technology?. <i>PLoS ONE</i> , 2015, 10, e0136033. | 2.5 | 28 |
| 46 | Expression of ERCC1 predicts clinical outcome in locoregionally advanced nasopharyngeal carcinoma treated with cisplatin-based induction chemotherapy. <i>Oral Oncology</i> , 2012, 48, 964-968. | 1.5 | 27 |
| 47 | Different Prognostic Values of Plasma Epstein-Barr Virus DNA and Maximal Standardized Uptake Value of 18F-FDG PET/CT for Nasopharyngeal Carcinoma Patients with Recurrence. <i>PLoS ONE</i> , 2015, 10, e0122756. | 2.5 | 27 |
| 48 | CDC42-interacting protein 4 promotes metastasis of nasopharyngeal carcinoma by mediating invadopodia formation and activating EGFR signaling. <i>Journal of Experimental and Clinical Cancer Research</i> , 2017, 36, 21. | 8.6 | 26 |
| 49 | Proposal for a new risk classification system for nasopharyngeal carcinoma patients with post-radiation nasopharyngeal necrosis. <i>Oral Oncology</i> , 2017, 67, 83-88. | 1.5 | 26 |
| 50 | Development and validation of the immune signature to predict distant metastasis in patients with nasopharyngeal carcinoma. , 2020, 8, e000205. | | 26 |
| 51 | Deintensified Chemoradiotherapy for Pretreatment Epstein-Barr Virus DNA-Selected Low-Risk Locoregionally Advanced Nasopharyngeal Carcinoma: A Phase II Randomized Noninferiority Trial. <i>Journal of Clinical Oncology</i> , 2022, 40, 1163-1173. | 1.6 | 25 |
| 52 | Prognostic implications of dynamic serum lactate dehydrogenase assessments in nasopharyngeal carcinoma patients treated with intensity-modulated radiotherapy. <i>Scientific Reports</i> , 2016, 6, 22326. | 3.3 | 24 |
| 53 | FMNL1 mediates nasopharyngeal carcinoma cell aggressiveness by epigenetically upregulating MTA1. <i>Oncogene</i> , 2018, 37, 6243-6258. | 5.9 | 24 |
| 54 | The diagnostic and prognostic values of plasma Epstein-Barr virus DNA for residual cervical lymphadenopathy in nasopharyngeal carcinoma patients: a retrospective study. <i>Cancer Communications</i> , 2019, 39, 1-13. | 9.2 | 24 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 55 | Advances in pathogenesis and precision medicine for nasopharyngeal carcinoma. <i>MedComm</i> , 2021, 2, 175-206. | 7.2 | 24 |
| 56 | Construction of a comprehensive nutritional index and its correlation with quality of life and survival in patients with nasopharyngeal carcinoma undergoing IMRT: A prospective study. <i>Oral Oncology</i> , 2019, 98, 62-68. | 1.5 | 23 |
| 57 | Combining pretreatment plasma Epstein-Barr virus DNA level and cervical node necrosis improves prognostic stratification in patients with nasopharyngeal carcinoma: A cohort study. <i>Cancer Medicine</i> , 2019, 8, 6841-6852. | 2.8 | 22 |
| 58 | Effect of Induction Chemotherapy With Paclitaxel, Cisplatin, and Capecitabine vs Cisplatin and Fluorouracil on Failure-Free Survival for Patients With Stage IVA to IVB Nasopharyngeal Carcinoma. <i>JAMA Oncology</i> , 2022, 8, 706. | 7.1 | 22 |
| 59 | The impact of the cumulative dose of cisplatin during concurrent chemoradiotherapy on the clinical outcomes of patients with advanced-stage nasopharyngeal carcinoma in an era of intensity-modulated radiotherapy. <i>BMC Cancer</i> , 2015, 15, 977. | 2.6 | 21 |
| 60 | Elevated High-Sensitivity C-Reactive Protein Levels Predict Decreased Survival for Nasopharyngeal Carcinoma Patients in the Intensity-Modulated Radiotherapy Era. <i>PLoS ONE</i> , 2015, 10, e0122965. | 2.5 | 21 |
| 61 | A curative-intent endoscopic surgery for postradiation nasopharyngeal necrosis in patients with nasopharyngeal carcinoma. <i>Cancer Communications</i> , 2018, 38, 1-11. | 9.2 | 21 |
| 62 | Establishment of an innovative staging system for extramedullary plasmacytoma. <i>BMC Cancer</i> , 2016, 16, 777. | 2.6 | 20 |
| 63 | Beneficial effects of anti-EGFR agents, Cetuximab or Nimotuzumab, in combination with concurrent chemoradiotherapy in advanced nasopharyngeal carcinoma. <i>Oral Oncology</i> , 2018, 80, 1-8. | 1.5 | 20 |
| 64 | Surgery for isolated regional failure in nasopharyngeal carcinoma after radiation: Selective or comprehensive neck dissection. <i>Laryngoscope</i> , 2019, 129, 387-395. | 2.0 | 20 |
| 65 | Therapeutic targeting of the endothelin receptor in human nasopharyngeal carcinoma. <i>Cancer Science</i> , 2006, 97, 1388-1395. | 3.9 | 19 |
| 66 | Targeting the IRAK1-S100A9 Axis Overcomes Resistance to Paclitaxel in Nasopharyngeal Carcinoma. <i>Cancer Research</i> , 2021, 81, 1413-1425. | 0.9 | 19 |
| 67 | Concurrent chemoradiotherapy with 3-weekly versus weekly cisplatin in patients with locoregionally advanced nasopharyngeal carcinoma: A phase 3 multicentre randomised controlled trial (ChiCTR-TRC-12001979).. <i>Journal of Clinical Oncology</i> , 2017, 35, 6006-6006. | 1.6 | 19 |
| 68 | Induction Chemotherapy Plus Concurrent Chemoradiotherapy Versus Concurrent Chemoradiotherapy Alone in Locoregionally Advanced Nasopharyngeal Carcinoma in Children and Adolescents: A Matched Cohort Analysis. <i>Cancer Research and Treatment</i> , 2018, 50, 1304-1315. | 3.0 | 19 |
| 69 | Efficacy of controlled-release oxycodone for reducing pain due to oral mucositis in nasopharyngeal carcinoma patients treated with concurrent chemoradiotherapy: a prospective clinical trial. <i>Supportive Care in Cancer</i> , 2019, 27, 3759-3767. | 2.2 | 18 |
| 70 | Subdivision of Nasopharyngeal Carcinoma Patients with Bone-Only Metastasis at Diagnosis for Prediction of Survival and Treatment Guidance. <i>Cancer Research and Treatment</i> , 2019, 51, 1259-1268. | 3.0 | 18 |
| 71 | Expression of EIF5A2 associates with poor survival of nasopharyngeal carcinoma patients treated with induction chemotherapy. <i>BMC Cancer</i> , 2016, 16, 669. | 2.6 | 17 |
| 72 | With or without reirradiation in advanced local recurrent nasopharyngeal carcinoma: a case-control study. <i>BMC Cancer</i> , 2016, 16, 774. | 2.6 | 17 |

| # | ARTICLE | IF | CITATIONS |
|----|--|------|-----------|
| 73 | Effect of local treatment for metastasis and its sequence with chemotherapy on prognosis of post-treatment metastatic nasopharyngeal carcinoma patients. <i>Oral Oncology</i> , 2019, 92, 40-45. | 1.5 | 17 |
| 74 | Deep learning signatures reveal multiscale intratumor heterogeneity associated with biological functions and survival in recurrent nasopharyngeal carcinoma. <i>European Journal of Nuclear Medicine and Molecular Imaging</i> , 2022, 49, 2972-2982. | 6.4 | 17 |
| 75 | Endoscopic microwave coagulation therapy for early recurrent T1 nasopharyngeal carcinoma. <i>European Journal of Cancer</i> , 2009, 45, 1107-1110. | 2.8 | 16 |
| 76 | Endoscope-Guided Interstitial Intensity-Modulated Brachytherapy and Intracavitary Brachytherapy as Boost Radiation for Primary Early T Stage Nasopharyngeal Carcinoma. <i>PLoS ONE</i> , 2014, 9, e90048. | 2.5 | 16 |
| 77 | The role of capecitabine as maintenance therapy in <i>de novo</i> metastatic nasopharyngeal carcinoma: A propensity score matching study. <i>Cancer Communications</i> , 2020, 40, 32-42. | 9.2 | 16 |
| 78 | The impact of smoking on the clinical outcome of locoregionally advanced nasopharyngeal carcinoma after chemoradiotherapy. <i>Radiation Oncology</i> , 2014, 9, 246. | 2.7 | 15 |
| 79 | Induction chemotherapy followed by concurrent chemoradiotherapy versus concurrent chemoradiotherapy alone in stage III-IVb nasopharyngeal carcinoma patients with Epstein-Barr virus DNA ≥ 4000 copies/ml: a matched study. <i>Oncotarget</i> , 2016, 7, 29739-29748. | 1.8 | 15 |
| 80 | A Randomized Controlled Trial Comparing Two Different Schedules for Cisplatin Treatment in Patients with Locoregionally Advanced Nasopharyngeal Cancer. <i>Clinical Cancer Research</i> , 2021, 27, 4186-4194. | 7.0 | 15 |
| 81 | RBFOX2/GOLIM4 Splicing Axis Activates Vesicular Transport Pathway to Promote Nasopharyngeal Carcinogenesis. <i>Advanced Science</i> , 2021, 8, e2004852. | 11.2 | 15 |
| 82 | Comparison of the treatment outcomes of intensity-modulated radiotherapy and two-dimensional conventional radiotherapy in nasopharyngeal carcinoma patients with parapharyngeal space extension. <i>Radiotherapy and Oncology</i> , 2015, 116, 167-173. | 0.6 | 14 |
| 83 | Pretreatment Serum Amyloid A and C-reactive Protein Comparing with Epstein-Barr Virus DNA as Prognostic Indicators in Patients with Nasopharyngeal Carcinoma: A Prospective Study. <i>Cancer Research and Treatment</i> , 2018, 50, 701-711. | 3.0 | 14 |
| 84 | LOX expression in primary nasopharyngeal carcinoma: correlation with prognostic parameters and outcome. <i>Oncotarget</i> , 2016, 7, 8200-8207. | 1.8 | 14 |
| 85 | Liposomal paclitaxel versus docetaxel in induction chemotherapy using Taxanes, cisplatin and 5-fluorouracil for locally advanced nasopharyngeal carcinoma. <i>BMC Cancer</i> , 2018, 18, 1279. | 2.6 | 13 |
| 86 | Pretreatment quality of life as a predictor of survival for patients with nasopharyngeal carcinoma treated with IMRT. <i>BMC Cancer</i> , 2018, 18, 114. | 2.6 | 13 |
| 87 | <p>The development of a nomogram to predict post-radiation necrosis in nasopharyngeal carcinoma patients: a large-scale cohort study</p>. <i>Cancer Management and Research</i> , 2019, Volume 11, 6253-6263. | 1.9 | 13 |
| 88 | Establishment and validation of a nomogram for predicting survival in patients with de novo metastatic nasopharyngeal carcinoma. <i>Oral Oncology</i> , 2019, 94, 73-79. | 1.5 | 12 |
| 89 | Establishment and validation of a nomogram for predicting the benefit of concurrent chemotherapy in stage II nasopharyngeal carcinoma: A study based on a phase III randomized clinical trial with 10-year follow-up. <i>Oral Oncology</i> , 2020, 100, 104490. | 1.5 | 12 |
| 90 | Genetic Variations in Radiation and Chemotherapy Drug Action Pathways and Survival in Locoregionally Advanced Nasopharyngeal Carcinoma Treated with Chemoradiotherapy. <i>PLoS ONE</i> , 2013, 8, e82750. | 2.5 | 12 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 91 | Advanced-Stage Nasopharyngeal Carcinoma: Restaging System after Neoadjuvant Chemotherapy on the Basis of MR Imaging Determines Survival. <i>Radiology</i> , 2017, 282, 171-181. | 7.3 | 11 |
| 92 | Patterns of Failure and Survival Trends Of 720 Patients with Stage I Nasopharyngeal Carcinoma Diagnosed from 1990-2012: A Large-scale Retrospective Cohort Study. <i>Journal of Cancer</i> , 2018, 9, 1308-1317. | 2.5 | 11 |
| 93 | The impact of Adult Comorbidity Evaluation-27 on the clinical outcome of elderly nasopharyngeal carcinoma patients treated with chemoradiotherapy or radiotherapy: a matched cohort analysis. <i>Journal of Cancer</i> , 2019, 10, 5614-5621. | 2.5 | 11 |
| 94 | Subdivision of de-novo metastatic nasopharyngeal carcinoma based on tumor burden and pretreatment EBV DNA for therapeutic guidance of locoregional radiotherapy. <i>BMC Cancer</i> , 2021, 21, 534. | 2.6 | 11 |
| 95 | Patterns of Failure and Survival Trends in 3,808 Patients with Stage II Nasopharyngeal Carcinoma Diagnosed from 1990 to 2012: A Large-Scale Retrospective Cohort Study. <i>Cancer Research and Treatment</i> , 2019, 51, 1449-1463. | 3.0 | 11 |
| 96 | Stratification of Candidates for Induction Chemotherapy in Stage III-IV Nasopharyngeal Carcinoma: A Large Cohort Study Based on a Comprehensive Prognostic Model. <i>Frontiers in Oncology</i> , 2020, 10, 255. | 2.8 | 10 |
| 97 | Establishment of a prognostic nomogram to identify optimal candidates for local treatment among patients with local recurrent nasopharyngeal carcinoma. <i>Oral Oncology</i> , 2020, 106, 104711. | 1.5 | 10 |
| 98 | Combining plasma Epstein-Barr virus DNA and nodal maximal standard uptake values of 18F-fluoro-2-deoxy-D-glucose positron emission tomography improved prognostic stratification to predict distant metastasis for locoregionally advanced nasopharyngeal carcinoma. <i>Oncotarget</i> , 2015, 6, 38296-38307. | 1.8 | 10 |
| 99 | Identifying distinct risks of treatment failure in nasopharyngeal carcinoma: A study based on the dynamic changes in peripheral blood lymphocytes, monocytes, N classification, and plasma Epstein-Barr virus DNA. <i>Head and Neck</i> , 2021, , . | 2.0 | 10 |
| 100 | Development and validation of a transcriptomics-based gene signature to predict distant metastasis and guide induction chemotherapy in locoregionally advanced nasopharyngeal carcinoma. <i>European Journal of Cancer</i> , 2022, 163, 26-34. | 2.8 | 10 |
| 101 | Functional polymorphisms of matrix metalloproteinase-9 and survival in patients with locoregionally advanced nasopharyngeal carcinoma treated with chemoradiotherapy. <i>Medical Oncology</i> , 2013, 30, 685. | 2.5 | 9 |
| 102 | Effect of Concurrent Chemoradiotherapy With Nedaplatin vs Cisplatin on the Long-term Outcomes of Survival and Toxic Effects Among Patients With Stage II to IVB Nasopharyngeal Carcinoma. <i>JAMA Network Open</i> , 2021, 4, e2138470. | 5.9 | 9 |
| 103 | Pregnancy associated nasopharyngeal carcinoma: A retrospective case-control analysis of maternal survival outcomes. <i>Radiotherapy and Oncology</i> , 2015, 116, 125-130. | 0.6 | 8 |
| 104 | Association between XRCC3 Thr241Met polymorphism and nasopharyngeal carcinoma risk: evidence from a large-scale case-control study and a meta-analysis. <i>Tumor Biology</i> , 2016, 37, 14825-14830. | 1.8 | 8 |
| 105 | Maximal standard uptake values of 18F-fluoro-2-deoxy-D-glucose positron emission tomography compared with Epstein-Barr virus DNA as prognostic indicators in de novo metastatic nasopharyngeal carcinoma patients. <i>BMC Cancer</i> , 2019, 19, 908. | 2.6 | 8 |
| 106 | Establishment and validation of two nomograms to predict the benefit of concurrent chemotherapy in stage II-IVa nasopharyngeal carcinoma patients with different risk factors: Analysis based on a large cohort. <i>Cancer Medicine</i> , 2020, 9, 1661-1670. | 2.8 | 8 |
| 107 | Comparing three induction chemotherapy regimens for patients with locoregionally advanced nasopharyngeal carcinoma based on TNM stage and plasma Epstein-Barr virus DNA level. <i>BMC Cancer</i> , 2020, 20, 89. | 2.6 | 8 |
| 108 | Geriatric nutritional risk index as an independent prognostic factor in locally advanced nasopharyngeal carcinoma treated using radical concurrent chemoradiotherapy: a retrospective cohort study. <i>Annals of Translational Medicine</i> , 2021, 9, 532-532. | 1.7 | 8 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 109 | Radiation-induced hypothyroidism in patients with nasopharyngeal carcinoma treated with intensity-modulated radiation therapy with or without chemotherapy: Development of a nomogram based on the equivalent dose. <i>Oral Oncology</i> , 2021, 120, 105378. | 1.5 | 8 |
| 110 | Nomogram for the prediction of primary distant metastasis of nasopharyngeal carcinoma to guide individualized application of FDG PET/CT. <i>European Journal of Nuclear Medicine and Molecular Imaging</i> , 2021, 48, 2586-2598. | 6.4 | 8 |
| 111 | Optimal cumulative cisplatin dose in nasopharyngeal carcinoma patients based on plasma Epstein-Barr virus DNA level after induction chemotherapy. <i>Aging</i> , 2020, 12, 4931-4944. | 3.1 | 8 |
| 112 | Reporting Quality of Randomized, Controlled Trials Evaluating Combined Chemoradiotherapy in Nasopharyngeal Carcinoma. <i>International Journal of Radiation Oncology Biology Physics</i> , 2017, 98, 170-176. | 0.8 | 7 |
| 113 | Association of MCP-1 promoter polymorphism with susceptibility to nasopharyngeal carcinoma. <i>Journal of Cellular Biochemistry</i> , 2019, 120, 6661-6670. | 2.6 | 7 |
| 114 | Optimizing the Treatment Pattern for De Novo Metastatic Nasopharyngeal Carcinoma Patients: A Large-Scale Retrospective Cohort Study. <i>Frontiers in Oncology</i> , 2020, 10, 543646. | 2.8 | 7 |
| 115 | Intensive Local Radiotherapy Is Associated With Better Local Control and Prolonged Survival in Bone-Metastatic Nasopharyngeal Carcinoma Patients. <i>Frontiers in Oncology</i> , 2020, 10, 378. | 2.8 | 7 |
| 116 | Construction of a comprehensive nutritional index and comparison of its prognostic performance with the PNI and NRI for survival in older patients with nasopharyngeal carcinoma: a retrospective study. <i>Supportive Care in Cancer</i> , 2021, 29, 5371-5381. | 2.2 | 7 |
| 117 | Impact of smoking on survival in nasopharyngeal carcinoma: A cohort study with 23,325 patients diagnosed from 1990 to 2016. <i>Radiotherapy and Oncology</i> , 2021, 162, 7-17. | 0.6 | 7 |
| 118 | Identification of surrogate endpoints in patients with locoregionally advanced nasopharyngeal carcinoma receiving neoadjuvant chemotherapy plus concurrent chemoradiotherapy versus concurrent chemoradiotherapy alone. <i>BMC Cancer</i> , 2015, 15, 930. | 2.6 | 6 |
| 119 | Nomogram Predicting the Benefits of Adding Concurrent Chemotherapy to Intensity-Modulated Radiotherapy After Induction Chemotherapy in Stages II-IVb Nasopharyngeal Carcinoma. <i>Frontiers in Oncology</i> , 2020, 10, 539321. | 2.8 | 6 |
| 120 | Optimal sequencing of chemotherapy with chemoradiotherapy based on TNM stage classification and EBV DNA in locoregionally advanced nasopharyngeal carcinoma. <i>Cancer Communications</i> , 2019, 39, 1-3. | 9.2 | 5 |
| 121 | Low value of whole-body dual-modality [18F]fluorodeoxyglucose positron emission tomography/computed tomography in primary staging of stage II nasopharyngeal carcinoma: a nest case-control study. <i>European Radiology</i> , 2021, 31, 5222-5233. | 4.5 | 5 |
| 122 | Increased Angiogenin Expression Correlates With Radiation Resistance and Predicts Poor Survival for Patients With Nasopharyngeal Carcinoma. <i>Frontiers in Pharmacology</i> , 2021, 12, 627935. | 3.5 | 5 |
| 123 | Establishment and validation of a prognostic nomogram to predict early metastasis in nasopharyngeal carcinoma patients within six months after radiotherapy and to guide intensive treatment. <i>Radiotherapy and Oncology</i> , 2021, 162, 202-211. | 0.6 | 5 |
| 124 | Efficacy of Transnasal Endoscopic Fine-Needle Aspiration Biopsy in Diagnosing Submucosal Nasopharyngeal Carcinoma. <i>Laryngoscope</i> , 2021, 131, 1798-1804. | 2.0 | 4 |
| 125 | Do all patients with locoregionally advanced nasopharyngeal carcinoma benefit from the maintenance chemotherapy using S-1/capecitabine?. <i>Oral Oncology</i> , 2021, 122, 105539. | 1.5 | 4 |
| 126 | Prognostic effect of pregnancy on young female patients with nasopharyngeal carcinoma: results from a matched cohort analysis. <i>Oncotarget</i> , 2016, 7, 21913-21921. | 1.8 | 4 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 127 | Percent change in apparent diffusion coefficient and plasma EBV DNA after induction chemotherapy identifies distinct prognostic response phenotypes in advanced nasopharyngeal carcinoma. <i>BMC Cancer</i> , 2021, 21, 1320. | 2.6 | 4 |
| 128 | Proteomic Analysis of a Nasopharyngeal Carcinoma Cell Line and a Nasopharyngeal Epithelial Cell Line. <i>Tumori</i> , 2015, 101, 676-683. | 1.1 | 3 |
| 129 | The prognosis of neck residue nasopharyngeal carcinoma (NPC) patients: results from a case-cohort study. <i>Journal of Cancer</i> , 2018, 9, 1765-1772. | 2.5 | 3 |
| 130 | Identifying optimal candidates for induction chemotherapy among stage IIa-IVa nasopharyngeal carcinoma based on pretreatment Epstein-Barr virus DNA and nodal maximal standard uptake values of [¹⁸ F]fluorodeoxyglucose positron emission tomography. <i>Cancer Medicine</i> , 2020, 9, 8852-8863. | 2.8 | 3 |
| 131 | Management of suboptimal response to induction chemotherapy in locoregionally advanced nasopharyngeal carcinoma: Re-induction therapy or direct to Radiotherapy?. <i>Radiotherapy and Oncology</i> , 2021, 163, 185-191. | 0.6 | 3 |
| 132 | Induction or adjuvant chemotherapy plus concurrent chemoradiotherapy versus concurrent chemoradiotherapy alone in paediatric nasopharyngeal carcinoma in the IMRT era: A recursive partitioning risk stratification analysis based on EBV DNA. <i>European Journal of Cancer</i> , 2021, 159, 133-143. | 2.8 | 3 |
| 133 | Establishment and validation of a recursive partitioning analysis based prognostic model for guiding re-radiotherapy in locally recurrent nasopharyngeal carcinoma patients. <i>Radiotherapy and Oncology</i> , 2022, 168, 61-68. | 0.6 | 3 |
| 134 | Association of Treatment Advances With Survival Rates in Pediatric Patients With Nasopharyngeal Carcinoma in China, 1989-2020. <i>JAMA Network Open</i> , 2022, 5, e220173. | 5.9 | 3 |
| 135 | Role of zoledronic acid in nasopharyngeal carcinoma patients with bone-only metastasis at diagnosis. <i>Oral Oncology</i> , 2019, 97, 31-36. | 1.5 | 2 |
| 136 | Development and validation of a normal tissue complication probability model for acquired nasal cavity stenosis and atresia after radical radiotherapy for nasopharyngeal carcinoma. <i>Radiotherapy and Oncology</i> , 2021, 160, 9-17. | 0.6 | 2 |
| 137 | Impact of salvage radiotherapy on survival of patients with advanced locally recurrent nasopharyngeal carcinoma: Derivation and validation of a predictive model. <i>Radiotherapy and Oncology</i> , 2022, 167, 252-260. | 0.6 | 2 |
| 138 | Cost-Effectiveness analysis of combining plasma Epstein-Barr virus DNA testing and different surveillance imaging modalities for nasopharyngeal carcinoma patients in first remission. <i>Oral Oncology</i> , 2022, 128, 105851. | 1.5 | 2 |
| 139 | Longitudinal Trend of Health-Related Quality of Life During Concurrent Chemoradiotherapy and Survival in Patients With Stage IIa-IVb Nasopharyngeal Carcinoma. <i>Frontiers in Oncology</i> , 2020, 10, 579292. | 2.8 | 1 |
| 140 | Management of first-line palliative chemotherapy for post-treatment metastasis after gemcitabine plus cisplatin induction chemotherapy: Gemcitabine plus cisplatin and non-gemcitabine plus cisplatin chemotherapy. <i>Head and Neck</i> , 2022, 44, 113-121. | 2.0 | 1 |
| 141 | Definitive radiation therapy and liver local therapy in de novo liver metastatic nasopharyngeal carcinoma: Large cohort study. <i>Head and Neck</i> , 2022, , . | 2.0 | 1 |
| 142 | Determining the suitability of definitive radiation therapy in patients with metastatic nasopharyngeal carcinoma based on PET/CT: a large cohort study. <i>European Radiology</i> , 2022, , 1. | 4.5 | 1 |
| 143 | Alpha-fetoprotein-producing recurrent nasopharyngeal carcinoma: A case report. <i>SAGE Open Medical Case Reports</i> , 2021, 9, 2050313X2110577. | 0.3 | 0 |
| 144 | Construction and validation of a biochemical signature to predict the prognosis and the benefit of induction chemotherapy in patients with nasopharyngeal carcinoma.. <i>American Journal of Cancer Research</i> , 2022, 12, 1635-1647. | 1.4 | 0 |