
Heidrun Ellinger-Ziegelbauer

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/7299032/publications.pdf Version: 2024-02-01



Heidrun

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 1 | Urinary miRNA Profiles in Chronic Kidney Injury—Benefits of Extracellular Vesicle Enrichment and miRNAs as Potential Biomarkers for Renal Fibrosis, Glomerular Injury, and Endothelial Dysfunction. Toxicological Sciences, 2022, , . | 3.1 | 2 |
| 2 | A Collaborative Initiative to Establish Genomic Biomarkers for Assessing Tumorigenic Potential to Reduce Reliance on Conventional Rodent Carcinogenicity Studies. Toxicological Sciences, 2022, 188, 4-16. | 3.1 | 7 |
| 3 | Methodological considerations for measuring biofluid-based microRNA biomarkers. Critical Reviews in Toxicology, 2021, 51, 264-282. | 3.9 | 13 |
| 4 | Urinary miRNA Biomarkers of Drug-Induced Kidney Injury and Their Site Specificity Within the Nephron. Toxicological Sciences, 2021, 180, 1-16. | 3.1 | 19 |
| 5 | A cross-sector call to improve carcinogenicity risk assessment through use of genomic methodologies. Regulatory Toxicology and Pharmacology, 2020, 110, 104526. | 2.7 | 21 |
| 6 | Energy metabolism modulation byÂbiguanides in comparison with rotenone in rat liver and heart. Archives of Toxicology, 2019, 93, 2603-2615. | 4.2 | 6 |
| 7 | Prediction of human drug-induced liver injury (DILI) in relation to oral doses and blood concentrations. Archives of Toxicology, 2019, 93, 1609-1637. | 4.2 | 86 |
| 8 | TGx-DDI, a Transcriptomic Biomarker for Genotoxicity Hazard Assessment of Pharmaceuticals and Environmental Chemicals. Frontiers in Big Data, 2019, 2, 36. | 2.9 | 15 |
| 9 | Toxicogenomics directory of rat hepatotoxicants in vivo and in cultivated hepatocytes. Archives of Toxicology, 2018, 92, 3517-3533. | 4.2 | 46 |
| 10 | Mechanistic Investigations of the Mitochondrial Complex I Inhibitor Rotenone in the Context of Pharmacological and Safety Evaluation. Scientific Reports, 2017, 7, 45465. | 3.3 | 196 |
| 11 | Xenobiotic CAR Activators Induce Dlk1-Dio3 Locus Noncoding RNA Expression in Mouse Liver. Toxicological Sciences, 2017, 158, 367-378. | 3.1 | 7 |
| 12 | Quantitative targeted bile acid profiling as new markers for DILI in a model of methapyrilene-induced liver injury in rats. Toxicology, 2017, 386, 1-10. | 4.2 | 22 |
| 13 | Development and validation of a high-throughput transcriptomic biomarker to address 21st century genetic toxicology needs. Proceedings of the National Academy of Sciences of the United States of America, 2017, 114, E10881-E10889. | 7.1 | 70 |
| 14 | Time-matched analysis of DNA adduct formation and early gene expression as predictive tool for renal carcinogenesis in methylazoxymethanol acetate treated Eker rats. Archives of Toxicology, 2017, 91, 3427-3438. | 4.2 | 8 |
| 15 | Non-Lethal Endotoxin Injection: A Rat Model of Hypercoagulability. PLoS ONE, 2017, 12, e0169976. | 2.5 | 28 |
| 16 | Absolute Measurement of Cardiac Injury-Induced microRNAs in Biofluids across Multiple Test Sites. Toxicological Sciences, 2016, 154, 115-125. | 3.1 | 9 |
| 17 | Beyond miR-122: Identification of MicroRNA Alterations in Blood During a Time Course of Hepatobiliary Injury and Biliary Hyperplasia in Rats. Toxicological Sciences, 2016, 150, 3-14. | 3.1 | 33 |
| 18 | Glomerulonephritis-induced changes in kidney gene expression in rats. Genomics Data, 2015, 6, 81-82. | 1.3 | 0 |

Heidrun

| # | Article | IF | CITATIONS |
|----|--|-----|-----------|
| 19 | Glomerulonephritis-Induced Changes in Urinary and Kidney MicroRNA Profiles in Rats. Toxicological Sciences, 2015, 145, 348-359. | 3.1 | 18 |
| 20 | Evaluation of Toxicogenomics Approaches for Assessing the Risk of Nongenotoxic Carcinogenicity in Rat Liver. PLoS ONE, 2014, 9, e97678. | 2.5 | 17 |
| 21 | Comparison of the Mesoscale Discovery and Luminex multiplex platforms for measurement of urinary biomarkers in a cisplatin rat kidney injury model. Journal of Pharmacological and Toxicological Methods, 2014, 69, 196-204. | 0.7 | 23 |
| 22 | Phenobarbital Induces Cell Cycle Transcriptional Responses in Mouse LiverÂHumanized for ConstitutiveÂAndrostane and Pregnane X Receptors. Toxicological Sciences, 2014, 139, 501-511. | 3.1 | 60 |
| 23 | Urinary microRNA profiling for identification of biomarkers after cisplatin-induced kidney injury. Toxicology, 2014, 324, 147-157. | 4.2 | 66 |
| 24 | Cross-Platform Toxicogenomics for the Prediction of Non-Genotoxic Hepatocarcinogenesis in Rat. PLoS ONE, 2014, 9, e97640. | 2.5 | 44 |
| 25 | Pharmacokinetics explain in vivo/in vitro discrepancies of carcinogen-induced gene expression alterations in rat liver and cultivated hepatocytes. Archives of Toxicology, 2013, 87, 337-345. | 4.2 | 49 |
| 26 | Comparison of hepatocarcinogen-induced gene expression profiles in conventional primary rat hepatocytes with in vivo rat liver. Archives of Toxicology, 2012, 86, 1399-1411. | 4.2 | 23 |
| 27 | Comparison of genotoxicant-modified transcriptomic responses in conventional and epigenetically stabilized primary rat hepatocytes with in vivo rat liver data. Archives of Toxicology, 2012, 86, 1703-1715. | 4.2 | 15 |
| 28 | Transcriptomic alterations induced by Ochratoxin A in rat and human renal proximal tubular in vitro models and comparison to a rat in vivo model. Archives of Toxicology, 2012, 86, 571-589. | 4.2 | 42 |
| 29 | Testosterone response of hepatic gene expression in female mice having acquired testosterone-unresponsive immunity to Plasmodium chabaudi malaria. Steroids, 2011, 76, 1204-1212. | 1.8 | 15 |
| 30 | The Role of Residual Gadolinium in the Induction of Nephrogenic Systemic Fibrosis-Like Skin Lesions in Rats. Investigative Radiology, 2011, 46, 48-56. | 6.2 | 39 |
| 31 | The enhanced value of combining conventional and "omics―analyses in early assessment of drug-induced hepatobiliary injury. Toxicology and Applied Pharmacology, 2011, 252, 97-111. | 2.8 | 58 |
| 32 | EU Framework 6 Project: Predictive Toxicology (PredTox)—overview and outcome. Toxicology and Applied Pharmacology, 2011, 252, 73-84. | 2.8 | 84 |
| 33 | An elastic network model to identify characteristic stress response genes. Computational Biology and Chemistry, 2010, 34, 193-202. | 2.3 | 1 |
| 34 | Performance of Novel Kidney Biomarkers in Preclinical Toxicity Studies. Toxicological Sciences, 2010, 116, 8-22. | 3.1 | 101 |
| 35 | Characterization and Interlaboratory Comparison of a Gene Expression Signature for Differentiating Genotoxic Mechanisms. Toxicological Sciences, 2009, 110, 341-352. | 3.1 | 72 |
| 36 | Pulmonary toxicity of multi-walled carbon nanotubes (Baytubes®) relative to α-quartz following a single 6h inhalation exposure of rats and a 3 months post-exposure period. Toxicology, 2009, 266, 16-29. | 4.2 | 81 |

Heidrun

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 37 | Inter-laboratory comparison of human renal proximal tubule (HK-2) transcriptome alterations due to Cyclosporine A exposure and medium exhaustion. Toxicology in Vitro, 2009, 23, 486-499. | 2.4 | 36 |
| 38 | Application of toxicogenomics to study mechanisms of genotoxicity and carcinogenicity. Toxicology Letters, 2009, 186, 36-44. | 0.8 | 126 |
| 39 | Molecular Characterization of Preneoplastic Lesions Provides Insight on the Development of Renal Tumors. American Journal of Pathology, 2009, 175, 1686-1698. | 3.8 | 19 |
| 40 | Prediction of a carcinogenic potential of rat hepatocarcinogens using toxicogenomics analysis of short-term in vivo studies. Mutation Research - Fundamental and Molecular Mechanisms of Mutagenesis, 2008, 637, 23-39. | 1.0 | 214 |
| 41 | The carcinoGENOMICS project: Critical selection of model compounds for the development of omics-based in vitro carcinogenicity screening assays. Mutation Research - Reviews in Mutation Research, 2008, 659, 202-210. | 5.5 | 60 |
| 42 | Carcinogen-Specific Gene Expression Profiles in Short-term Treated Eker and Wild-type Rats Indicative of Pathways Involved in Renal Tumorigenesis. Cancer Research, 2007, 67, 4052-4068. | 0.9 | 56 |
| 43 | Establishment of a protocol for the gene expression analysis of laser microdissected rat kidney samples with affymetrix genechips. Toxicology and Applied Pharmacology, 2006, 217, 134-142. | 2.8 | 17 |
| 44 | Comparison of the expression profiles induced by genotoxic and nongenotoxic carcinogens in rat liver. Mutation Research - Fundamental and Molecular Mechanisms of Mutagenesis, 2005, 575, 61-84. | 1.0 | 197 |
| 45 | Characteristic Expression Profiles Induced by Genotoxic Carcinogens in Rat Liver. Toxicological Sciences, 2004, 77, 19-34. | 3.1 | 155 |
| 46 | Ste20-like kinase (SLK), a regulatory kinase for polo-like kinase (Plk) during the G2/M transition in somatic cells. Genes To Cells, 2000, 5, 491-498. | 1.2 | 76 |
| 47 | Physical and Functional Interaction of Filamin (Actin-binding Protein-280) and Tumor Necrosis Factor Receptor-associated Factor 2. Journal of Biological Chemistry, 2000, 275, 271-278. | 3.4 | 112 |
| 48 | Cell Cycle Arrest and Reversion of Ras-Induced Transformation by a Conditionally Activated Form of Mitogen-Activated Protein Kinase Kinase Kinase 3. Molecular and Cellular Biology, 1999, 19, 3857-3868. | 2.3 | 84 |
| 49 | Direct Activation of the Stress-activated Protein Kinase (SAPK) and Extracellular Signal-regulated Protein Kinase (ERK) Pathways by an Inducible Mitogen-activated Protein Kinase/ERK Kinase Kinase 3 (MEKK) Derivative. Journal of Biological Chemistry, 1997, 272, 2668-2674. | 3.4 | 91 |