

# Peter W Sorensen

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7334393/publications.pdf>

Version: 2024-02-01

98  
papers

5,757  
citations

71102

41  
h-index

82547

72  
g-index

115  
all docs

115  
docs citations

115  
times ranked

3890  
citing authors

#	ARTICLE	IF	CITATIONS
1	Migration, homing and spatial ecology of common carp in interconnected lakes. <i>Ecology of Freshwater Fish</i> , 2022, 31, 164-176.	1.4	13
2	Numeric Simulation Demonstrates That the Upstream Movement of Invasive Bigheaded Carp Can Be Blocked at Sets of Mississippi River Locks-and-Dams Using a Combination of Optimized Spillway Gate Operations, Lock Deterrents, and Carp Removal. <i>Fishes</i> , 2021, 6, 10.	1.7	9
3	Hormonal Prostaglandin F <sub>2</sub> ± Mediates Behavioral Responsiveness to a Species-Specific Multi-component Male Hormonal Sex Pheromone in a Female Fish. <i>Integrative and Comparative Biology</i> , 2021, 61, 193-204.	2.0	5
4	Global Inland Capture and Culture Finfisheries Follow Different Trends When Evaluated by the Human Development Index. <i>Sustainability</i> , 2021, 13, 8420.	3.2	2
5	Introduction to the Biology and Control of Invasive Fishes and a Special Issue on This Topic. <i>Fishes</i> , 2021, 6, 69.	1.7	4
6	Monitoring upstream fish passage through a Mississippi River lock and dam reveals species differences in lock chamber usage and supports a fish passage model which describes velocity-dependent passage through spillway gates. <i>River Research and Applications</i> , 2020, 36, 36-46.	1.7	14
7	Common Carp Are Initially Repelled by a Broadband Outboard Motor Sound in a Lock Chamber but Habituate Rapidly. <i>North American Journal of Fisheries Management</i> , 2020, 40, 1499-1509.	1.0	6
8	Otolith Microchemistry of Common Carp Reflects Capture Location and Differentiates Nurseries in an Interconnected Lake System of the North American Midwest. <i>North American Journal of Fisheries Management</i> , 2020, 40, 1100-1118.	1.0	2
9	Case Studies Demonstrate That Common Carp Can Be Sustainably Reduced by Exploiting Source-Sink Dynamics in Midwestern Lakes. <i>Fishes</i> , 2020, 5, 36.	1.7	14
10	A complex sound coupled with an air curtain blocks invasive carp passage without habituation in a laboratory flume. <i>Biological Invasions</i> , 2019, 21, 2837-2855.	2.4	21
11	Nonlinear relationship between Silver Carp density and their eDNA concentration in a large river. <i>PLoS ONE</i> , 2019, 14, e0218823.	2.5	41
12	The Effect of Modifying a CFD-AB Approach on Fish Passage through a Model Hydraulic Dam. <i>Water (Switzerland)</i> , 2019, 11, 1776.	2.7	6
13	A Blend of F Prostaglandins Functions as an Attractive Sex Pheromone in Silver Carp. <i>Fishes</i> , 2019, 4, 27.	1.7	9
14	The Chemical Sensitivity and Electrical Activity of Individual Olfactory Sensory Neurons to a Range of Sex Pheromones and Food Odors in the Goldfish. <i>Chemical Senses</i> , 2018, 43, 249-260.	2.0	19
15	Source-sink dynamics explain the distribution and persistence of an invasive population of common carp across a model Midwestern watershed. <i>Biological Invasions</i> , 2018, 20, 1961-1976.	2.4	25
16	High levels of circulating prostaglandin F <sub>2</sub> ± associated with ovulation stimulate female sexual receptivity and spawning behavior in the goldfish ( <i>Carassius auratus</i> ). <i>General and Comparative Endocrinology</i> , 2018, 267, 128-136.	1.8	22
17	Attracting Common Carp to a bait site with food reveals strong positive relationships between fish density, feeding activity, environmental <sc>DNA</sc>, and sex pheromone release that could be used in invasive fish management. <i>Ecology and Evolution</i> , 2018, 8, 6714-6727.	1.9	34
18	Chemical Cues which Include Amino Acids Mediate Species-Specific Feeding Behavior in Invasive Filter-Feeding Bigheaded Carps. <i>Journal of Chemical Ecology</i> , 2017, 43, 374-384.	1.8	10

#	ARTICLE	IF	CITATIONS
19	Silver, bighead, and common carp orient to acoustic particle motion when avoiding a complex sound. PLoS ONE, 2017, 12, e0180110.	2.5	20
20	Invasive Bighead and Silver Carps Form Different Sized Shoals that Readily Intermix. PLoS ONE, 2016, 11, e0157174.	2.5	14
21	Biological invasion by a benthivorous fish reduced the cover and species richness of aquatic plants in most lakes of a large North American ecoregion. Global Change Biology, 2016, 22, 3937-3947.	9.5	47
22	Male-typical courtship, spawning behavior, and olfactory sensitivity are induced to different extents by androgens in the goldfish suggesting they are controlled by different neuroendocrine mechanisms. General and Comparative Endocrinology, 2016, 232, 160-173.	1.8	28
23	Different Migratory Strategies of Invasive Common Carp and Native Northern Pike in the American Midwest Suggest an Opportunity for Selective Management Strategies. North American Journal of Fisheries Management, 2016, 36, 769-779.	1.0	19
24	Theory and Application of Semiochemicals in Nuisance Fish Control. Journal of Chemical Ecology, 2016, 42, 698-715.	1.8	31
25	Environment shapes the fecal microbiome of invasive carp species. Microbiome, 2016, 4, 44.	11.1	166
26	Effects of Temperature and Trophic State on Degradation of Environmental DNA in Lake Water. Environmental Science & Technology, 2016, 50, 1859-1867.	10.0	210
27	Effects of common carp ( <i>Cyprinus carpio</i> ) on sediment mixing depth and mobile phosphorus mass in the active sediment layer of a shallow lake. Hydrobiologia, 2016, 763, 23-33.	2.0	48
28	Optimizing techniques to capture and extract environmental <sc>DNA</sc> for detection and quantification of fish. Molecular Ecology Resources, 2016, 16, 56-68.	4.8	171
29	Partial migration to seasonallyâ€unstable habitat facilitates biological invasions in a predatorâ€dominated system. Oikos, 2015, 124, 1520-1526.	2.7	19
30	Effects of common carp on phosphorus concentrations, water clarity, and vegetation density: a whole system experiment in a thermally stratified lake. Hydrobiologia, 2015, 746, 303-311.	2.0	42
31	Lamprey Spawning Migration. , 2015, , 215-263.		67
32	The Relationship between the Distribution of Common Carp and Their Environmental DNA in a Small Lake. PLoS ONE, 2014, 9, e112611.	2.5	172
33	Anatomical and physiological studies of bigheaded carps demonstrate that the epibranchial organ functions as a pharyngeal taste organ. Journal of Experimental Biology, 2014, 217, 3945-54.	1.7	19
34	<i>Guidelines for Use of Fishes in Research</i>â€Revised and Expanded, 2014. Fisheries, 2014, 39, 415-416.	0.8	35
35	Chemical Analysis of Aquatic Pheromones in Fish. Methods in Molecular Biology, 2013, 1068, 55-69.	0.9	3
36	Direct Field and Laboratory Evidence that a Combination of Egg and Larval Predation Controls Recruitment of Invasive Common Carp in Many Lakes of the Upper Mississippi River Basin. Transactions of the American Fisheries Society, 2013, 142, 1134-1140.	1.4	22

#	ARTICLE	IF	CITATIONS
37	Behavioral Analysis of Pheromones in Fish. <i>Methods in Molecular Biology</i> , 2013, 1068, 293-305.	0.9	1
38	Variation in native micro-predator abundance explains recruitment of a mobile invasive fish, the common carp, in a naturally unstable environment. <i>Biological Invasions</i> , 2012, 14, 1919-1929.	2.4	49
39	Using Boat Electrofishing to Estimate the Abundance of Invasive Common Carp in Small Midwestern Lakes. <i>North American Journal of Fisheries Management</i> , 2012, 32, 817-822.	1.0	36
40	Common Carp Implanted with Prostaglandin F <sub>2</sub> ± Release a Sex Pheromone Complex that Attracts Conspecific Males in Both the Laboratory and Field. <i>Journal of Chemical Ecology</i> , 2012, 38, 127-134.	1.8	22
41	Olfactory-mediated stream-finding behavior of migratory adult sea lamprey ( <i>Petromyzon marinus</i> ). <i>Canadian Journal of Fisheries and Aquatic Sciences</i> , 2011, 68, 523-533.	1.4	54
42	Behavioral and genomic impacts of a wastewater effluent on the fathead minnow. <i>Aquatic Toxicology</i> , 2011, 101, 38-48.	4.0	80
43	Behavioral responses of adult male and female fathead minnows to a model estrogenic effluent and its effects on exposure regime and reproductive success. <i>Aquatic Toxicology</i> , 2011, 101, 521-528.	4.0	14
44	Synthesis and olfactory activity of unnatural, sulfated 5 <sup>β</sup> -bile acid derivatives in the sea lamprey ( <i>Petromyzon marinus</i> ). <i>Steroids</i> , 2011, 76, 291-300.	1.8	18
45	A Multi-Component Species Identifying Pheromone in the Goldfish. <i>Journal of Chemical Ecology</i> , 2011, 37, 219-227.	1.8	39
46	Polar Metabolites Synergize the Activity of Prostaglandin F <sub>2</sub> ± in a Species-Specific Hormonal Sex Pheromone Released by Ovulated Common Carp. <i>Journal of Chemical Ecology</i> , 2011, 37, 695-704.	1.8	40
47	Pheromones in Vertebrates. , 2010, , 225-262.		15
48	Bold minnows consistently approach danger in the field and lab in response to either chemical or visual indicators of predation risk. <i>Behavioral Ecology and Sociobiology</i> , 2010, 64, 381-387.	1.4	23
49	Recruitment and abundance of an invasive fish, the common carp, is driven by its propensity to invade and reproduce in basins that experience winter-time hypoxia in interconnected lakes. <i>Biological Invasions</i> , 2010, 12, 1101-1112.	2.4	106
50	Cognitive aspects of food searching behavior in free-ranging wild Common Carp. <i>Environmental Biology of Fishes</i> , 2010, 88, 295-300.	1.0	42
51	Production and fate of the sea lamprey migratory pheromone. <i>Fish Physiology and Biochemistry</i> , 2010, 36, 1013-1020.	2.3	21
52	Importance of the olfactory sense to migratory sea lampreys <i>Petromyzon marinus</i> seeking riverine spawning habitat. <i>Journal of Fish Biology</i> , 2010, 76, 949-964.	1.6	52
53	Olfactory Sensitivity of Pacific Lampreys to Lamprey Bile Acids. <i>Transactions of the American Fisheries Society</i> , 2009, 138, 144-152.	1.4	34
54	Effects of a rapidly increasing population of common carp on vegetative cover and waterfowl in a recently restored Midwestern shallow lake. <i>Hydrobiologia</i> , 2009, 632, 235-245.	2.0	115

#	ARTICLE	IF	CITATIONS
55	A Sterol-Like Odorant in the Urine of Mozambique Tilapia Males Likely Signals Social Dominance to Females. <i>Journal of Chemical Ecology</i> , 2008, 34, 438-449.	1.8	68
56	Neural Processing, Perception, and Behavioral Responses to Natural Chemical Stimuli by Fish and Crustaceans. <i>Journal of Chemical Ecology</i> , 2008, 34, 898-914.	1.8	159
57	Isolation and Biological Activity of the Multi-Component Sea Lamprey Migratory Pheromone. <i>Journal of Chemical Ecology</i> , 2008, 34, 1259-1267.	1.8	57
58	High-Potency Olfactory Receptor Agonists Discovered by Virtual High-Throughput Screening: Molecular Probes for Receptor Structure and Olfactory Function. <i>Neuron</i> , 2008, 60, 767-774.	8.1	26
59	Temporal Variation in the Estrogenicity of a Sewage Treatment Plant Effluent and Its Biological Significance. <i>Environmental Science &amp; Technology</i> , 2008, 42, 3421-3427.	10.0	54
60	Common carp and goldfish discern conspecific identity using chemical cues. <i>Behaviour</i> , 2008, 145, 1409-1425.	0.8	29
61	Details of the Structure Determination of the Sulfated Steroids PSDS and PADS:Â New Components of the Sea Lamprey ( <i>Petromyzonmarinus</i> ) Migratory Pheromone. <i>Journal of Organic Chemistry</i> , 2007, 72, 7544-7550.	3.2	41
62	A critical review of the discovery and application of a migratory pheromone in an invasive fish, the sea lamprey <i>Petromyzon marinus</i> L.. <i>Journal of Fish Biology</i> , 2007, 71, 100-114.	1.6	51
63	Female goldfish signal spawning readiness by altering when and where they release a urinary pheromone. <i>Animal Behaviour</i> , 2007, 74, 1329-1338.	1.9	69
64	ENVIRONMENTAL ESTROGENS SUPPRESS HORMONES, BEHAVIOR, AND REPRODUCTIVE FITNESS IN MALE FATHEAD MINNOWS. <i>Environmental Toxicology and Chemistry</i> , 2007, 26, 271.	4.3	118
65	A field test verifies that pheromones can be useful for sea lamprey ( <i>Petromyzon marinus</i> ) control in the Great Lakes. <i>Canadian Journal of Fisheries and Aquatic Sciences</i> , 2006, 63, 475-479.	1.4	76
66	A Practical Method for Obtaining Useful Quantities of Pheromones from Sea Lamprey and Other Fishes for Identification and Control. <i>Journal of Great Lakes Research</i> , 2006, 32, 832.	1.9	22
67	Mixture of new sulfated steroids functions as a migratory pheromone in the sea lamprey. <i>Nature Chemical Biology</i> , 2005, 1, 324-328.	8.0	222
68	Sexually mature male goldfish release large quantities of androstenedione into the water where it functions as a pheromone. <i>General and Comparative Endocrinology</i> , 2005, 140, 164-175.	1.8	121
69	Biologically Relevant Concentrations of Petromyzonol Sulfate, a Component of the Sea Lamprey Migratory Pheromone, Measured in Stream Water. <i>Journal of Chemical Ecology</i> , 2005, 31, 2205-2210.	1.8	36
70	Stable Isotope Analysis of Amphidromous Hawaiian Gobies Suggests Their Larvae Spend a Substantial Period of Time in Freshwater River Plumes. <i>Environmental Biology of Fishes</i> , 2005, 74, 31-42.	1.0	42
71	Second Messenger Systems Mediating Sex Pheromone and Amino Acid Sensitivity in Goldfish Olfactory Receptor Neurons. <i>Chemical Senses</i> , 2005, 30, i315-i316.	2.0	14
72	Reproductive Pheromones. <i>Fish Physiology</i> , 2005, , 359-412.	0.8	47

#	ARTICLE	IF	CITATIONS
73	Evidence that 4-pregnen-17,20 <sup>12</sup> ,21-triol-3-one functions as a maturation-inducing hormone and pheromonal precursor in the percid fish, <i>Gymnocephalus cernuus</i> . <i>General and Comparative Endocrinology</i> , 2004, 139, 1-11.	1.8	26
74	Evidence That Petromyzontid Lampreys Employ a Common Migratory Pheromone That Is Partially Comprised of Bile Acids. <i>Journal of Chemical Ecology</i> , 2004, 30, 2091-2110.	1.8	91
75	Brief review of fish pheromones and discussion of their possible uses in the control of non <sup>1</sup> indigenous teleost fishes. <i>New Zealand Journal of Marine and Freshwater Research</i> , 2004, 38, 399-417.	2.0	141
76	The Chemical Ecology and Potential Application of the Sea Lamprey Migratory Pheromone. <i>Journal of Great Lakes Research</i> , 2003, 29, 66-84.	1.9	88
77	Possible Applications of Pheromones in an Integrated Sea Lamprey Management Program. <i>Journal of Great Lakes Research</i> , 2003, 29, 794-800.	1.9	41
78	Spawning Interactions between Sympatric Brown and Brook Trout May Contribute to Species Replacement. <i>Transactions of the American Fisheries Society</i> , 2002, 131, 569-576.	1.4	19
79	Hormonal Pheromones in Fish. , 2002, , 375-434.		29
80	Ten <sup>1</sup> week exposure to treated sewage discharge has relatively minor, variable effects on reproductive behavior and sperm production in goldfish. <i>Environmental Toxicology and Chemistry</i> , 2002, 21, 2185-2190.	4.3	51
81	Hormonal and pheromonal control of spawning behavior in the goldfish. <i>Fish Physiology and Biochemistry</i> , 2002, 26, 71-84.	2.3	196
82	Laboratory assessment of the role of a larval pheromone and natural stream odor in spawning stream localization by migratory sea lamprey ( <i>Petromyzon marinus</i> ). <i>Canadian Journal of Fisheries and Aquatic Sciences</i> , 2001, 58, 2374-2385.	1.4	99
83	The three steroidal components of the goldfish preovulatory pheromone signal evoke different behaviors in males. <i>Comparative Biochemistry and Physiology - B Biochemistry and Molecular Biology</i> , 2001, 129, 645-651.	1.6	96
84	Title is missing!. <i>Fish Physiology and Biochemistry</i> , 2001, 24, 15-30.	2.3	77
85	Injured Eurasian Ruffe, <i>Gymnocephalus cernuus</i> , Release an Alarm Pheromone that Could be Used to Control their Dispersal. <i>Journal of Great Lakes Research</i> , 2000, 26, 183-195.	1.9	26
86	Direct behavioral evidence that unique bile acids released by larval sea lamprey ( <i>Petromyzon</i> ) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 2. 2000, 57, 557-569.	1.4	145
87	Functional Identification of a Goldfish Odorant Receptor. <i>Neuron</i> , 1999, 23, 487-498.	8.1	224
88	Evolution and Specialization of Fish Hormonal Pheromones. , 1999, , 15-47.		60
89	Discrimination of pheromonal cues in fish: emerging parallels with insects. <i>Current Opinion in Neurobiology</i> , 1998, 8, 458-467.	4.2	75
90	High rate of redd superimposition by brook trout ( <i>Salvelinus fontinalis</i> ) and brown trout ( <i>Salmo trutta</i> ) in a Minnesota stream cannot be explained by habitat availability alone. <i>Canadian Journal of Fisheries and Aquatic Sciences</i> , 1998, 55, 2310-2316.	1.4	97

#	ARTICLE	IF	CITATIONS
91	Management Briefs: Effects of Electroshocking on the Sexual Behavior of Goldfish and Brook Trout. North American Journal of Fisheries Management, 1994, 14, 862-865.	1.0	5
92	Exposure to the pheromone 17 $\beta$ ,20 $\alpha$ -dihydroxy-4-pregnen-3-one enhances the behavioural spawning success, sperm production and sperm motility of male goldfish. Animal Behaviour, 1993, 46, 245-256.	1.9	93
93	Sex pheromones selectively stimulate the medial olfactory tracts of male goldfish. Brain Research, 1991, 558, 343-347.	2.2	77
94	Differing behavioral and endocrinological effects of two female sex pheromones on male goldfish. Hormones and Behavior, 1989, 23, 317-332.	2.1	99
95	F Prostaglandins Function as Potent Olfactory Stimulants that Comprise the Postovulatory Female Sex Pheromone in Goldfish. Biology of Reproduction, 1988, 39, 1039-1050.	2.7	285
96	Extreme olfactory sensitivity of mature and gonadally-regressed goldfish to a potent steroidal pheromone, 17 $\beta$ ,20 $\alpha$ -dihydroxy-4-pregnen-3-one. Journal of Comparative Physiology A: Neuroethology, Sensory, Neural, and Behavioral Physiology, 1987, 160, 305-313.	1.6	107
97	Origins of the freshwater attractant(s) of migrating elvers of the American eel, <i>Anguilla rostrata</i> . Environmental Biology of Fishes, 1986, 17, 185-200.	1.0	67
98	Movements of a model fish, the common carp, through a generic Mississippi River lock and dam demonstrate how fish swimming performance, behavior, and discharge-driven flow fields determine fish passage rates in ways that can be predicted and modified using fish passage models. River Research and Applications, 0, , .	1.7	4