## Maya Evenden

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/7340396/publications.pdf

Version: 2024-02-01

361413 454955 1,413 103 20 30 citations h-index g-index papers 103 103 103 1089 docs citations times ranked citing authors all docs

#	Article	IF	CITATIONS
1	OUP accepted manuscript. Environmental Entomology, 2022, , .	1.4	О
2	Effect of semiochemical exposure on flight propensity and flight capacity of Dendroctonus ponderosae in laboratory bioassays. Arthropod-Plant Interactions, 2021, 15, 551-562.	1.1	1
3	Management of Pea Leaf Weevil (Coleoptera: Curculionidae) and Development of a Nominal Threshold in Faba Beans. Journal of Economic Entomology, 2021, 114, 1597-1606.	1.8	2
4	Host Plant Volatile Lures Attract Apanteles polychrosidis (Hymenoptera: Braconidae) to Ash Trees Infested With Caloptilia fraxinella (Lepidoptera: Gracillariidae). Frontiers in Ecology and Evolution, 2021, 9, .	2.2	1
5	Seasonal emergence patterns of Sitodiplosis mosellana (Diptera: Cecidomyiidae) in the Peace River region, Alberta, Canada. Canadian Entomologist, 2021, 153, 222-236.	0.8	1
6	Identification of genes and gene expression associated with dispersal capacity in the mountain pine beetle, <i>Dendroctonus ponderosae</i> Hopkins (Coleoptera: Curculionidae). Peerl, 2021, 9, e12382.	2.0	1
7	Influence of Host Plant Species and Fertilization Regime on Larval Performance and Feeding Preference of the Redbacked Cutworm and the Pale Western Cutworm (Lepidoptera: Noctuidae). Journal of Economic Entomology, 2020, 113, 731-741.	1.8	3
8	Effect of Environmental Conditions on Flight Capacity in Mountain Pine Beetle (Coleoptera:) Tj ETQq0 0 0 rgBT /	/Overlock	10 Tf 50 462 T
9	Infection of canola by the root pathogen Plasmodiophora brassicae increases resistance to aboveground herbivory by bertha armyworm, Mamestra configurata Walker (Lepidoptera: Noctuidae). Plant Science, 2020, 300, 110625.	3.6	6
10	Bumble Bees (Hymenoptera: Apidae) Respond to Moth (Lepidoptera: Noctuidae) Pheromone Components, Leading to Bee Bycatch in Monitoring Traps Targeting Moth Pests. Frontiers in Ecology and Evolution, 2020, 8, .	2.2	10
11	Local and Landscape-Scale Features Influence Bumble Bee (Hymenoptera: Apidae) Bycatch in Bertha Armyworm Mamestra configurata (Lepidoptera: Noctuidae) Pheromone-Baited Monitoring Traps. Environmental Entomology, 2020, 49, 1127-1136.	1.4	6
12	Patterns of Diversity in the Symbiotic Mite Assemblage of the Mountain Pine Beetle, Dendroctonus Ponderosae Hopkins. Forests, 2020, $11$ , $1102$ .	2.1	4
13	Mechanisms and consequences of flight polyphenisms in an outbreaking bark beetle species. Journal of Experimental Biology, 2020, 223, .	1.7	4
14	Assessment of Available Tools for Monitoring Wheat Midge (Diptera: Cecidomyiidae). Environmental Entomology, 2020, 49, 627-637.	1.4	6
15	Modeling the dispersal–reproduction trade-off in an expanding population. Theoretical Population Biology, 2020, 134, 147-159.	1.1	4
16	Bugs 101: Insect–Human Interactions; Developing and Implementing a General Entomology MOOC (Massive Open Online Course). American Entomologist, 2020, 66, 55-60.	0.2	0
17	Influence of crop variety and fertilization on oviposition preference and larval performance of a generalist herbivore, the true armyworm, Mythimna unipuncta. Entomologia Experimentalis Et Applicata, 2020, 168, 266-278.	1.4	2
18	Factors influencing dispersal by flight in bark beetles (Coleoptera: Curculionidae: Scolytinae): from genes to landscapes. Canadian Journal of Forest Research, 2019, 49, 1024-1041.	1.7	31

#	Article	IF	Citations
19	Energy use by the mountain pine beetle (Coleoptera: Curculionidae: Scolytinae) for dispersal by flight. Physiological Entomology, 2019, 44, 200-208.	1.5	9
20	Testing for trade-offs between flight and reproduction in the mountain pine beetle (Coleoptera:) Tj ETQq0 0 0 r	gBT/Qver	lock <sub>5</sub> 10 Tf 50 7
21	Herbivoreâ€induced plants do not affect oviposition but do affect fitness of subsequent herbivores on canola. Entomologia Experimentalis Et Applicata, 2019, 167, 341-349.	1.4	6
22	Efficacy of Chlorantraniliprole Seed Treatments Against Armyworm ( <i>Mythimna unipuncta</i> ) Tj ETQq0 0 0 188-195.	rgBT /Over 1.8	rlock 10 Tf 50 15
23	Morphological variation associated with dispersal capacity in a treeâ€killing bark beetle <i>Dendroctonus ponderosae</i> Hopkins. Agricultural and Forest Entomology, 2019, 21, 79-87.	1.3	12
24	Wing polymorphisms of Pterostichus melanarius (Coleoptera: Carabidae) (Illiger, 1978) in Alberta pulse crops. Alberta Academic Review, 2019, 2, 23-24.	0.0	1
25	State-Dependent Plasticity in Response to Host-Plant Volatiles in a Long-Lived Moth, Caloptilia fraxinella (Lepidoptera: Gracillariidae). Journal of Chemical Ecology, 2018, 44, 276-287.	1.8	7
26	Progress Toward Integrated Pest Management of Pea Leaf Weevil: A Review. Annals of the Entomological Society of America, 2018, 111, 144-153.	2.5	22
27	Evaluation of Semiochemical-Baited Traps for Monitoring the Pea Leaf Weevil, Sitona lineatus (Coleoptera: Curculionidae) in Field Pea Crops. Environmental Entomology, 2018, 47, 93-106.	1.4	13
28	Canola Nutrition and Variety Affect Oviposition and Offspring Performance in the Generalist Herbivore, Mamestra configurata (Lepidoptera: Noctuidae). Journal of Economic Entomology, 2018, 111, 1702-1710.	1.8	8
29	Semiochemical-Based Management of the Pea Leaf Weevil (Coleoptera: Curculionidae). Annals of the Entomological Society of America, 2018, 111, 154-160.	2.5	10
30	Olfactory hostâ€finding behaviour of <i>Oulema melanopus</i> (Coleoptera: Chrysomelidae) and its parasitoid, <i>Tetrastichus julis</i> (Hymenoptera: Eulophidae). Journal of Applied Entomology, 2017, 141, 740-750.	1.8	3
31	The effect of cold storage of mass-reared codling moths (Lepidoptera: Tortricidae) on subsequent flight capacity. Canadian Entomologist, 2017, 149, 391-398.	0.8	10
32	Water-deficit and fungal infection can differentially affect the production of different classes of defense compounds in two host pines of mountain pine beetle. Tree Physiology, 2017, 37, 338-350.	3.1	35
33	The influence of Canadian research on semiochemical-based management of forest insect pests in Canada—ERRATUM. Canadian Entomologist, 2017, 149, 139-139.	0.8	0
34	The Effect of Water Limitation on Volatile Emission, Tree Defense Response, and Brood Success of Dendroctonus ponderosae in Two Pine Hosts, Lodgepole, and Jack Pine. Frontiers in Ecology and Evolution, 2016, 4, .	2.2	26
35	The roles of juvenile hormone and biogenic amines on pheromone response plasticity and diapause termination in male <i>Caloptilia fraxinella</i> . Entomologia Experimentalis Et Applicata, 2016, 158, 184-201.	1.4	3
36	Host plant preference and offspring performance of a leafâ€mining moth, <i><scp>C</scp>aloptilia fraxinella</i> , on two <i><scp>F</scp>raxinus</i> species. Entomologia Experimentalis Et Applicata, 2016, 159, 311-326.	1.4	5

#	Article	IF	Citations
37	Potential for semiochemical-based monitoring of the pea leaf weevil (Coleoptera: Curculionidae) on field pea (Fabaceae) in the Canadian Prairie Provinces. Canadian Entomologist, 2016, 148, 595-602.	0.8	12
38	The influence of Canadian research on semiochemical-based management of forest insect pests in Canada. Canadian Entomologist, 2016, 148, S170-S209.	0.8	9
39	Genetic diversity and population structure identify the potential source of the invasive red clover casebearer moth, Coleophora deauratella, in North America. Biological Invasions, 2016, 18, 3595-3609.	2.4	11
40	Size and protein content of accessory glands in adult male <i>Caloptilia fraxinella</i> in different physiological states. Physiological Entomology, 2016, 41, 74-82.	1.5	5
41	Aggregation and a strong <scp>A</scp> llee effect in a cooperative outbreak insect. Ecological Applications, 2016, 26, 2623-2636.	3.8	25
42	Nutrition, sex and season contribute to variation in fat and glycerol levels in the longâ€lived moth <i>Caloptilia fraxinella</i> . Physiological Entomology, 2016, 41, 67-73.	1.5	0
43	Mating disruption of <i>Coleophora deauratella </i> (Lepidoptera: Coleophoridae) using laminate flakes in red clover seed production fields. Pest Management Science, 2015, 71, 1149-1157.	3.4	3
44	Forest tent caterpillar, Malacosoma disstria (Lepidoptera: Lasiocampidae), mate-finding behavior is greatest at intermediate population densities: implications for interpretation of moth capture in pheromone-baited traps. Frontiers in Ecology and Evolution, 2015, 3, .	2.2	9
45	Differential parasitism by a generalist parasitoid is mediated by volatile organic chemicals of the herbivore's host. Arthropod-Plant Interactions, 2015, 9, 515-527.	1.1	6
46	Resource Allocation to Flight in an Outbreaking Forest Defoliator Malacosoma disstria. Environmental Entomology, 2015, 44, 835-845.	1.4	13
47	Challenges of Mating Disruption Using Aerosol-Emitting Pheromone Puffers in Red Clover Seed Production Fields to Control Coleophora deauratella (Lepidoptera: Coleophoridae). Environmental Entomology, 2015, 44, 34-43.	1.4	10
48	Diversity and abundance of arthropod by-catch in semiochemical-baited traps targeting apple clearwing moth (Lepidoptera: Sesiidae) in organic and conventional apple orchards in British Columbia, Canada. Canadian Entomologist, 2015, 147, 227-243.	0.8	12
49	The costs of colour: plasticity of melanin pigmentation in an outbreaking polymorphic forest moth. Entomologia Experimentalis Et Applicata, 2015, 154, 242-250.	1.4	10
50	Environmental conditions terminate reproductive diapause and influence pheromone perception in the longâ€lived moth <i><scp>C</scp>aloptilia fraxinella</i> . Physiological Entomology, 2015, 40, 30-42.	1.5	12
51	The use of plant volatiles for host location by an ash (Fraxinus) specialist, Caloptilia fraxinella. Chemoecology, 2014, 24, 229-242.	1.1	7
52	Factors Influencing Male <l>Plutella xylostella</l> (Lepidoptera: Plutellidae) Capture Rates in Sex Pheromone-Baited Traps on Canola in Western Canada. Journal of Economic Entomology, 2014, 107, 2067-2076.	1.8	9
53	Relationships among male <i>Coleophora deauratella</i> ( <scp>L</scp> epidoptera:) Tj ETQq1 1 0.784314 rgBT phenology. Agricultural and Forest Entomology, 2014, 16, 207-215.	Γ/Overlock 1.3	2 10 Tf 50 107 10

Factors Influencing Flight Capacity of the Mountain Pine Beetle (Coleoptera: Curculionidae:) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 62 Td 1.4

#	Article	IF	Citations
55	Chemical similarity between historical and novel host plants promotes range and host expansion of the mountain pine beetle in a naÃ-ve host ecosystem. New Phytologist, 2014, 201, 940-950.	7.3	115
56	Efficacy and Mechanisms of Communication Disruption of the Red Clover Casebearer Moth (Coleophora deauratella) with Complete and Partial Pheromone Formulations. Journal of Chemical Ecology, 2014, 40, 577-589.	1.8	7
57	The potential for pheromone-based monitoring to predict larval populations of diamondback moth, Plutella xylostella (L.), in canola (Brassica napus L.). Crop Protection, 2013, 45, 89-97.	2.1	14
58	When mating disruption does not disrupt mating: fitness consequences of delayed mating in moths. Entomologia Experimentalis Et Applicata, $2013$ , $146$ , $50$ - $65$ .	1.4	38
59	The Lodgepole × Jack Pine Hybrid Zone in Alberta, Canada: A Stepping Stone for the Mountain Pine Beetle on its Journey East Across the Boreal Forest?. Journal of Chemical Ecology, 2013, 39, 1209-1220.	1.8	32
60	Parasitoid complex and bionomics of (i) Apanteles polychrosidis (li) (Hymenoptera: Braconidae) on the ash leaf-cone roller (Lepidoptera: Gracillariidae). Canadian Entomologist, 2013, 145, 416-429.	0.8	5
61	Factors Affecting Pheromone-Baited Trap Capture of Male <l>Coleophora deauratella</l> , an Invasive Pest of Clover in Canada. Journal of Economic Entomology, 2013, 106, 844-854.	1.8	19
62	Influence of Seeding Rate, Nitrogen Management, and Micronutrient Blend Applications on Pith Expression in Solidâ€Stemmed Spring Wheat. Crop Science, 2012, 52, 1316-1329.	1.8	23
63	Developmental and Behavioral Effects and Retention of Incremental Rates of Rubidium Fed to Grapholita molesta (Lepidoptera: Tortricidae) in Dietary Medium. Journal of Entomological Science, 2012, 47, 316-326.	0.3	1
64	Smallâ€plot studies comparing pheromone and juice baits for massâ€trapping invasive <i><scp>S</scp>ynanthedon myopaeformis</i> in <scp>C</scp> anada. Entomologia Experimentalis Et Applicata, 2012, 145, 102-114.	1.4	5
65	The effect of flight on reproduction in an outbreaking forest lepidopteran. Physiological Entomology, 2012, 37, 219-226.	1.5	16
66	Do Interactions between Residue Management and Direct Seeding System Affect Wheat Stem Sawfly and Grain Yield?. Agronomy Journal, 2011, 103, 1635-1644.	1.8	6
67	Effect of Water Stress and Fungal Inoculation on Monoterpene Emission from an Historical and a New Pine Host of the Mountain Pine Beetle. Journal of Chemical Ecology, 2011, 37, 1013-1026.	1.8	47
68	Biology and management of North American cone-feeding <i>Dioryctria</i> species. Canadian Entomologist, 2011, 143, 1-34.	0.8	13
69	Phoretic mite associates of mountain pine beetle at the leading edge of an infestation in northwestern Alberta, Canada. Canadian Entomologist, 2011, 143, 44-55.	0.8	14
70	Sex pheromone of the red clover casebearer moth, <i>Coleophora deauratella</i> , an invasive pest of clover in Canada. Entomologia Experimentalis Et Applicata, 2010, 137, 255-261.	1.4	10
71	Assessment of Commercially Available Pheromone Lures for Monitoring Diamondback Moth (Lepidoptera: Plutellidae) in Canola. Journal of Economic Entomology, 2010, 103, 654-661.	1.8	9
72	Factors influencing flight potential of <i>Choristoneura conflictana</i> . Physiological Entomology, 2009, 34, 71-78.	1.5	21

#	Article	IF	Citations
73	Peripheral and behavioral plasticity of pheromone response and its hormonal control in a long-lived moth. Journal of Experimental Biology, 2009, 212, 2000-2006.	1.7	19
74	Biology of <i>Caloptilia fraxinella</i> (Lepidoptera: Gracillariidae) on ornamental green ash, <i>Fraxinus pennsylvanica</i> (Oleaceae). Canadian Entomologist, 2009, 141, 31-39.	0.8	10
75	Effect of horticultural oil on oviposition behaviour and egg survival in the obliquebanded leafroller (Lepidoptera: Tortricidae). Canadian Entomologist, 2009, 141, 86-94.	0.8	15
76	Development of a Combined Sex Pheromone-based Monitoring System for <i>Malacosoma disstria </i> (Lepidoptera: Lasoicampidae) and <i>Choristoneura conflictana </i> (Lepidoptera:) Tj ETQq0 0 0 rgBT /0	Ov <b>e</b> rłock 1	.0 <b>⊉</b> 650 617
77	Cocoonâ€spinning larvae of Oriental fruit moth and Indianmeal moth do not produce aggregation pheromone. Agricultural and Forest Entomology, 2009, 11, 205-212.	1.3	3
78	Mechanisms of Pheromone Communication Disruption in Choristoneura rosaceana Exposed to Microencapsulated (Z)-11-tetradecenyl Acetate Formulated with and Without Horticultural Oil. Journal of Chemical Ecology, 2008, 34, 1096-1106.	1.8	9
79	Attractiveness and toxicity of an attracticide formulation on adult males of ash leaf cone roller, CaloptiliaÂfraxinella. Entomologia Experimentalis Et Applicata, 2008, 127, 30-38.	1.4	4
80	Plasticity of male response to sex pheromone depends on physiological state in a long-lived moth. Animal Behaviour, 2008, 75, 663-672.	1.9	16
81	Ecological applications of pheromone trapping of <i>Malacosoma disstria </i> and <i>Choristoneura conflictana </i> . Canadian Entomologist, 2008, 140, 573-581.	0.8	10
82	Disruption of Pheromone Communication of <i>Choristoneura rosaceana</i> (Lepidoptera:) Tj ETQq0 0 0 rgBT /GENVIRONMENTAL Environmental Entomology, 2007, 36, 1189-1198.	Overlock 1 1.4	0 Tf 50 387 <sup>1</sup>
83	Effects of nutrition and methoprene treatment upon reproductive diapause in Caloptilia fraxinella (Lepidoptera: Gracillariidae). Physiological Entomology, 2007, 32, 275-282.	1.5	42
84	Sex pheromone of the large aspen tortrix, Choristoneura conflictana(Lepidoptera: Tortricidae). Chemoecology, 2006, 16, 115-122.	1.1	7
85	Body Size, Age, and Disease Influence Female Reproductive Performance in <i>Choristoneura conflictana</i> (Lepidoptera: Tortricidae). Annals of the Entomological Society of America, 2006, 99, 837-844.	2.5	21
86	Male Oriental Fruit Moth Response to a Combined Pheromone-Based Attracticide Formulation Targeting Both Oriental Fruit Moth and Codling Moth (Lepidoptera: Tortricidae). Journal of Economic Entomology, 2005, 98, 317-325.	1.8	22
87	Effects of Exposure to Pheromone and Insecticide Constituents of an Attracticide Formulation on Reproductive Behavior of Oriental Fruit Moth (Lepidoptera: Tortricidae). Journal of Economic Entomology, 2005, 98, 334-341.	1.8	6
88	Potential for combining sex pheromones for the forest tent caterpillar (Lepidoptera: Lasiocampidae) and the large aspen tortrix (Lepidoptera: Tortricidae) within monitoring traps targeting both species. Canadian Entomologist, 2005, 137, 615-619.	0.8	6
89	Male Oriental Fruit Moth Response to a Combined Pheromone-Based Attracticide Formulation Targeting Both Oriental Fruit Moth and Codling Moth (Lepidoptera: Tortricidae). Journal of Economic Entomology, 2005, 98, 317-325.	1.8	2
90	Effects of Exposure to Pheromone and Insecticide Constituents of an Attracticide Formulation on Reproductive Behavior of Oriental Fruit Moth (Lepidoptera: Tortricidae). Journal of Economic Entomology, 2005, 98, 334-341.	1.8	2

#	Article	IF	CITATIONS
91	Initial Development of an Attracticide Formulation Against the Oriental Fruit Moth, <i>Grapholita molesta</i> (Lepidoptera: Tortricidae). Environmental Entomology, 2004, 33, 213-220.	1.4	18
92	Factors influencing the effectiveness of an attracticide formulation against the Oriental fruit moth, Grapholita molesta. Entomologia Experimentalis Et Applicata, 2004, 112, 89-97.	1.4	23
93	Assessing the Mating Status of Male Obliquebanded Leafrollers <1>Choristoneura rosaceana 1 (Lepidoptera: Tortricidae) by Dissection of Male and Female Moths. Annals of the Entomological Society of America, 2003, 96, 217-224.	2.5	9
94	Inheritance and evolution of male response to sex pheromone in Trichoplusia ni (Lepidoptera:) Tj ETQq0 0 0 rgBT	Overlock	10 Tf 50 622
95	Potential for the evolution of resistance to pheromone-based mating disruption tested using two pheromone strains of the cabbage looper, Trichoplusia ni. Entomologia Experimentalis Et Applicata, 2001, 100, 131-134.	1.4	27
96	Title is missing!. Journal of Insect Behavior, 2000, 13, 499-510.	0.7	26
97	Mating Disruption of Two Sympatric, Orchard-Inhabiting Tortricids, Choristoneura rosaceana and Pandemis limitata (Lepidoptera: Tortricidae), with Pheromone Components of Both Species' Blends. Journal of Economic Entomology, 1999, 92, 380-390.	1.8	41
98	Pheromone-mediated mating disruption of Choristoneura rosaceana: is the most attractive blend really the most effective?. Entomologia Experimentalis Et Applicata, 1999, 90, 37-47.	1.4	39
99	Title is missing!. Journal of Chemical Ecology, 1999, 25, 501-517.	1.8	33
100	A synomone imparting distinct sex pheromone communication channels for Choristoneura rosaceana (Harris) and Pandemis limitata (Robinson) (Lepidoptera: Tortricidae). Chemoecology, 1999, 9, 73-80.	1.1	19
101	Predictive Capabilities of a Pheromone-Based Monitoring System for Western Hemlock Looper (Lepidoptera: Geometridae). Environmental Entomology, 1995, 24, 933-943.	1.4	22
102	Development of a Pheromone-Based Monitoring System for Western Hemlock Looper (Lepidoptera:) Tj ETQq0 0 923-932.	0 rgBT /Ov 1.4	verlock 10 Tf ! 21

Biology and management of the generalist herbivore, the bertha armyworm, iMamestra configurata/i (Lepidoptera: Noctuidae), on canola in western Canada.. , 0, , 114-129.