

Pietro Caironi

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7415312/publications.pdf>

Version: 2024-02-01

23
papers

1,705
citations

623734

14
h-index

642732

23
g-index

23
all docs

23
docs citations

23
times ranked

2193
citing authors

#	ARTICLE	IF	CITATIONS
1	A combined role for low vitamin D and low albumin circulating levels as strong predictors of worse outcome in COVID-19 patients. <i>Irish Journal of Medical Science</i> , 2023, 192, 423-430.	1.5	5
2	Emergency room comprehensive assessment of demographic, radiological, laboratory and clinical data of patients with COVID-19: determination of its prognostic value for in-hospital mortality. <i>Internal and Emergency Medicine</i> , 2022, 17, 205-214.	2.0	9
3	Lung Response to a Higher Positive End-Expiratory Pressure in Mechanically Ventilated Patients With COVID-19. <i>Chest</i> , 2022, 161, 979-988.	0.8	30
4	The Impact of Serum Albumin Levels on COVID-19 Mortality. <i>Infectious Disease Reports</i> , 2022, 14, 278-286.	3.1	12
5	PCSK9 is associated with mortality in patients with septic shock: data from the ALBIOS study. <i>Journal of Internal Medicine</i> , 2021, 289, 179-192.	6.0	27
6	Effects of neuromuscular electrical stimulation therapy on physical function in patients with COVID-19 associated pneumonia: Study protocol of a randomized controlled trial. <i>Contemporary Clinical Trials Communications</i> , 2021, 21, 100742.	1.1	3
7	Fluid-induced harm in the hospital: look beyond volume and start considering sodium. From physiology towards recommendations for daily practice in hospitalized adults. <i>Annals of Intensive Care</i> , 2021, 11, 79.	4.6	22
8	Paradoxical Effect of Chest Wall Compression on Respiratory System Compliance. <i>Chest</i> , 2021, 160, 1335-1339.	0.8	27
9	Higher levels of IgA and IgG at sepsis onset are associated with higher mortality: results from the Albumin Italian Outcome Sepsis (ALBIOS) trial. <i>Annals of Intensive Care</i> , 2021, 11, 161.	4.6	6
10	Orthodeoxia and its implications on awake-proning in COVID-19 pneumonia. <i>Critical Care</i> , 2021, 25, 429.	5.8	4
11	Extracorporeal Chloride Removal by Electrodialysis. A Novel Approach to Correct Acidemia. <i>American Journal of Respiratory and Critical Care Medicine</i> , 2020, 201, 799-813.	5.6	16
12	Early osteopontin levels predict mortality in patients with septic shock. <i>European Journal of Internal Medicine</i> , 2020, 78, 113-120.	2.2	15
13	The role of resistin and myeloperoxidase in severe sepsis and septic shock: Results from the ALBIOS trial. <i>European Journal of Clinical Investigation</i> , 2020, 50, e13333.	3.4	17
14	D-dimer corrected for thrombin and plasmin generation is a strong predictor of mortality in patients with sepsis. <i>Blood Transfusion</i> , 2020, 18, 304-311.	0.4	16
15	Low D-dimer levels in sepsis: Good or bad?. <i>Thrombosis Research</i> , 2019, 174, 13-15.	1.7	30
16	Persistence of Central Venous Oxygen Desaturation During Early Sepsis Is Associated With Higher Mortality. <i>Chest</i> , 2018, 154, 1291-1300.	0.8	18
17	Circulating Biologically Active Adrenomedullin (bio-ADM) Predicts Hemodynamic Support Requirement and Mortality During Sepsis. <i>Chest</i> , 2017, 152, 312-320.	0.8	59
18	Pentraxin 3 in patients with severe sepsis or shock: the ALBIOS trial. <i>European Journal of Clinical Investigation</i> , 2017, 47, 73-83.	3.4	71

#	ARTICLE	IF	CITATIONS
19	Sequential N-Terminal Pro-B-Type Natriuretic Peptide and High-Sensitivity Cardiac Troponin Measurements During Albumin Replacement in Patients With Severe Sepsis or Septic Shock*. <i>Critical Care Medicine</i> , 2016, 44, 707-716.	0.9	75
20	Cardiac dysfunction and circulating cardiac markers during sepsis. <i>Minerva Anestesiologica</i> , 2016, 82, 697-710.	1.0	7
21	Circulating presepsin (soluble CD14 subtype) as a marker of host response in patients with severe sepsis or septic shock: data from the multicenter, randomized ALBIOS trial. <i>Intensive Care Medicine</i> , 2015, 41, 12-20.	8.2	114
22	Albumin Replacement in Patients with Severe Sepsis or Septic Shock. <i>New England Journal of Medicine</i> , 2014, 370, 1412-1421.	27.0	947
23	Presepsin (soluble CD14 subtype) and procalcitonin levels for mortality prediction in sepsis: data from the Albumin Italian Outcome Sepsis trial. <i>Critical Care</i> , 2014, 18, R6.	5.8	175