

Daniel W Armstrong

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7420808/publications.pdf>

Version: 2024-02-01

475
papers

29,306
citations

4120

87
h-index

7931

149
g-index

480
all docs

480
docs citations

480
times ranked

13050
citing authors

#	ARTICLE	IF	CITATIONS
1	Characterizing Ionic Liquids On the Basis of Multiple Solvation Interactions. Journal of the American Chemical Society, 2002, 124, 14247-14254.	6.6	1,036
2	Ionic Liquids in Separations. Accounts of Chemical Research, 2007, 40, 1079-1086.	7.6	967
3	Structure and Properties of High Stability Geminal Dicationic Ionic Liquids. Journal of the American Chemical Society, 2005, 127, 593-604.	6.6	712
4	Macrocyclic Antibiotics as a New Class of Chiral Selectors for Liquid Chromatography. Analytical Chemistry, 1994, 66, 1473-1484.	3.2	702
5	Ionic liquids in analytical chemistry. Analytica Chimica Acta, 2010, 661, 1-16.	2.6	670
6	Examination of Ionic Liquids and Their Interaction with Molecules, When Used as Stationary Phases in Gas Chromatography. Analytical Chemistry, 1999, 71, 3873-3876.	3.2	615
7	High-Stability Ionic Liquids. A New Class of Stationary Phases for Gas Chromatography. Analytical Chemistry, 2003, 75, 4851-4858.	3.2	455
8	Ionic Liquids as Matrixes for Matrix-Assisted Laser Desorption/Ionization Mass Spectrometry. Analytical Chemistry, 2001, 73, 3679-3686.	3.2	453
9	Ionic Liquids in Analytical Chemistry. Analytical Chemistry, 2006, 78, 2892-2902.	3.2	433
10	Partitioning behavior of solutes eluted with micellar mobile phases in liquid chromatography. Analytical Chemistry, 1981, 53, 1662-1666.	3.2	426
11	Immobilized Ionic Liquids as High-Selectivity/High-Temperature/High-Stability Gas Chromatography Stationary Phases. Analytical Chemistry, 2005, 77, 6453-6462.	3.2	388
12	A covalently bonded teicoplanin chiral stationary phase for HPLC enantioseparations. Chirality, 1995, 7, 474-497.	1.3	327
13	Evaluation of the macrocyclic antibiotic vancomycin as a chiral selector for capillary electrophoresis. Chirality, 1994, 6, 496-509.	1.3	302
14	Facile liquid chromatographic enantioresolution of native amino acids and peptides using a teicoplanin chiral stationary phase. Journal of Chromatography A, 1996, 731, 123-137.	1.8	294
15	Liquid chromatographic separation of enantiomers using a chiral .beta.-cyclodextrin-bonded stationary phase and conventional aqueous-organic mobile phases. Analytical Chemistry, 1985, 57, 237-242.	3.2	275
16	Chiral Ionic Liquids as Stationary Phases in Gas Chromatography. Analytical Chemistry, 2004, 76, 6819-6822.	3.2	275
17	Examination of the origin, variation, and proper use of expressions for the estimation of association constants by capillary electrophoresis. Journal of Chromatography A, 1996, 721, 173-186.	1.8	273
18	Chiral ionic liquids: Synthesis and applications. Chirality, 2005, 17, 281-292.	1.3	272

#	ARTICLE	IF	CITATIONS
19	Comparison and Modeling Study of Vancomycin, Ristocetin A, and Teicoplanin for CE Enantioseparations. <i>Analytical Chemistry</i> , 1996, 68, 2501-2514.	3.2	239
20	Use of a Macrocyclic Antibiotic, Rifamycin B, and Indirect Detection for the Resolution of Racemic Amino Alcohols by CE. <i>Analytical Chemistry</i> , 1994, 66, 1690-1695.	3.2	230
21	Unsymmetrical Dicationic Ionic Liquids: Manipulation of Physicochemical Properties Using Specific Structural Architectures. <i>Chemistry of Materials</i> , 2007, 19, 5848-5850.	3.2	216
22	Role of the Carbohydrate Moieties in Chiral Recognition on Teicoplanin-Based LC Stationary Phases. <i>Analytical Chemistry</i> , 2000, 72, 1767-1780.	3.2	213
23	Use of an Aqueous Micellar Mobile Phase for Separation of Phenols and Polynuclear Aromatic Hydrocarbons via HPLC. <i>Journal of Liquid Chromatography and Related Technologies</i> , 1980, 3, 657-662.	0.9	208
24	Multiple enantioselective retention mechanisms on derivatized cyclodextrin gas chromatographic chiral stationary phases.. <i>Analytical Chemistry</i> , 1992, 64, 873-879.	3.2	205
25	Chiral Stationary Phases for High Performance Liquid Chromatographic Separation of Enantiomers: A Mini-Review. <i>Journal of Liquid Chromatography and Related Technologies</i> , 1984, 7, 353-376.	0.9	203
26	Derivatized cyclodextrins for normal-phase liquid chromatographic separation of enantiomers. <i>Analytical Chemistry</i> , 1990, 62, 1610-1615.	3.2	199
27	Improved Cyclodextrin Chiral Phases: A Comparison and Review. <i>Journal of Liquid Chromatography and Related Technologies</i> , 1986, 9, 407-423.	0.9	197
28	Using Geminal Dicationic Ionic Liquids as Solvents for High-Temperature Organic Reactions. <i>Organic Letters</i> , 2005, 7, 4205-4208.	2.4	197
29	Micelles in Separations: Practical and Theoretical Review. <i>Separation and Purification Reviews</i> , 1985, 14, 213-304.	0.8	194
30	Separation of Metallocene Enantiomers by Liquid Chromatography: Chiral Recognition Via Cyclodextrin Bonded Phases. <i>Analytical Chemistry</i> , 1985, 57, 481-484.	3.2	183
31	Methods for the determination of binding constants by capillary electrophoresis. <i>Electrophoresis</i> , 2001, 22, 1419-1427.	1.3	182
32	Separating Microbes in the Manner of Molecules. 1. Capillary Electrokinetic Approaches. <i>Analytical Chemistry</i> , 1999, 71, 5465-5469.	3.2	178
33	Derivatized cyclodextrins immobilized on fused-silica capillaries for enantiomeric separations via capillary electrophoresis, gas chromatography, or supercritical fluid chromatography. <i>Analytical Chemistry</i> , 1993, 65, 1114-1117.	3.2	173
34	Highly Efficient Asymmetric Direct Stoichiometric Aldol Reactions on/in Water. <i>Angewandte Chemie - International Edition</i> , 2007, 46, 9073-9077.	7.2	173
35	Liquid chromatographic separation of diastereomers and structural isomers on cyclodextrin-bonded phases. <i>Analytical Chemistry</i> , 1985, 57, 234-237.	3.2	171
36	Mechanism of Signal Suppression by Anionic Surfactants in Capillary Electrophoresis~Electrospray Ionization Mass Spectrometry. <i>Analytical Chemistry</i> , 1996, 68, 3493-3497.	3.2	171

#	ARTICLE	IF	CITATIONS
37	Separation and analysis of colloidal/nano-particles including microorganisms by capillary electrophoresis: a fundamental review. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2004, 800, 7-25.	1.2	166
38	Characterization of phosphonium ionic liquids through a linear solvation energy relationship and their use as GLC stationary phases. <i>Analytical and Bioanalytical Chemistry</i> , 2008, 390, 1605-1617.	1.9	163
39	Polar-liquid, derivatized cyclodextrin stationary phases for the capillary gas chromatography separation of enantiomers. <i>Analytical Chemistry</i> , 1990, 62, 914-923.	3.2	159
40	Development of New HPLC Chiral Stationary Phases Based on Native and Derivatized Cyclofructans. <i>Analytical Chemistry</i> , 2009, 81, 10215-10226.	3.2	157
41	Highly enantioselective capillary electrophoretic separations with dilute solutions of the macrocyclic antibiotic ristocetin A. <i>Journal of Chromatography A</i> , 1995, 689, 285-304.	1.8	153
42	Methods for the estimation of binding constants by capillary electrophoresis. <i>Electrophoresis</i> , 1997, 18, 2194-2202.	1.3	153
43	Advances in high-throughput and high-efficiency chiral liquid chromatographic separations. <i>Journal of Chromatography A</i> , 2016, 1467, 2-18.	1.8	153
44	PEG-linked geminal dicationic ionic liquids as selective, high-stability gas chromatographic stationary phases. <i>Analytical and Bioanalytical Chemistry</i> , 2007, 389, 2265-2275.	1.9	152
45	Highly enantioselective HPLC separations using the covalently bonded macrocyclic antibiotic, ristocetin A, chiral stationary phase. , 1998, 10, 434-483.		149
46	(S)-2-Hydroxypropyl- β -cyclodextrin, a new chiral stationary phase for reversed-phase liquid chromatography. <i>Journal of Chromatography A</i> , 1990, 513, 181-194.	1.8	147
47	Trigonal Tricationic Ionic Liquids: A Generation of Gas Chromatographic Stationary Phases. <i>Analytical Chemistry</i> , 2009, 81, 160-173.	3.2	146
48	Direct liquid chromatographic separation of racemates with an α -cyclodextrin bonded phase. <i>Analytical Chemistry</i> , 1987, 59, 2594-2596.	3.2	142
49	Gone in Seconds: Praxis, Performance, and Peculiarities of Ultrafast Chiral Liquid Chromatography with Superficially Porous Particles. <i>Analytical Chemistry</i> , 2015, 87, 9137-9148.	3.2	140
50	Ionic matrices for matrix-assisted laser desorption/ionization time-of-flight detection of DNA oligomers. <i>Rapid Communications in Mass Spectrometry</i> , 2003, 17, 553-560.	0.7	139
51	Capillary electrophoretic enantioseparations using macrocyclic antibiotics as chiral selectors. <i>Electrophoresis</i> , 1997, 18, 2331-2342.	1.3	137
52	Use of Chiral Ionic Liquids as Solvents for the Enantioselective Photoisomerization of Dibenzobicyclo[2.2.2]octatrienes. <i>Organic Letters</i> , 2005, 7, 335-337.	2.4	137
53	Use of CE for the determination of binding constants. <i>Electrophoresis</i> , 2010, 31, 17-27.	1.3	136
54	Dicationic ionic liquid stationary phase for GC-MS analysis of volatile compounds in herbal plants. <i>Analytical and Bioanalytical Chemistry</i> , 2007, 388, 889-899.	1.9	133

#	ARTICLE	IF	CITATIONS
55	Enzyme-Catalyzed Hydrolysis of Cellulose in Ionic Liquids: A Green Approach Toward the Production of Biofuels. <i>Journal of Physical Chemistry B</i> , 2010, 114, 8221-8227.	1.2	127
56	Ionic Liquids in Analytical Chemistry. <i>Annual Review of Analytical Chemistry</i> , 2009, 2, 145-168.	2.8	125
57	(R)- and (S)-Naphthylethylcarbamate-substituted β -cyclo-dextrin bonded stationary phases for the reversed-phase liquid chromatographic separation of enantiomers. <i>Journal of Chromatography A</i> , 1991, 539, 83-90.	1.8	124
58	Substituent effects on the binding of phenols to cyclodextrins in aqueous solution. <i>The Journal of Physical Chemistry</i> , 1989, 93, 6863-6867.	2.9	122
59	Effect of Micelles and Mixed Micelles on Efficiency and Selectivity of Antibiotic-Based Capillary Electrophoretic Enantioseparations. <i>Analytical Chemistry</i> , 1995, 67, 2088-2095.	3.2	121
60	High-performance liquid chromatographic separation of enantiomers of unusual amino acids on a teicoplanin chiral stationary phase. <i>Journal of Chromatography A</i> , 1998, 793, 283-296.	1.8	120
61	Separation, Identification, and Characterization of Microorganisms by Capillary Electrophoresis. <i>Microbiology and Molecular Biology Reviews</i> , 2003, 67, 38-51.	2.9	120
62	Chiral ionic liquids: A compendium of syntheses and applications (2005-2012). <i>Chirality</i> , 2012, 24, 17-53.	1.3	119
63	Ultrafast chiral separations for high throughput enantiopurity analysis. <i>Chemical Communications</i> , 2017, 53, 509-512.	2.2	117
64	Super/subcritical fluid chromatography chiral separations with macrocyclic glycopeptide stationary phases. <i>Journal of Chromatography A</i> , 2002, 978, 185-204.	1.8	113
65	Evaluation of the liquid chromatographic separation of monosaccharides, disaccharides, trisaccharides, tetrasaccharides, deoxysaccharides and sugar alcohols with stable cyclodextrin bonded phase columns. <i>Journal of Chromatography A</i> , 1989, 462, 219-232.	1.8	110
66	Thin Layer Chromatographic Separation of Ortho, Meta, and Para Substituted Benzoic Acids and Phenols with Aqueous Solutions of β -Cyclodextrin. <i>Analytical Letters</i> , 1980, 13, 1093-1104.	1.0	106
67	Use of hydroxypropyl- and hydroxyethyl-derivatized β -cyclodextrins for the thin-layer chromatographic separation of enantiomers and diastereomers. <i>Journal of Chromatography A</i> , 1988, 452, 323-330.	1.8	106
68	Enrichment of enantiomers and other isomers with aqueous liquid membranes containing cyclodextrin carriers. <i>Analytical Chemistry</i> , 1987, 59, 2237-2241.	3.2	105
69	Nicotine enantiomers and oxidative stress. <i>Toxicology</i> , 1998, 130, 155-165.	2.0	105
70	Bonded ionic liquid polymeric material for solid-phase microextraction GC analysis. <i>Analytical and Bioanalytical Chemistry</i> , 2010, 396, 511-524.	1.9	105
71	Micellar effects on molecular diffusion: theoretical and chromatographic considerations. <i>Analytical Chemistry</i> , 1986, 58, 579-582.	3.2	104
72	Enantiomeric separation of fluorescent, 6-aminoquinolyl-N-hydroxysuccinimidyl carbamate, tagged amino acids. <i>Journal of Chromatography A</i> , 1993, 641, 257-265.	1.8	104

#	ARTICLE	IF	CITATIONS
73	Theoretical considerations concerning the separation of enantiomeric solutes by liquid chromatography. <i>Analytical Chemistry</i> , 1988, 60, 522-528.	3.2	103
74	Effects of temperature on retention of chiral compounds on a ristocetin A chiral stationary phase. <i>Journal of Chromatography A</i> , 2002, 958, 89-107.	1.8	103
75	D-amino acid levels in human physiological fluids. <i>Chirality</i> , 1993, 5, 375-378.	1.3	102
76	Ultrafast Chiral Chromatography as the Second Dimension in Two-Dimensional Liquid Chromatography Experiments. <i>Analytical Chemistry</i> , 2017, 89, 3545-3553.	3.2	102
77	Determination of Cell Viability in Single or Mixed Samples Using Capillary Electrophoresis Laser-Induced Fluorescence Microfluidic Systems. <i>Analytical Chemistry</i> , 2001, 73, 4551-4557.	3.2	101
78	Evaluation of a Vancomycin Chiral Stationary Phase in Capillary Electrochromatography Using Polar Organic and Reversed-Phase Modes. <i>Analytical Chemistry</i> , 2000, 72, 4394-4401.	3.2	99
79	Effect of the configuration of the substituents of derivatized β -cyclodextrin bonded phases on enantioselectivity in normal-phase liquid chromatography. <i>Journal of Chromatography A</i> , 1991, 540, 113-128.	1.8	98
80	Capillary electrophoretic enantiomeric separations using the glycopeptide antibiotic, teicoplanin. <i>Chirality</i> , 1996, 8, 88-107.	1.3	98
81	Relevance of enantiomeric separations in food and beverage analyses. <i>Journal of Agricultural and Food Chemistry</i> , 1990, 38, 1674-1677.	2.4	97
82	Towards a second generation of ionic liquid matrices (ILMs) for MALDI-MS of Peptides, proteins, and carbohydrates. <i>Journal of the American Society for Mass Spectrometry</i> , 2009, 20, 1790-1800.	1.2	96
83	Rapid Identification of the Bacterial Pathogens Responsible for Urinary Tract Infections Using Direct Injection CE. <i>Analytical Chemistry</i> , 2000, 72, 4474-4476.	3.2	95
84	Reversing enantioselectivity in capillary gas chromatography with polar and nonpolar cyclodextrin derivative phases. <i>Analytical Chemistry</i> , 1990, 62, 214-217.	3.2	94
85	Evaluation of partition coefficients to micelles and cyclodextrins via planar chromatography. <i>Journal of the American Chemical Society</i> , 1983, 105, 2962-2964.	6.6	93
86	Effective enantiomeric separations of racemic primary amines by the isopropyl carbamate-cyclofructan6 chiral stationary phase. <i>Journal of Chromatography A</i> , 2010, 1217, 4904-4918.	1.8	93
87	β -Amino Acid Levels in Perfused Mouse Brain Tissue and Blood: A Comparative Study. <i>ACS Chemical Neuroscience</i> , 2017, 8, 1251-1261.	1.7	93
88	A General, Positive Ion Mode ESI-MS Approach for the Analysis of Singly Charged Inorganic and Organic Anions Using a Dicationic Reagent. <i>Analytical Chemistry</i> , 2007, 79, 7346-7352.	3.2	92
89	Rapid CE microbial assays for consumer products that contain active bacteria. <i>FEMS Microbiology Letters</i> , 2001, 194, 33-37.	0.7	88
90	Pseudophase Liquid Chromatography: Applications to TLC. <i>Journal of Liquid Chromatography and Related Technologies</i> , 1980, 3, 895-900.	0.9	87

#	ARTICLE	IF	CITATIONS
91	Evaluation of the macrocyclic antibiotic avoparcin as a new chiral selector for HPLC. , 1998, 10, 627-660.		87
92	Gas-Phase Ion Association Provides Increased Selectivity and Sensitivity for Measuring Perchlorate by Mass Spectrometry. Analytical Chemistry, 2005, 77, 4829-4835.	3.2	84
93	Empirical procedure that uses molecular structure to predict enantioselectivity of chiral stationary phases. Analytical Chemistry, 1992, 64, 395-404.	3.2	82
94	Salient Sub-Second Separations. Analytical Chemistry, 2016, 88, 8821-8826.	3.2	82
95	Development of dinitrophenylated cyclodextrin derivatives for enhanced enantiomeric separations by high-performance liquid chromatography. Journal of Chromatography A, 2006, 1115, 19-45.	1.8	80
96	On-Chip Drop-to-Drop Liquid Microextraction Coupled with Real-Time Concentration Monitoring Technique. Analytical Chemistry, 2011, 83, 1658-1664.	3.2	80
97	Comprehensive two-dimensional gas chromatography using a high-temperature phosphonium ionic liquid column. Analytical and Bioanalytical Chemistry, 2008, 390, 323-332.	1.9	79
98	Evaluation of A Chiral Crown Etherlc Column For the Separation of Racemic Amines. Journal of Liquid Chromatography and Related Technologies, 1991, 14, 9-28.	0.9	78
99	A Fundamental Study on Electrowetting by Traditional and Multifunctional Ionic Liquids: Possible Use in Electrowetting on Dielectric-Based Microfluidic Applications. Analytical Chemistry, 2008, 80, 7690-7698.	3.2	77
100	Ionic cyclodextrins in ionic liquid matrices as chiral stationary phases for gas chromatography. Journal of Chromatography A, 2010, 1217, 5261-5273.	1.8	77
101	Synthesis of Thermally Stable Geminal Dicationic Ionic Liquids and Related Ionic Compounds: An Examination of Physicochemical Properties by Structural Modification. Chemistry of Materials, 2016, 28, 4315-4323.	3.2	77
102	Enantiomeric resolution and chiral recognition of racemic nicotine and nicotine analogs by .beta.-cyclodextrin complexation. Structure-enantiomeric resolution relationships in host-guest interactions. Analytical Chemistry, 1988, 60, 2120-2127.	3.2	76
103	Evaluation of the Enantiomeric Separation of Dipeptides Using a Chiral Crown Ether LC Column. Journal of Liquid Chromatography and Related Technologies, 1991, 14, 3673-3683.	0.9	75
104	Sampling frequency, response times and embedded signal filtration in fast, high efficiency liquid chromatography: A tutorial. Analytica Chimica Acta, 2016, 907, 31-44.	2.6	75
105	Cyclodextrin chiral stationary phases for liquid chromatographic separations of drug stereoisomers. Journal of Pharmaceutical and Biomedical Analysis, 1990, 8, 123-130.	1.4	74
106	Separation and characterization of underivatized oligosaccharides using liquid chromatography and liquid chromatographyâ€“electrospray ionization mass spectrometry. Journal of Chromatography A, 2005, 1079, 146-152.	1.8	74
107	Superficially porous particles vs. fully porous particles for bonded high performance liquid chromatographic chiral stationary phases: Isopropyl cyclofructan 6. Journal of Chromatography A, 2014, 1363, 89-95.	1.8	74
108	Single-Cell Detection:Â Test of Microbial Contamination Using Capillary Electrophoresis. Analytical Chemistry, 2007, 79, 1720-1724.	3.2	73

#	ARTICLE	IF	CITATIONS
109	Cyclofructan 6 based stationary phases for hydrophilic interaction liquid chromatography. <i>Journal of Chromatography A</i> , 2011, 1218, 270-279.	1.8	73
110	High efficiency, narrow particle size distribution, sub-2 μ m based macrocyclic glycopeptide chiral stationary phases in HPLC and SFC. <i>Analytica Chimica Acta</i> , 2015, 898, 128-137.	2.6	73
111	Synthesis, rapid resolution, and determination of absolute configuration of racemic 2,2'-binaphthylidyl crown ethers and analogs via β -cyclodextrin complexation. <i>Journal of Organic Chemistry</i> , 1985, 50, 5556-5559.	1.7	72
112	Planar chromatographic separation of enantiomers and diastereomers with cyclodextrin mobile phase additives. <i>Journal of Chromatography A</i> , 1988, 448, 345-354.	1.8	72
113	Efficient enantioselective separation and determination of trace impurities in secondary amino acids (i.e., imino acids). <i>Journal of Chromatography A</i> , 1992, 623, 33-41.	1.8	72
114	Practice and mechanism of HPLC oligosaccharide separation with a cyclodextrin bonded phase. <i>Talanta</i> , 1998, 47, 1001-1012.	2.9	72
115	Fundamental and Practical Insights on the Packing of Modern High-Efficiency Analytical and Capillary Columns. <i>Analytical Chemistry</i> , 2017, 89, 8177-8191.	3.2	72
116	Gas chromatography–vacuum ultraviolet spectroscopy for analysis of fatty acid methyl esters. <i>Food Chemistry</i> , 2016, 194, 265-271.	4.2	70
117	Elucidation of vancomycin's enantioselective binding site using its copper complex. <i>Chirality</i> , 1996, 8, 590-595.	1.3	69
118	Product Review: Chiral Stationary Phases for HPLC. <i>Analytical Chemistry</i> , 2001, 73, 557 A-561 A.	3.2	67
119	Chiral separation of monoterpenes using mixtures of sulfated β -cyclodextrins and β -cyclodextrin as chiral additives in the reversed-polarity capillary electrophoresis mode. <i>Journal of Chromatography A</i> , 1997, 759, 149-155.	1.8	66
120	Mechanistic Aspects in the Generation of Apparent Ultrahigh Efficiencies for Colloidal (Microbial) Electrokinetic Separations. <i>Analytical Chemistry</i> , 2002, 74, 5523-5530.	3.2	66
121	Considerations on HILIC and polar organic solvent-based separations: Use of cyclodextrin and macrocyclic glycopeptide stationary phases. <i>Journal of Separation Science</i> , 2008, 31, 1980-1990.	1.3	66
122	Enantiomeric composition of nicotine in smokeless tobacco, medicinal products, and commercial reagents. <i>Chirality</i> , 1998, 10, 587-591.	1.3	65
123	Capillary Electrophoretic Method for the Detection of Bacterial Contamination. <i>Analytical Chemistry</i> , 2006, 78, 4759-4767.	3.2	65
124	Separation of chiral sulfoxides by liquid chromatography using macrocyclic glycopeptide chiral stationary phases. <i>Journal of Chromatography A</i> , 2002, 955, 53-69.	1.8	63
125	Coronatine Facilitates <i>Pseudomonas syringae</i> Infection of <i>Arabidopsis</i> Leaves at Night. <i>Frontiers in Plant Science</i> , 2016, 7, 880.	1.7	63
126	Monitoring the migration behavior of living microorganisms in capillary electrophoresis using laser-induced fluorescence detection with a charge-coupled device imaging system. <i>Electrophoresis</i> , 2002, 23, 2048.	1.3	61

#	ARTICLE	IF	CITATIONS
127	Trigonal Tricationic Ionic Liquids: Molecular Engineering of Trications to Control Physicochemical Properties. <i>Chemistry of Materials</i> , 2008, 20, 4182-4184.	3.2	61
128	Increasing selectivity in comprehensive three-dimensional gas chromatography via an ionic liquid stationary phase column in one dimension. <i>Journal of Chromatography A</i> , 2010, 1217, 3144-3149.	1.8	60
129	Ultrafast separation of fluorinated and desfluorinated pharmaceuticals using highly efficient and selective chiral selectors bonded to superficially porous particles. <i>Journal of Chromatography A</i> , 2015, 1426, 241-247.	1.8	59
130	Quinine bonded to superficially porous particles for high-efficiency and ultrafast liquid and supercritical fluid chromatography. <i>Analytica Chimica Acta</i> , 2017, 963, 164-174.	2.6	58
131	Separation of Mycotoxins, Polycyclic Aromatic Hydrocarbons, Quinones, and Heterocyclic Compounds on Cyclodextrin Bonded Phases: An Alternative LC Packing. <i>Journal of Liquid Chromatography and Related Technologies</i> , 1985, 8, 261-269.	0.9	57
132	Rapid determination of complex mixtures by dual-column gas chromatography with a novel stationary phase combination and spectrometric detection. <i>Journal of Chromatography A</i> , 2006, 1135, 230-240.	1.8	57
133	Centrifugal Partition Chromatography. I. General Features. <i>Journal of Liquid Chromatography and Related Technologies</i> , 1988, 11, 547-566.	0.9	56
134	Evaluation of the macrocyclic glycopeptide A-40,926 as a high-performance liquid chromatographic chiral selector and comparison with teicoplanin chiral stationary phase. <i>Journal of Chromatography A</i> , 2000, 897, 113-129.	1.8	55
135	The Effect of AC Frequency on the Electrowetting Behavior of Ionic Liquids. <i>Analytical Chemistry</i> , 2010, 82, 3146-3154.	3.2	55
136	Rapid Analysis of Ethanol and Water in Commercial Products Using Ionic Liquid Capillary Gas Chromatography with Thermal Conductivity Detection and/or Barrier Discharge Ionization Detection. <i>Journal of Agricultural and Food Chemistry</i> , 2014, 62, 1832-1838.	2.4	55
137	Separation of optical isomers of scopolamine, cocaine, homatropine, and atropine. <i>Analytical Biochemistry</i> , 1987, 167, 261-264.	1.1	53
138	Analysis of native amino acid and peptide enantiomers by high-performance liquid chromatography/atmospheric pressure chemical ionization mass spectrometry. <i>Journal of Mass Spectrometry</i> , 2004, 39, 177-187.	0.7	53
139	Evaluation of dicationic reagents for their use in detection of anions using positive ion mode ESI-MS via gas phase ion association. <i>Journal of the American Society for Mass Spectrometry</i> , 2008, 19, 261-269.	1.2	53
140	Nanopore Stochastic Detection of a Liquid Explosive Component and Sensitizers Using Boromycin and an Ionic Liquid Supporting Electrolyte. <i>Analytical Chemistry</i> , 2009, 81, 460-464.	3.2	53
141	Evaluation of aromatic-derivatized cyclofructans 6 and 7 as HPLC chiral selectors. <i>Analyst, The</i> , 2011, 136, 787-800.	1.7	53
142	Mechanism of enhancement of analyte sensitivity by surfactants in flame atomic spectrometry. <i>Analytical Chemistry</i> , 1982, 54, 1325-1329.	3.2	52
143	Analysis of derivatized and underivatized theanine enantiomers by high-performance liquid chromatography/atmospheric pressure ionization-mass spectrometry. <i>Rapid Communications in Mass Spectrometry</i> , 2004, 18, 251-256.	0.7	52
144	Enantiomeric Separations of Ruthenium(II) Polypyridyl Complexes Using High-Performance Liquid Chromatography (HPLC) with Cyclodextrin Chiral Stationary Phases (CSPs). <i>Inorganic Chemistry</i> , 2007, 46, 10312-10320.	1.9	52

#	ARTICLE	IF	CITATIONS
145	Synthesis and examination of sulfated cyclofructans as a novel class of chiral selectors for CE. <i>Electrophoresis</i> , 2009, 30, 3897-3909.	1.3	52
146	Analysis of Long-Chain Unsaturated Fatty Acids by Ionic Liquid Gas Chromatography. <i>Journal of Agricultural and Food Chemistry</i> , 2016, 64, 1422-1432.	2.4	52
147	Evaluation of enantiomeric purity of selected amino acids in honey. <i>Chirality</i> , 1994, 6, 270-276.	1.3	51
148	Two-dimensional high performance liquid chromatography for determination of homocysteine, methionine and cysteine enantiomers in human serum. <i>Journal of Chromatography A</i> , 2015, 1408, 118-124.	1.8	51
149	Determination and use of Rohrschneider-McReynolds constants for chiral stationary phases used in capillary gas chromatography. <i>Analytical Chemistry</i> , 1995, 67, 849-857.	3.2	50
150	Derivatized vancomycin stationary phases for LC chiral separations. <i>Talanta</i> , 1996, 43, 1767-1782.	2.9	50
151	High-performance liquid chromatographic separation of enantiomers of synthetic amino acids on a ristocetin A chiral stationary phase. <i>Journal of Chromatography A</i> , 2000, 904, 1-15.	1.8	50
152	RLIP76 in Defense of Radiation Poisoning. <i>International Journal of Radiation Oncology Biology Physics</i> , 2008, 72, 553-561.	0.4	50
153	Development and evaluation of new zwitterionic Hydrophilic interaction liquid chromatography stationary phases based on 3-P,P-diphenylphosphonium-propylsulfonate. <i>Journal of Chromatography A</i> , 2011, 1218, 8075-8082.	1.8	49
154	Separations at the Speed of Sensors. <i>Analytical Chemistry</i> , 2018, 90, 3349-3356.	3.2	49
155	Enantiomeric composition of nornicotine, anatabine, and anabasine in tobacco. <i>Chirality</i> , 1999, 11, 82-84.	1.3	48
156	Evaluating the Use of Tricationic Reagents for the Detection of Doubly Charged Anions in the Positive Mode by ESI-MS. <i>Analytical Chemistry</i> , 2008, 80, 2612-2616.	3.2	48
157	Macrocyclic glycopeptide chiral selectors bonded to core-shell particles enables enantiopurity analysis of the entire verubecestat synthetic route. <i>Journal of Chromatography A</i> , 2018, 1539, 87-92.	1.8	48
158	CE Resolution of Neutral and Anionic Racemates with Glycopeptide Antibiotics and Micelles. <i>Journal of Liquid Chromatography and Related Technologies</i> , 1995, 18, 3659-3674.	0.9	46
159	Comparison of the separation efficiencies of Chirobiotic T and TAG columns in the separation of unusual amino acids. <i>Journal of Chromatography A</i> , 2004, 1031, 159-170.	1.8	46
160	Determination of Trace Water Content in Petroleum and Petroleum Products. <i>Analytical Chemistry</i> , 2016, 88, 8194-8201.	3.2	46
161	Enantiomeric impurities in chiral catalysts, auxiliaries and synthons used in enantioselective synthesis. <i>Tetrahedron: Asymmetry</i> , 1998, 9, 2043-2064.	1.8	45
162	Comparison of superficially porous and fully porous silica supports used for a cyclofructan 6 hydrophilic interaction liquid chromatographic stationary phase. <i>Journal of Chromatography A</i> , 2014, 1365, 124-130.	1.8	45

#	ARTICLE	IF	CITATIONS
163	Ramifications and Insights on the Role of Water in Chiral Sub/Supercritical Fluid Chromatography. <i>Analytical Chemistry</i> , 2019, 91, 14672-14680.	3.2	45
164	β -Cyclodextrin chiral stationary phases for liquid chromatography. Effect of the spacer arm on chiral recognition. <i>Talanta</i> , 1993, 40, 1367-1373.	2.9	44
165	Direct and indirect high-performance liquid chromatographic enantioseparation of β -amino acids. <i>Journal of Chromatography A</i> , 2004, 1031, 171-178.	1.8	44
166	A second-generation ionic liquid matrix-assisted laser desorption/ionization matrix for effective mass spectrometric analysis of biodegradable polymers. <i>Rapid Communications in Mass Spectrometry</i> , 2009, 23, 3409-3422.	0.7	44
167	1,3-Dimethylamylamine (DMAA) in supplements and geranium products: natural or synthetic?. <i>Drug Testing and Analysis</i> , 2012, 4, 986-990.	1.6	44
168	Evaluation and comparison of a methylated teicoplanin aglycone to teicoplanin aglycone and natural teicoplanin chiral stationary phases. <i>Journal of Separation Science</i> , 2006, 29, 429-445.	1.3	43
169	The use of cyclofructans as novel chiral selectors for gas chromatography. <i>Analyst</i> , 2010, 135, 1076.	1.7	43
170	Sulfonated cyclofructan 6 based stationary phase for hydrophilic interaction chromatography. <i>Journal of Separation Science</i> , 2011, 34, 1636-1647.	1.3	43
171	Enhanced comprehensive two-dimensional gas chromatographic resolution of polychlorinated biphenyls on a non-polar polysiloxane and an ionic liquid column series. <i>Journal of Chromatography A</i> , 2011, 1218, 746-751.	1.8	43
172	Instrumental Idiosyncrasies Affecting the Performance of Ultrafast Chiral and Achiral Sub/Supercritical Fluid Chromatography. <i>Analytical Chemistry</i> , 2016, 88, 8664-8672.	3.2	43
173	Enantiomeric impurities in chiral catalysts, auxiliaries, synthons and resolving agents. Part 2. <i>Tetrahedron: Asymmetry</i> , 1999, 10, 37-60.	1.8	42
174	Centrifugal Partition Chromatography. V. Octanol-Water Partition Coefficients, Direct and Indirect Determination. <i>Journal of Liquid Chromatography and Related Technologies</i> , 1988, 11, 1441-1456.	0.9	41
175	Liquid chromatographic separation of anomeric forms of saccharides with cyclodextrin bonded phases. <i>Chirality</i> , 1989, 1, 27-37.	1.3	41
176	Evaluation of Flexible Linear Tricationic Salts as Gas-Phase Ion-Pairing Reagents for the Detection of Divalent Anions in Positive Mode ESI-MS. <i>Analytical Chemistry</i> , 2008, 80, 8828-8834.	3.2	41
177	Physicochemical properties of branched-chain dicationic ionic liquids. <i>Journal of Molecular Liquids</i> , 2018, 256, 247-255.	2.3	41
178	Practice, Mechanism and Theory of Reversed Phase TLC Polymer Fractionation. <i>Journal of Liquid Chromatography and Related Technologies</i> , 1983, 6, 1-22.	0.9	40
179	Facile Resolution OF N-tert-Butoxy-Carbonyl Amino Acids: The Importance of Enantiomeric Purity in Peptide Synthesis. <i>Journal of Liquid Chromatography and Related Technologies</i> , 1992, 15, 1411-1429.	0.9	40
180	Effective methodologies for enantiomeric separations of 150 pharmacology and toxicology related 1 $^{\circ}$, 2 $^{\circ}$, and 3 $^{\circ}$ amines with core-shell chiral stationary phases. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2018, 155, 70-81.	1.4	40

#	ARTICLE	IF	CITATIONS
181	Replacing methanol with azeotropic ethanol as the co-solvent for improved chiral separations with supercritical fluid chromatography (SFC). <i>Green Chemistry</i> , 2020, 22, 1249-1257.	4.6	40
182	Stereochemistry of pipecolic acid found in the urine and plasma of subjects with peroxisomal deficiencies. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 1993, 11, 881-886.	1.4	39
183	Separation of enantiomers by capillary electrophoresis using pentosan polysulfate. <i>Electrophoresis</i> , 1999, 20, 162-170.	1.3	39
184	Water determination in active pharmaceutical ingredients using ionic liquid headspace gas chromatography and two different detection protocols. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2014, 94, 111-117.	1.4	39
185	Quantification of aminobutyric acids and their clinical applications as biomarkers for osteoporosis. <i>Communications Biology</i> , 2020, 3, 39.	2.0	39
186	Cyclodextrin-based liquid chromatographic enantiomeric separation of chiral dihydrofurocoumarins, an emerging class of medicinal compounds. <i>Journal of Chromatography A</i> , 2003, 1011, 37-47.	1.8	38
187	Chromatographic evaluation of poly (trans-1,2-cyclohexanediy-bis acrylamide) as a chiral stationary phase for HPLC. <i>Journal of Chromatography A</i> , 2005, 1066, 55-70.	1.8	38
188	Enantiomeric separation of biaryl atropisomers using cyclofructan based chiral stationary phases. <i>Journal of Chromatography A</i> , 2014, 1357, 172-181.	1.8	38
189	Separation of chiral furan derivatives by liquid chromatography using cyclodextrin-based chiral stationary phases. <i>Journal of Chromatography A</i> , 2005, 1063, 111-120.	1.8	37
190	Characterization of new R-naphthylethyl cyclofructan 6 chiral stationary phase and its comparison with R-naphthylethyl β -cyclodextrin-based column. <i>Journal of Chromatography A</i> , 2011, 1218, 1393-1398.	1.8	37
191	Direct and sensitive determination of glyphosate and aminomethylphosphonic acid in environmental water samples by high performance liquid chromatography coupled to electrospray tandem mass spectrometry. <i>Journal of Chromatography A</i> , 2016, 1443, 93-100.	1.8	37
192	Fast super/subcritical fluid chromatographic enantioseparations on superficially porous particles bonded with broad selectivity chiral selectors relative to fully porous particles. <i>Journal of Chromatography A</i> , 2019, 1605, 360339.	1.8	37
193	Use of microcolumn liquid chromatography with a chiral stationary phase for the separation of low-resolution enantiomers. <i>Journal of Chromatography A</i> , 1987, 389, 256-260.	1.8	36
194	Characterization and utilization of a novel triflate ionic liquid stationary phase for use in comprehensive two-dimensional gas chromatography. <i>Journal of Separation Science</i> , 2008, 31, 3429-3436.	1.3	36
195	The use of cationic surfactants and ionic liquids in the detection of microbial contamination by capillary electrophoresis. <i>Electrophoresis</i> , 2008, 29, 2587-2592.	1.3	36
196	The Enantiomeric Separation of Tetrahydrobenzimidazoles Cyclodextrins and Cyclofructans. <i>Chirality</i> , 2013, 25, 133-140.	1.3	36
197	Cellular and cell-free studies of catalytic DNA cleavage by ruthenium polypyridyl complexes containing redox-active intercalating ligands. <i>Chemical Science</i> , 2017, 8, 3726-3740.	3.7	36
198	Examination of the Varied and Changing Ethanol Content of Commercial Kombucha Products. <i>Food Analytical Methods</i> , 2017, 10, 4062-4067.	1.3	36

#	ARTICLE	IF	CITATIONS
199	Variations of l- and d-amino acid levels in the brain of wild-type and mutant mice lacking d-amino acid oxidase activity. <i>Analytical and Bioanalytical Chemistry</i> , 2018, 410, 2971-2979.	1.9	36
200	Altered profiles and metabolism of l- and d-amino acids in cultured human breast cancer cells vs. non-tumorigenic human breast epithelial cells. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2019, 164, 421-429.	1.4	36
201	Enantiomeric resolution of racemic nicotine and nicotine analogues by microcolumn liquid chromatography with β -cyclodextrin inclusion complexes. <i>Journal of Chromatography A</i> , 1987, 411, 490-493.	1.8	35
202	Acylation effects on chiral recognition of racemic amines and alcohols by new polar and non-polar cyclodextrin derivative gas chromatographic phases. <i>Journal of Chromatography A</i> , 1990, 502, 154-159.	1.8	35
203	Chiral Separations Using Native and Functionalized Cyclodextrin-Bonded Stationary Phases in High-Pressure Liquid Chromatography. <i>ACS Symposium Series</i> , 1991, , 67-100.	0.5	35
204	Ionic liquid-alkane association in dilute solutions. <i>Theoretical Chemistry Accounts</i> , 2006, 117, 127-135.	0.5	35
205	Effect of silica gel modification with cyclofructans on properties of hydrophilic interaction liquid chromatography stationary phases. <i>Journal of Chromatography A</i> , 2012, 1257, 58-65.	1.8	35
206	On the use of ionic liquid capillary columns for analysis of aromatic hydrocarbons in low-boiling petrochemical products by one-dimensional and comprehensive two-dimensional gas chromatography. <i>Journal of Chromatography A</i> , 2013, 1301, 225-236.	1.8	35
207	On-line preconcentration of weak electrolytes by electrokinetic accumulation in CE: Experiment and simulation. <i>Electrophoresis</i> , 2007, 28, 1540-1547.	1.3	34
208	Carboxylated cyclofructan 6 as a hydrolytically stable high efficiency stationary phase for hydrophilic interaction liquid chromatography and mixed mode separations. <i>Analytical Methods</i> , 2016, 8, 6038-6045.	1.3	34
209	Evaluation of freeD-glutamate in processed foods. <i>Chirality</i> , 1994, 6, 277-282.	1.3	33
210	Avoparcin, a new macrocyclic antibiotic chiral run buffer additive for capillary electrophoresis. <i>Electrophoresis</i> , 1999, 20, 2438-2457.	1.3	33
211	High Efficiency Liquid and Supercritical Fluid-Based Enantiomeric Separations: An Overview. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2004, 27, 1121-1178.	0.5	33
212	Estimation of association constants between oral malodor components and various native and derivatized cyclodextrins. <i>Analytica Chimica Acta</i> , 2006, 557, 184-190.	2.6	33
213	Use of a polar ionic liquid as second column for the comprehensive two-dimensional GC separation of PCBs. <i>Journal of Chromatography A</i> , 2010, 1217, 5859-5867.	1.8	33
214	Mechanisms of ESI-MS Selectivity and Sensitivity Enhancements When Detecting Anions in the Positive Mode Using Cationic Pairing Agents. <i>Analytical Chemistry</i> , 2010, 82, 9066-9073.	3.2	33
215	Effect of Selector Coverage and Mobile Phase Composition on Enantiomeric Separations with Ristocetin A Chiral Stationary Phases. <i>Microchemical Journal</i> , 1999, 62, 26-49.	2.3	32
216	CE-ESI-MS analysis of singly charged inorganic and organic anions using a dicationic reagent as a complexing agent. <i>Electrophoresis</i> , 2009, 30, 3918-3925.	1.3	32

#	ARTICLE	IF	CITATIONS
217	High-performance liquid chromatography with paired ion electrospray ionization (PIESI) tandem mass spectrometry for the highly sensitive determination of acidic pesticides in water. <i>Analytica Chimica Acta</i> , 2013, 792, 1-9.	2.6	32
218	Thermodynamic studies of a zwitterionic stationary phase in hydrophilic interaction liquid chromatography. <i>Journal of Chromatography A</i> , 2013, 1272, 81-89.	1.8	32
219	Separation of nucleotides by hydrophilic interaction chromatography using the FRULIC-N column. <i>Analytical and Bioanalytical Chemistry</i> , 2013, 405, 8837-8848.	1.9	32
220	Evaluation of nicotine in tobacco-free nicotine commercial products. <i>Drug Testing and Analysis</i> , 2017, 9, 944-948.	1.6	32
221	Gram Scale Conversion of <i>R</i> -BINAM to <i>R</i> -NOBIN. <i>Journal of Organic Chemistry</i> , 2016, 81, 1295-1299.	1.7	31
222	Chiral surfaces: The many faces of chiral recognition. <i>Current Opinion in Colloid and Interface Science</i> , 2017, 32, 94-107.	3.4	31
223	Enantioselective UHPLC Screening Combined with <i>In Silico</i> Modeling for Streamlined Development of Ultrafast Enantiopurity Assays. <i>Analytical Chemistry</i> , 2022, 94, 1804-1812.	3.2	31
224	Centrifugal Partition Chromatography. IV. Preparative Sample Purification and Partition Coefficient Determination. <i>Journal of Liquid Chromatography and Related Technologies</i> , 1988, 11, 1187-1204.	0.9	30
225	Direct Measurement of Octanol-Water Partition Coefficients Using Centrifugal Partition Chromatography with a Back-Flushing Technique. <i>Journal of Liquid Chromatography and Related Technologies</i> , 1990, 13, 3061-3077.	0.9	30
226	Resolution of enantiomeric hydrocarbon biomarkers of geochemical importance. <i>Analytical Chemistry</i> , 1991, 63, 2858-2861.	3.2	30
227	In vitro study of the metabolic effects of D-amino acids. , 1996, 8, 24-29.		30
228	Retention and Selectivity of Teicoplanin Stationary Phases after Copper Complexation and Isotopic Exchange. <i>Analytical Chemistry</i> , 2001, 73, 5499-5508.	3.2	30
229	Cyclodextrin-Based Chiral Stationary Phases for Liquid Chromatography: A Twenty-Year Overview <i></i></i> . , 2004, 243, 061-112.		30
230	Resolution of Enantiomers in Solution and Determination of the Chirality of Extended Metal Atom Chains. <i>Inorganic Chemistry</i> , 2007, 46, 1535-1537.	1.9	30
231	Combined capillary electrophoresis and DNA fluorescence <i>in situ</i> hybridization for rapid molecular identification of <i>Salmonella</i> Typhimurium in mixed culture. <i>Electrophoresis</i> , 2008, 29, 2477-2484.	1.3	30
232	Enantiomeric separation of chiral ruthenium(II) complexes using capillary electrophoresis. <i>Chirality</i> , 2009, 21, 208-217.	1.3	30
233	Characterization of cyclodextrin-based chiral stationary phases by linear free energy relationship. <i>Journal of Separation Science</i> , 2011, 34, 2639-2644.	1.3	30
234	Study on the use of boromycin as a chiral selector in capillary electrophoresis. <i>Journal of Chromatography A</i> , 2012, 1237, 128-132.	1.8	30

#	ARTICLE	IF	CITATIONS
235	An insight into the use of dimethylphenyl carbamate cyclofructan 7 chiral stationary phase in supercritical fluid chromatography: The basic comparison with HPLC. <i>Journal of Separation Science</i> , 2013, 36, 1711-1719.	1.3	30
236	Mechanism and Sensitivity of Anion Detection Using Rationally Designed Unsymmetrical Dications in Paired Ion Electrospray Ionization Mass Spectrometry. <i>Analytical Chemistry</i> , 2014, 86, 2665-2672.	3.2	30
237	Thermal racemization of biaryl atropisomers. <i>Tetrahedron: Asymmetry</i> , 2017, 28, 1557-1561.	1.8	30
238	Linear Tricationic Room-Temperature Ionic Liquids: Synthesis, Physicochemical Properties, and Electrowetting Properties. <i>ACS Applied Materials & Interfaces</i> , 2009, 1, 2126-2133.	4.0	29
239	High-performance liquid chromatographic enantioseparation of monoterpene-based 2-amino carboxylic acids on macrocyclic glycopeptide-based phases. <i>Journal of Chromatography A</i> , 2010, 1217, 6956-6963.	1.8	29
240	Enantiomeric separations of illicit drugs and controlled substances using cyclofructan-based (LARIHC) and cyclobond I 2000 RSP HPLC chiral stationary phases. <i>Drug Testing and Analysis</i> , 2014, 6, 542-551.	1.6	29
241	Centrifugal Partition Chromatography. II. Selectivity Efficiency. <i>Journal of Liquid Chromatography and Related Technologies</i> , 1988, 11, 567-583.	0.9	28
242	Enantiomeric composition of monoterpenes in conifer resins. <i>Tetrahedron: Asymmetry</i> , 1997, 8, 3977-3984.	1.8	28
243	Influence of Chiral Ionic Liquids on Stereoselective Fluorescence Quenching by Photoinduced Electron Transfer in a Naproxen Dyad. <i>Journal of Physical Chemistry B</i> , 2009, 113, 10825-10829.	1.2	28
244	Use of ion pairing reagents for sensitive detection and separation of phospholipids in the positive ion mode LC-ESI-MS. <i>Analyst</i> , The, 2011, 136, 1586.	1.7	28
245	Enantioselective comprehensive two-dimensional gas chromatography. A route to elucidate the authenticity and origin of <i>Rosa damascena</i> Miller essential oils. <i>Journal of Separation Science</i> , 2015, 38, 3397-3403.	1.3	28
246	Dicationic ionic liquid thermal decomposition pathways. <i>Analytical and Bioanalytical Chemistry</i> , 2018, 410, 4645-4655.	1.9	28
247	Separation of Cyclodextrins Using Cyclodextrin Bonded Phases. <i>Journal of Liquid Chromatography and Related Technologies</i> , 1988, 11, 3295-3304.	0.9	27
248	HPLC Enantioseparation Of Di- and Tripeptides on Cyclodextrin Bonded Stationary Phases After Derivatization with 6-Aminoquinolyl-N-hydroxysuccinimidyl Carbamate (AQC). <i>Journal of Liquid Chromatography and Related Technologies</i> , 1994, 17, 483-497.	0.9	27
249	Direct chiral separation of unnatural amino acids by high-performance liquid chromatography on a ristocetin a-bonded stationary phase. <i>Chirality</i> , 2001, 13, 648-656.	1.3	27
250	Composition and chirality of amino acids in aerosol/dust from laboratory and residential enclosures. <i>Chirality</i> , 2001, 13, 153-158.	1.3	27
251	The determination of glutathione-4-hydroxynonenal (GSHNE), E-4-hydroxynonenal (HNE), and E-1-hydroxynon-2-en-4-one (HNO) in mouse liver tissue by LC-ESI-MS. <i>Analytical and Bioanalytical Chemistry</i> , 2008, 392, 1325-1333.	1.9	27
252	Separation of inorganic and small organic anions by CE using phosphonium-based mono- and dicationic reagents. <i>Electrophoresis</i> , 2009, 30, 3955-3963.	1.3	27

#	ARTICLE	IF	CITATIONS
253	Enantiomeric Separations of Ruthenium (II) Polypyridyl Complexes Using HPLC With Cyclofructan Chiral Stationary Phases. <i>Chirality</i> , 2015, 27, 64-70.	1.3	27
254	Water Determination in Solid Pharmaceutical Products Utilizing Ionic Liquids and Headspace Gas Chromatography. <i>Journal of Pharmaceutical Sciences</i> , 2016, 105, 2288-2292.	1.6	27
255	Enantiomeric Separations of Chiral Sulfonic and Phosphoric Acids with Barium-Doped Cyclofructan Selectors via an Ion Interaction Mechanism. <i>Analytical Chemistry</i> , 2014, 86, 1282-1290.	3.2	26
256	A comprehensive methodology for the chiral separation of 40 tobacco alkaloids and their carcinogenic Δ^2 -(R,S)-tobacco-specific nitrosamine metabolites. <i>Talanta</i> , 2018, 181, 132-141.	2.9	26
257	Direct Octanol Water Partition Coefficient Determination Using Co-Current Chromatography. <i>Journal of Liquid Chromatography and Related Technologies</i> , 1992, 15, 2769-2785.	0.9	25
258	Evaluation of the concentration and enantiomeric purity of selected free amino acids in fermented malt beverages (beers). , 1996, 8, 49-57.		25
259	RESOLUTION OF CHIRAL THIOL COMPOUNDS DERIVATIZED WITH N-(1-PYRENYL)-MALEIMIDE AND THIOLGLO Δ^3 . <i>Journal of Liquid Chromatography and Related Technologies</i> , 2000, 23, 1941-1952.	0.5	25
260	Separation of the enantiomers of substituted dihydrofurocoumarins by HPLC using macrocyclic glycopeptide chiral stationary phases. <i>Analytical and Bioanalytical Chemistry</i> , 2003, 377, 639-654.	1.9	25
261	Enantiomeric impurities in chiral synthons, catalysts, and auxiliaries: Part 3. <i>Tetrahedron: Asymmetry</i> , 2006, 17, 2821-2832.	1.8	25
262	Comparison of performance of Chirobiotic T, T2 and TAG columns in the separation of Δ^2 and Δ^3 homoamino acids. <i>Journal of Separation Science</i> , 2008, 31, 3688-3697.	1.3	25
263	GC-MS analysis of crocetane, phytane and some of their stereoisomers using cyclodextrin-based stationary phases. <i>Organic Geochemistry</i> , 2009, 40, 283-286.	0.9	25
264	Rapid identification of <i>Candida albicans</i> in blood by combined capillary electrophoresis and fluorescence <i>in situ</i> hybridization. <i>Electrophoresis</i> , 2010, 31, 2849-2853.	1.3	25
265	Problems and Pitfalls in the Analysis of Amygdalin and Its Epimer. <i>Journal of Agricultural and Food Chemistry</i> , 2015, 63, 8966-8973.	2.4	25
266	The utilisation of two detectors for the determination of water in honey using headspace gas chromatography. <i>Food Chemistry</i> , 2016, 205, 23-27.	4.2	25
267	Study of complexation between cyclofructans and alkali metal cations by electrospray ionization mass spectrometry and density functional theory calculations. <i>International Journal of Mass Spectrometry</i> , 2010, 291, 118-124.	0.7	24
268	Enantiomeric separations of Δ^1 -aryl ketones with cyclofructan chiral stationary phases via high performance liquid chromatography and supercritical fluid chromatography. <i>Journal of Chromatography A</i> , 2016, 1427, 45-54.	1.8	24
269	Variation of anionic moieties of dicationic ionic liquid GC stationary phases: Effect on stability and selectivity. <i>Analytica Chimica Acta</i> , 2018, 1042, 155-164.	2.6	24
270	Arsenic sequestration by iron oxide coated geopolymer microspheres. <i>Journal of Cleaner Production</i> , 2021, 291, 125931.	4.6	24

#	ARTICLE	IF	CITATIONS
271	Chromatography of Amino Acids on Reversed Phase Thin Layer Plates. <i>Journal of Liquid Chromatography and Related Technologies</i> , 1983, 6, 95-108.	0.9	23
272	Separation of carotenes on cyclodextrin-bonded phases. <i>Journal of Chromatography A</i> , 1990, 499, 627-635.	1.8	23
273	Positive mode electrospray ionization mass spectrometry of bisphosphonates using dicationic and tricationic ion-pairing agents. <i>Analytica Chimica Acta</i> , 2009, 633, 232-237.	2.6	23
274	High-performance liquid chromatographic enantioseparation of amino compounds on newly developed cyclofructan-based chiral stationary phases. <i>Journal of Separation Science</i> , 2012, 35, 617-624.	1.3	23
275	Isopropyl derivative of cyclofructan 6 as chiral selector in liquid chromatography and capillary electrophoresis. <i>Journal of Chromatography A</i> , 2014, 1338, 197-200.	1.8	23
276	Superficially Porous Particle Based Hydroxypropyl- β -cyclodextrin Stationary Phase for High-Efficiency Enantiomeric Separations. <i>Chirality</i> , 2015, 27, 788-794.	1.3	23
277	Mass spectrometric detection of trace anions: The evolution of paired-ion electrospray ionization (PIESI). <i>Mass Spectrometry Reviews</i> , 2016, 35, 201-218.	2.8	23
278	Examination of Selectivities of Thermally Stable Geminal Dicationic Ionic Liquids by Structural Modification. <i>Chromatographia</i> , 2017, 80, 1563-1574.	0.7	23
279	Direct high-performance liquid chromatographic enantioseparation of β -lactam stereoisomers. <i>Chirality</i> , 2005, 17, 193-200.	1.3	22
280	Fast detection of <i>Candida albicans</i> and/or bacteria in blood plasma by "sample-self-focusing" using capillary electrophoresis-laser-induced fluorescence. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2010, 53, 75-80.	1.4	22
281	A Tunable Ionic Liquid Based RC Filter Using Electrowetting: A New Concept. <i>ACS Applied Materials & Interfaces</i> , 2010, 2, 1785-1787.	4.0	22
282	High-Performance Liquid Chromatographic Resolution of Neutral and Cationic Hetero[6]Helicenes. <i>Chirality</i> , 2016, 28, 282-289.	1.3	22
283	Total peak shape analysis: detection and quantitation of concurrent fronting, tailing, and their effect on asymmetry measurements. <i>Journal of Chromatography A</i> , 2017, 1509, 163-170.	1.8	22
284	Branched-chain dicationic ionic liquids for fatty acid methyl ester assessment by gas chromatography. <i>Analytical and Bioanalytical Chemistry</i> , 2018, 410, 4633-4643.	1.9	22
285	Enantiomeric composition and prevalence of some bicyclic monoterpenoids in amber. <i>Chirality</i> , 1996, 8, 39-48.	1.3	21
286	Enantioseparation of chiral sulfoxides and sulfinate esters by capillary electrophoresis. <i>Electrophoresis</i> , 2002, 23, 1561.	1.3	21
287	Empirical Observations and Mechanistic Insights on the First Boron-Containing Chiral Selector for LC and Supercritical Fluid Chromatography. <i>Analytical Chemistry</i> , 2007, 79, 8125-8135.	3.2	21
288	Evaluation of tetracationic salts as gas-phase ion-pairing agents for the detection of trivalent anions in positive mode electrospray ionization mass spectrometry. <i>Rapid Communications in Mass Spectrometry</i> , 2010, 24, 1113-1123.	0.7	21

#	ARTICLE	IF	CITATIONS
289	Stereoselective separation of spiroindoline phytoalexins on R-naphthylethyl cyclofructan 6-based chiral stationary phase. <i>Journal of Chromatography A</i> , 2013, 1272, 100-105.	1.8	21
290	Separation of peptides on superficially porous particle based macrocyclic glycopeptide liquid chromatography stationary phases: consideration of fast separations. <i>Analytical and Bioanalytical Chemistry</i> , 2017, 409, 2437-2447.	1.9	21
291	The utility of statistical moments in chromatography using trapezoidal and Simpson's rules of peak integration. <i>Journal of Separation Science</i> , 2019, 42, 1644-1657.	1.3	21
292	Increasing chromatographic resolution of analytical signals using derivative enhancement approach. <i>Talanta</i> , 2019, 192, 492-499.	2.9	21
293	Mass Spectrometry-Compatible Enantiomeric Separations of 100 Pesticides Using Core-Shell Chiral Stationary Phases and Evaluation of Iterative Curve Fitting Models for Overlapping Peaks. <i>Chromatographia</i> , 2019, 82, 221-233.	0.7	21
294	Enhancing supercritical fluid chromatographic efficiency: Predicting effects of small aqueous additives. <i>Analytica Chimica Acta</i> , 2020, 1120, 75-84.	2.6	21
295	A Comparison of Polymer Separation Efficiency and Resolution by Gradient LC, GPC and TLC. <i>Journal of Liquid Chromatography and Related Technologies</i> , 1984, 7, 29-43.	0.9	20
296	Enantioseparations of chiral ruthenium(II) polypyridyl complexes using HPLC with macrocyclic glycopeptide chiral stationary phases (CSPs). <i>Journal of Molecular Structure</i> , 2008, 890, 75-80.	1.8	20
297	Sterility testing by CE: A comparison of online preconcentration approaches in capillaries with greater internal diameters. <i>Electrophoresis</i> , 2009, 30, 3870-3876.	1.3	20
298	The evaluation and comparison of trigonal and linear tricationic ion-pairing reagents for the detection of anions in positive mode ESI-MS. <i>Journal of the American Society for Mass Spectrometry</i> , 2009, 20, 529-538.	1.2	20
299	Complete identification of all 20 relevant epimeric peptides in \hat{I}^2 -amyloid: a new HPLC-MS based analytical strategy for Alzheimer's research. <i>Chemical Communications</i> , 2020, 56, 1537-1540.	2.2	20
300	Enantioseparation performance of superficially porous particle vancomycin-based chiral stationary phases in supercritical fluid chromatography and high performance liquid chromatography; applicability for psychoactive substances. <i>Journal of Chromatography A</i> , 2021, 1637, 461846.	1.8	20
301	Separation of Enantiomers Using a \hat{I}^3 -Cyclodextrin Liquid Chromatographic Bonded Phase. <i>Journal of Liquid Chromatography and Related Technologies</i> , 1990, 13, 473-484.	0.9	19
302	Influence of Chiral Ionic Liquids on the Excited-State Properties of Naproxen Analogs. <i>Journal of Physical Chemistry B</i> , 2008, 112, 7555-7559.	1.2	19
303	Enantioselective host-guest complexation of Ru(II) trisdiimine complexes using neutral and anionic derivatized cyclodextrins. <i>Inorganica Chimica Acta</i> , 2009, 362, 3073-3078.	1.2	19
304	4,6-Di-O-pentyl-3-O-trifluoroacetyl/propionyl cyclofructan stationary phases for gas chromatographic enantiomeric separations. <i>Analyst</i> , The, 2011, 136, 2931.	1.7	19
305	Determination of methionine enantiomers by HPLC on the cyclofructan chiral stationary phase. <i>Analytical Methods</i> , 2015, 7, 4577-4582.	1.3	19
306	High-performance liquid chromatographic separation of paclitaxel intermediate phenylisoserine derivatives on macrocyclic glycopeptide and cyclofructan-based chiral stationary phases. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2015, 114, 312-320.	1.4	19

#	ARTICLE	IF	CITATIONS
307	Quantitative analysis of dicamba residues in raw agricultural commodities with the use of ion-pairing reagents in LC-ESI-MS/MS. <i>Talanta</i> , 2016, 149, 103-109.	2.9	19
308	Determination of the enantiomeric purity of scopolamine isolated from plant extract using achiral/chiral coupled column chromatography. <i>Biomedical Chromatography</i> , 1991, 5, 3-7.	0.8	18
309	Synthesis and chromatographic evaluation of new polymeric chiral stationary phases based on three (1S,2S)-(âˆ™)-1,2-diphenylethylenediamine derivatives in HPLC and SFC. <i>Analytical and Bioanalytical Chemistry</i> , 2011, 399, 2445-2461.	1.9	18
310	Chiral HPLC Separation on Derivatized Cyclofructan Versus Cyclodextrin Stationary Phases. <i>Analytical Letters</i> , 2012, 45, 2344-2358.	1.0	18
311	Enantioseparation of citalopram analogues with sulfated β -cyclodextrin by capillary electrophoresis. <i>Electrophoresis</i> , 2016, 37, 841-848.	1.3	18
312	Geopolymers as a New Class of High pH Stable Supports with Different Chromatographic Selectivity. <i>Analytical Chemistry</i> , 2018, 90, 8139-8146.	3.2	18
313	Power Law Approach as a Convenient Protocol for Improving Peak Shapes and Recovering Areas from Partially Resolved Peaks. <i>Chromatographia</i> , 2019, 82, 211-220.	0.7	18
314	Enantioselective potential of teicoplanin- and vancomycin-based superficially porous particles-packed columns for supercritical fluid chromatography. <i>Journal of Chromatography A</i> , 2020, 1612, 460687.	1.8	18
315	Extraction of Nonionic Surfactants from Waste Water Using Centrifugal Partition Chromatography. <i>Journal of Liquid Chromatography and Related Technologies</i> , 1992, 15, 2909-2925.	0.9	17
316	Removal of Organic Compounds from Water via Cloud-Point Extraction with Permethyl Hydroxypropyl- β -cyclodextrin. <i>Separation Science and Technology</i> , 1993, 28, 1009-1018.	1.3	17
317	The Evolution of Chiral Stationary Phases for Liquid Chromatography. <i>Journal of the Chinese Chemical Society</i> , 1998, 45, 581-590.	0.8	17
318	Evaluation of the enantiomeric composition of amino acids in tobacco. , 1999, 11, 669-673.		17
319	LC and LC-MS Separation of Peptides on Macrocyclic Glycopeptide Stationary Phases: Diastereomeric Series and Large Peptides. <i>Chromatographia</i> , 2007, 66, 461-468.	0.7	17
320	Detection of nucleotides in positive-mode electrospray ionization mass spectrometry using multiply-charged cationic ion-pairing reagents. <i>Analytical and Bioanalytical Chemistry</i> , 2010, 398, 367-376.	1.9	17
321	Chromatographic separation of racemic praziquantel and its residual determination in perch by LC-MS/MS. <i>Talanta</i> , 2017, 174, 380-386.	2.9	17
322	Synthesis of new C3 symmetric amino acid- and aminoalcohol- containing chiral stationary phases and application to HPLC enantioseparations. <i>Chirality</i> , 2018, 30, 74-84.	1.3	17
323	The theory and practice of ultrafast liquid chromatography: A tutorial. <i>Analytica Chimica Acta</i> , 2021, 1151, 238170.	2.6	17
324	Enantiomeric separation of neutral hydrophobic dihydroflavones by cyclodextrin-modified micellar capillary electrophoresis. <i>Electrophoresis</i> , 2004, 25, 2727-2734.	1.3	16

#	ARTICLE	IF	CITATIONS
325	Enantioseparation of extended metal atom chain complexes: Unique compounds of extraordinarily high specific rotation. <i>Chirality</i> , 2007, 19, 179-183.	1.3	16
326	Comparison of stationary phases for packed column supercritical fluid chromatography based upon ionic liquid motifs: a study of cation and anion effects. <i>Analytical and Bioanalytical Chemistry</i> , 2011, 400, 435-447.	1.9	16
327	Rapid, effective deprotection of tert-butoxycarbonyl (Boc) amino acids and peptides at high temperatures using a thermally stable ionic liquid. <i>RSC Advances</i> , 2015, 5, 95854-95856.	1.7	16
328	Mass spectrometry detection of basic drugs in fast chiral analyses with vancomycin stationary phases. <i>Journal of Pharmaceutical Analysis</i> , 2018, 8, 324-332.	2.4	16
329	Centrifugal Partition Chromatography. III. Physico-Chemical Properties of Ternary Liquid Systems. <i>Journal of Liquid Chromatography and Related Technologies</i> , 1988, 11, 1171-1185.	0.9	15
330	Use of Centrifugal Partition Chromatography and Proteins in The Preparative Separation of Amino Acid Enantiomers. <i>Journal of Liquid Chromatography and Related Technologies</i> , 1990, 13, 3571-3581.	0.9	15
331	Determination of Association Constants in Cyclodextrin or Vancomycin-Modified Micellar Capillary Electrophoresis. <i>Journal of Liquid Chromatography and Related Technologies</i> , 1997, 20, 203-216.	0.5	15
332	Enantioseparation of dihydrofurocoumarin derivatives by various separation modes of capillary electrophoresis. <i>Electrophoresis</i> , 2003, 24, 2650-2656.	1.3	15
333	High-performance liquid chromatographic chiral separation of β -amino acids. <i>Chirality</i> , 2009, 21, 787-798.	1.3	15
334	Chiral Recognition with Macrocyclic Glycopeptides: Mechanisms and Applications. , 2010, , 203-222.		15
335	Sensitive analysis of metal cations in positive ion mode electrospray ionization mass spectrometry using commercial chelating agents and cationic ion-pairing reagents. <i>Rapid Communications in Mass Spectrometry</i> , 2012, 26, 1005-1013.	0.7	15
336	Separation of therapeutic peptides with cyclodextrin and glycopeptide based columns in hydrophilic interaction liquid chromatography. <i>Journal of Chromatography A</i> , 2015, 1390, 50-61.	1.8	15
337	Hydroxypropyl beta cyclodextrin bonded superficially porous particle-based HILIC stationary phases. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2016, 39, 459-464.	0.5	15
338	Screening primary racemic amines for enantioseparation by derivatized polysaccharide and cyclodextrin columns. <i>Journal of Pharmaceutical Analysis</i> , 2016, 6, 345-355.	2.4	15
339	Reduced matrix effects for anionic compounds with paired ion electrospray ionization mass spectrometry. <i>Analytica Chimica Acta</i> , 2016, 912, 74-84.	2.6	15
340	Liquid chromatographic enantioseparation of carbocyclic β -amino acids possessing limonene skeleton on macrocyclic glycopeptide-based chiral stationary phases. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2017, 145, 119-126.	1.4	15
341	Chiral Gas Chromatography. , 2018, , 468-505.		15
342	Gas chromatography selectivity of new phosphonium-based dicationic ionic liquid stationary phases. <i>Journal of Separation Science</i> , 2018, 41, 4142-4148.	1.3	15

#	ARTICLE	IF	CITATIONS
343	Selective separations of peptides with sequence deletions, single amino acid polymorphisms, and/or epimeric centers using macrocyclic glycopeptide liquid chromatography stationary phases. <i>Journal of Chromatography A</i> , 2004, 1053, 89-99.	1.8	15
344	Centrifugal Partition Chromatography. vi. Temperature Effects. <i>Journal of Liquid Chromatography and Related Technologies</i> , 1988, 11, 1457-1474.	0.9	14
345	Sensitive enantiomeric separation of aliphatic and aromatic amines using aromatic anhydrides as non-chiral derivatizing agents. <i>Journal of Chromatography A</i> , 1994, 666, 485-491.	1.8	14
346	Chiral separations of indan, tetralin and benzosuberan derivatives by capillary electrophoresis. <i>Journal of Chromatography A</i> , 1998, 793, 135-143.	1.8	14
347	Optimization of the Synthesis of 2,6-Dinitro-4-trifluoromethylphenyl Ether Substituted Cyclodextrin Bonded Chiral Stationary Phases. <i>Chromatographia</i> , 2006, 64, 147-155.	0.7	14
348	Preparation and evaluation of a new synthetic polymeric chiral stationary phase for HPLC based on the trans-9,10-dihydro-9,10-ethanoanthracene-(11S,12S)-11,12-dicarboxylic acid bis-4-vinylphenylamide monomer. <i>Analytical and Bioanalytical Chemistry</i> , 2007, 387, 2681-2697.	1.9	14
349	Comparison of HPLC enantioseparation of substituted binaphthyls on CD, polysaccharide and synthetic polymer based chiral stationary phases. <i>Journal of Separation Science</i> , 2010, 33, 1244-1254.	1.3	14
350	Preparation and Evaluation of HPLC Chiral Stationary Phases Based on Cationic/Basic Derivatives of Cyclofructan 6. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2015, 38, 550-560.	0.5	14
351	Enantioselective comprehensive two-dimensional gas chromatography of lavender essential oil. <i>Journal of Separation Science</i> , 2016, 39, 4765-4772.	1.3	14
352	Optical enrichment of dansyl-rac-amino acids by formation of crystalline inclusion complexes with cyclodextrins. <i>Chirality</i> , 1989, 1, 137-141.	1.3	13
353	Quantitative and stereoisomeric determination of light biomarkers in crude oil and coal samples. <i>Geochimica Et Cosmochimica Acta</i> , 1998, 62, 1619-1630.	1.6	13
354	Evaluation of non-polar interactions in chiral recognition by alkylated β - and γ -cyclodextrin chiral stationary phases. <i>Journal of Separation Science</i> , 2002, 25, 45-52.	1.3	13
355	Enantiomeric Separations by HPLC Using Macrocyclic Glycopeptide-Based Chiral Stationary Phases: An Overview, 2004, 243, 113-172.		13
356	Selectivity Tuning in an HPLC Multicomponent Separation. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2005, 28, 1453-1471.	0.5	13
357	Enantiomeric Separation of Synthetic Amino Acids Using Capillary Zone Electrophoresis. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2007, 30, 1421-1436.	0.5	13
358	LC Enantiomeric Separation of Unusual Amino Acids Using Cyclodextrin-based Stationary Phases. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2007, 31, 219-230.	0.5	13
359	Enantiomeric Resolution of a Chiral Sulfoxide Series by LC on Synthetic Polymeric Columns with Multimodal Elution. <i>Chromatographia</i> , 2010, 71, 361-372.	0.7	13
360	The enantiomeric separation of 4,5-disubstituted imidazoles by HPLC and CE using cyclodextrin-based chiral selectors. <i>Supramolecular Chemistry</i> , 2010, 22, 758-767.	1.5	13

#	ARTICLE	IF	CITATIONS
361	Metal cation detection in positive ion mode electrospray ionization mass spectrometry using a tetracationic salt as a gas-phase ion-pairing agent: Evaluation of the effect of chelating agents on detection sensitivity. <i>Rapid Communications in Mass Spectrometry</i> , 2012, 26, 2885-2896.	0.7	13
362	Complexation of cyclofructans with transition metal ions studied by electrospray ionization mass spectrometry and collision-induced dissociation. <i>International Journal of Mass Spectrometry</i> , 2012, 323-324, 21-27.	0.7	13
363	Enantiomeric impurities in chiral catalysts, auxiliaries, and synthons used in enantioselective syntheses. Part 4. <i>Tetrahedron: Asymmetry</i> , 2013, 24, 1134-1141.	1.8	13
364	Development and evaluation of gas and liquid chromatographic methods for the analysis of fatty amines. <i>Journal of Separation Science</i> , 2014, 37, 558-565.	1.3	13
365	Enantiomeric separation of citalopram analogues by HPLC using macrocyclic glycopeptide and cyclodextrin based chiral stationary phases. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2016, 39, 154-160.	0.5	13
366	Evaluation of the Edman degradation product of vancomycin bonded to core-shell particles as a new HPLC chiral stationary phase. <i>Chirality</i> , 2018, 30, 1067-1078.	1.3	13
367	A Gas Chromatography-Molecular Rotational Resonance Spectroscopy Based System of Singular Specificity. <i>Angewandte Chemie - International Edition</i> , 2020, 59, 192-196.	7.2	13
368	High information spectroscopic detection techniques for gas chromatography. <i>Journal of Chromatography A</i> , 2022, 1676, 463255.	1.8	13
369	Interconversion of Oxazepam Enantiomers During HPLC Separation. Determination of Thermodynamic Parameters. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2006, 29, 2889-2900.	0.5	12
370	Evaluation of dalbavancin as chiral selector for HPLC and comparison with teicoplanin-based chiral stationary phases. <i>Chirality</i> , 2010, 22, 495-513.	1.3	12
371	Ionic Liquid Based Headspace Solid-Phase Microextraction-Gas Chromatography for the Determination of Volatile Polar Organic Compounds. <i>Separation Science and Technology</i> , 2010, 45, 2322-2328.	1.3	12
372	Enantiomeric separation of isochromene derivatives by high-performance liquid chromatography using cyclodextrin based stationary phases and principal component analysis of the separation data. <i>Journal of Chromatography A</i> , 2013, 1305, 94-101.	1.8	12
373	Enantiomeric separation of functionalized ethano-bridged Tröger bases using macrocyclic cyclofructan and cyclodextrin chiral selectors in high-performance liquid chromatography and capillary electrophoresis with application of principal component analysis. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2014, 955-956, 72-80.	1.2	12
374	On the use of quadrupole mass spectrometric detection for flow modulated comprehensive two-dimensional gas chromatography. <i>Journal of Chromatography A</i> , 2014, 1330, 51-60.	1.8	12
375	Paired-ion electrospray ionization-triple quadrupole tandem mass spectrometry for quantification of anionic surfactants in waters. <i>Talanta</i> , 2015, 143, 320-327.	2.9	12
376	Topiramate: A Review of Analytical Approaches for the Drug Substance, Its Impurities and Pharmaceutical Formulations. <i>Journal of Chromatographic Science</i> , 2016, 54, bmv120.	0.7	12
377	Extending the power transform approach for recovering areas of overlapping peaks. <i>Journal of Separation Science</i> , 2019, 42, 3604-3610.	1.3	12
378	Roles of N-methyl-d-aspartate receptors and d-amino acids in cancer cell viability. <i>Molecular Biology Reports</i> , 2020, 47, 6749-6758.	1.0	12

#	ARTICLE	IF	CITATIONS
379	Selectivity of a Native β -Cyclodextrin Column in the Separation of Catechins. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2005, 28, 1669-1678.	0.5	11
380	Separation and sensitive determination of sphingolipids at low femtomole level by using HPLC-PIESI-MS/MS. <i>Analyst</i> , 2014, 139, 4169-4175.	1.7	11
381	Chlorinated aromatic derivatives of cyclofructan 6 as HPLC chiral stationary phases. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2016, 39, 497-503.	0.5	11
382	HPLC Enantioseparation of Novel Spirobrassinin Analogs on the Cyclofructan Chiral Stationary Phases. <i>Chromatographia</i> , 2017, 80, 53-62.	0.7	11
383	Enantiorecognition ability of different chiral selectors for separation of liquid crystals in supercritical fluid chromatography; critical evaluation. <i>Journal of Chromatography A</i> , 2020, 1622, 461138.	1.8	11
384	Calibration of GC-FID and IR Spectrometric Methods for Determination of High Boiling Petroleum Hydrocarbons in Environmental Samples. <i>Water, Air, and Soil Pollution</i> , 2004, 153, 329-341.	1.1	10
385	LC Separation of β -Amino Acid Enantiomers. <i>Chromatographia</i> , 2010, 71, 13-19.	0.7	10
386	Study of a new chiral selector: Sodium arsenyl-(l)-(+)-tartrate for capillary electrophoresis. <i>Journal of Chromatography A</i> , 2010, 1217, 1139-1148.	1.8	10
387	Rapid determination of sample purity and composition by nanopore stochastic sensing. <i>Nanoscale</i> , 2011, 3, 4593.	2.8	10
388	The empirical comparison of cyclofructans and cyclodextrins as chiral selectors in capillary electrophoretic separation of atropisomers of (R,S)-1,1'-binaphthalene-2,2'-diyl hydrogen phosphate. <i>Journal of Separation Science</i> , 2016, 39, 973-979.		10
389	Evaluation of gas chromatography for the separation of a broad range of isotopic compounds. <i>Analytica Chimica Acta</i> , 2021, 1165, 338490.	2.6	10
390	Interactions of chiral molecules with an (r)-n-(3,5-dinitrobenzoyl) phenylglycine HPLC stationary phase. <i>Chirality</i> , 1990, 2, 38-42.	1.3	9
391	Cyclodextrin PLOT columns for the gas-solid chromatographic separation of light hydrocarbons and inorganic gases. <i>Journal of Separation Science</i> , 1994, 6, 151-157.	1.0	9
392	Coupling solid-phase microextraction and laser desorption ionization for rapid identification of biological material. <i>Rapid Communications in Mass Spectrometry</i> , 2012, 26, 853-862.	0.7	9
393	On the biosynthesis and optical activity of the flinderoles. <i>Tetrahedron Letters</i> , 2013, 54, 5892-5894.	0.7	9
394	Separation of 2-naphthol atropisomers on cyclofructan-based chiral stationary phases. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2016, 39, 710-717.	0.5	9
395	Development and validation of a stability-indicating HPLC method for topiramate using a mixed-mode column and charged aerosol detector. <i>Journal of Separation Science</i> , 2018, 41, 1716-1725.	1.3	9
396	Cyclofructans as Chiral Selectors: An Overview. <i>Methods in Molecular Biology</i> , 2019, 1985, 183-200.	0.4	9

#	ARTICLE	IF	CITATIONS
397	Enantiomeric separation of quorum sensing autoinducer homoserine lactones using GC-MS and LC-MS. <i>Analytical and Bioanalytical Chemistry</i> , 2020, 412, 2927-2937.	1.9	9
398	Development and validation of a fast HPLC method for methyl dopa enantiomers using superficially porous particle based macrocyclic glycopeptide stationary phase. <i>Microchemical Journal</i> , 2021, 164, 105957.	2.3	9
399	Halocarbon separations on a new GSC-PLOT column. <i>Journal of Separation Science</i> , 1996, 8, 83-87.	1.0	8
400	Generation of Fluorescent Adducts of Malondialdehyde and Amino Acids: Toward an Understanding of Lipofuscin. <i>Photochemistry and Photobiology</i> , 2004, 79, 21-25.	1.3	8
401	Separations of Cycloinooligosaccharides via Hydrophilic Interaction Chromatography (HILIC) and Ligand-Exchange Chromatography. <i>Separation Science and Technology</i> , 2010, 45, 447-452.	1.3	8
402	Enantioseparation of Novel Amino Analogs of Indole Phytoalexins on Macrocyclic Glycopeptide-Based Chiral Stationary Phase. <i>Chromatographia</i> , 2011, 74, 751-757.	0.7	8
403	STEREOSELECTIVE HPLC DETERMINATION OF THYROXINE ENANTIOMERS IN PHARMACEUTICALS. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2011, 34, 2304-2314.	0.5	8
404	Topiramate: a review of analytical approaches for biological matrices. <i>Biomedical Chromatography</i> , 2015, 29, 1461-1472.	0.8	8
405	Sensitive detection of anionic metabolites of drugs by positive ion mode HPLC-PIESI-MS. <i>International Journal of Mass Spectrometry</i> , 2015, 389, 14-25.	0.7	8
406	Separation of Enantiomers of Selected Sulfur-Containing Amino Acids by Using Serially Coupled Achiral-Chiral Columns. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2015, 38, 789-794.	0.5	8
407	Fabrication of interconnected macroporosity in geopolymers via inverse suspension polymerization. <i>Journal of the American Ceramic Society</i> , 2019, 102, 4405-4409.	1.9	8
408	Sensitive detection of topiramate degradation products by high performance liquid chromatography/electrospray ionization mass spectrometry using ion-pairing reagents and polarity switching. <i>Rapid Communications in Mass Spectrometry</i> , 2019, 33, 116-124.	0.7	8
409	High efficiency functionalized hydrophilic cyclofructans as stationary phases in sub/supercritical fluid chromatography. <i>Talanta</i> , 2021, 232, 122308.	2.9	8
410	Cyclofructans, a New Class of Chiral Stationary Phases. , 2010, , 77-96.		8
411	An examination of the effects of water on normal phase enantioseparations. <i>Analytica Chimica Acta</i> , 2022, 1200, 339608.	2.6	8
412	Effect of the Orientation of Amide Linkage Groups on the Enantioselectivity of Two Related Synthetic Polymeric Chiral Stationary Phases. <i>Chromatographia</i> , 2008, 67, 199-210.	0.7	7
413	Enantiomeric Separation of Isochromene Derivatives by Cyclodextrin-Modified Micellar Capillary Electrophoresis. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2008, 31, 2035-2052.	0.5	7
414	A liquid drop RC filter apparatus for detection. <i>Analytical and Bioanalytical Chemistry</i> , 2011, 401, 2669-2678.	1.9	7

#	ARTICLE	IF	CITATIONS
415	On retentivity tuning by flow in the second column of different comprehensive two dimensional gas chromatographic configurations. <i>Journal of Chromatography A</i> , 2011, 1218, 3186-3189.	1.8	7
416	Enantioseparation of flinderoles and borreverines by HPLC on Chirobiotic V and V2 stationary phases and by CE using cyclodextrin selectors. <i>Analytical and Bioanalytical Chemistry</i> , 2013, 405, 9169-9177.	1.9	7
417	Absolute configuration of an axially chiral sulfonate determined from its optical rotatory dispersion, electronic circular dichroism, and vibrational circular dichroism spectra. <i>Chirality</i> , 2017, 29, 670-676.	1.3	7
418	Comparison of reversed-phase, anion-exchange, and hydrophilic interaction HPLC for the analysis of nucleotides involved in biological enzymatic pathways. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2019, 42, 184-193.	0.5	7
419	Improving peak capacities over 100 in less than 60 seconds: operating above normal peak capacity limits with signal processing. <i>Analytical and Bioanalytical Chemistry</i> , 2020, 412, 1925-1932.	1.9	7
420	A Closer Examination of 6-Aminoquinolyl-N-Hydroxysuccinimidyl Carbamate Amino Acid Derivatization in HPLC with Multiple Detection Modes. <i>Chromatographia</i> , 2021, 84, 719-727.	0.7	7
421	Enantioseparation of α -amino acids by liquid chromatography using core-shell chiral stationary phases based on teicoplanin and teicoplanin aglycone. <i>Journal of Chromatography A</i> , 2021, 1653, 462383.	1.8	7
422	Cyclodextrin-Mediated Enantiomeric Separation of Chiral Dihydrofuroflavones, a Class of Compounds with Promising Pharmacological Activity. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2005, 28, 169-186.	0.5	6
423	Evaluation of Pentaproline-Based Chiral Stationary Phase by LC. <i>Chromatographia</i> , 2008, 67, 13-32.	0.7	6
424	Separation of Enantiomers and Control of Elution Order of β -Lactams by GC Using Cyclodextrin-Based Chiral Stationary Phases. <i>Chromatographia</i> , 2009, 69, 331-337.	0.7	6
425	Water determination. , 2014, , 223-241.		6
426	Separation Of Methionine Enantiomers By Using Teicoplanin And Cyclofructan Columns. <i>Nova Biotechnologica Et Chimica</i> , 2015, 14, 1-11.	0.1	6
427	Enantiomeric impurities in chiral catalysts, auxiliaries, and synthons used in enantioselective syntheses. Part 5. <i>Chirality</i> , 2019, 31, 688-699.	1.3	6
428	Synthetic aluminosilicate based geopolymers – Second generation geopolymer HPLC stationary phases. <i>Analytica Chimica Acta</i> , 2019, 1081, 209-217.	2.6	6
429	Triple Helical Ir(ppy) ₃ Phenylene Cage Prepared by Diol-Mediated Benzannulation: Synthesis, Resolution, Absolute Stereochemistry and Photophysical Properties. <i>Chemistry - A European Journal</i> , 2019, 25, 8719-8724.	1.7	6
430	Chiral resolution and absolute configuration determination of new metal-based photodynamic therapy antitumor agents. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2021, 204, 114233.	1.4	6
431	Enantiomeric Separation of New Chiral Azole Compounds. <i>Molecules</i> , 2021, 26, 213.	1.7	6
432	Enhancing Sensitivity for High-Selectivity Gas Chromatography-Molecular Rotational Resonance Spectroscopy. <i>Analytical Chemistry</i> , 2021, 93, 15525-15533.	3.2	6

#	ARTICLE	IF	CITATIONS
433	Enantiomeric Separation of Fused Polycycles by HPLC with Cyclodextrin and Macrocyclic Glycopeptide Chiral Stationary Phases. <i>Separation Science and Technology</i> , 2005, 40, 2745-2759.	1.3	5
434	NATIVE/DERIVATIZED CYCLOFRUCTAN 6 BOUND TO RESINS VIA "CLICK" CHEMISTRY AS STATIONARY PHASES FOR ACHIRAL/CHIRAL SEPARATIONS. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2014, 37, 2302-2326.	0.5	5
435	Computerized optimization of flows and temperature gradient in flow modulated comprehensive two-dimensional gas chromatography. <i>Journal of Chromatography A</i> , 2014, 1349, 135-138.	1.8	5
436	Construction the switch binding pattern of cyclofructan 6. <i>Tetrahedron</i> , 2015, 71, 3447-3452.	1.0	5
437	Enantiomeric separation of new phytoalexin analogs with cyclofructan chiral stationary phases in normal-phase mode. <i>Journal of Separation Science</i> , 2016, 39, 3669-3676.	1.3	5
438	Sensitive analysis of N-blocked amino acids using high-performance liquid chromatography with paired ion electrospray ionization mass spectrometry. <i>Analytical and Bioanalytical Chemistry</i> , 2018, 410, 4725-4735.	1.9	5
439	Improved rate of substrate oxidation catalyzed by genetically-engineered myoglobin. <i>Archives of Biochemistry and Biophysics</i> , 2018, 639, 44-51.	1.4	5
440	Chiral Liquid Chromatography. , 2018, , 507-564.		5
441	Improving visualization of trace components for quantification using a power law based integration approach. <i>Journal of Chromatography A</i> , 2018, 1574, 1-8.	1.8	5
442	Selective Depletion of Chiral 4-Hydroxypraziquantel Metabolites in Three Types of Aquaculture Fish by LC-MS/MS. <i>Journal of Agricultural and Food Chemistry</i> , 2019, 67, 4098-4104.	2.4	5
443	A Gas Chromatography-Molecular Rotational Resonance Spectroscopy Based System of Singular Specificity. <i>Angewandte Chemie</i> , 2020, 132, 198-202.	1.6	5
444	Rapid and selective separation of amyloid beta from its stereoisomeric point mutations implicated in neurodegenerative Alzheimer's disease. <i>Analytica Chimica Acta</i> , 2021, 1163, 338506.	2.6	5
445	Gas Chromatography Columns Using Ionic Liquids as Stationary Phase. <i>Green Chemistry and Sustainable Technology</i> , 2020, , 131-165.	0.4	5
446	Selective separations of peptides with sequence deletions, single amino acid polymorphisms, and/or epimeric centers using macrocyclic glycopeptide liquid chromatography stationary phases. <i>Journal of Chromatography A</i> , 2004, 1053, 89-99.	1.8	5
447	Cyclodextrin Purification with Hollow Fibers. <i>Separation Science and Technology</i> , 1991, 26, 515-527.	1.3	4
448	CE-ESI-MS separation of divalent organic and inorganic anions using a tricationic complexing reagent. <i>Electrophoresis</i> , 2012, 33, 734-740.	1.3	4
449	Separation of multiply charged anions by capillary electrophoresis using alkyl phosphonium pairing agents. <i>Electrophoresis</i> , 2012, 33, 1153-1161.	1.3	4
450	Enantiomeric Separations of "H"/"Me Aziridines Utilizing GC and HPLC. <i>Chromatographia</i> , 2014, 77, 1607-1612.	0.7	4

#	ARTICLE	IF	CITATIONS
451	Cation-enhanced capillary electrophoresis separation of atropisomer anions. <i>Electrophoresis</i> , 2015, 36, 2859-2865.	1.3	4
452	Production of both <i>N</i> - and <i>D</i> -acetyl-homoserine lactones by <i>Burkholderia cepacia</i> and <i>Vibrio fischeri</i> . <i>MicrobiologyOpen</i> , 2021, 10, e1242.	1.2	4
453	Enhanced carboxypeptidase efficacies and differentiation of peptide epimers. <i>Analytical Biochemistry</i> , 2022, 642, 114451.	1.1	4
454	Macrocyclic glycopeptides- and derivatized cyclofructan-based chiral stationary phases for the enantioseparation of fluorinated α -phenylalanine analogs. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2022, 219, 114912.	1.4	4
455	Effect of Organic Cosolvents on Enantio-Enrichments via Cyclodextrin-Based Precipitations: An Examination of Production Efficiency. <i>Separation Science and Technology</i> , 1995, 30, 2259-2276.	1.3	3
456	STUDY OF MECHANISM OF ENANTIOSEPARATION. II. HPLC CHIRAL ANALYSIS OF ALKOXYSUBSTITUTED ESTERS OF PHENYL CARBAMIC ACID. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2002, 25, 1711-1720.	0.5	3
457	Binding characteristics of native cyclofructan 6 and its derivatives with metal ions. <i>Supramolecular Chemistry</i> , 2014, 26, 705-713.	1.5	3
458	Determination of the interconversion energy barrier of three novel pentahelicene derivative enantiomers by dynamic high resolution liquid chromatography. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2017, 1051, 60-67.	1.2	3
459	Water determination. , 2020, , 459-477.		3
460	Insights into enantioselective separations of ionic metal complexes by sub/supercritical fluid chromatography. <i>Analytica Chimica Acta</i> , 2022, 1228, 340156.	2.6	3
461	Reprint of: Enantiomeric separation of functionalized ethano-bridged Tröger bases using macrocyclic cyclofructan and cyclodextrin chiral selectors in high-performance liquid chromatography and capillary electrophoresis with application of principal component analysis. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2014, 968, 40-48.	1.2	2
462	Ionizable Cyclofructan 6-Based Stationary Phases for Hydrophilic Interaction Liquid Chromatography Using Superficially Porous Particles. <i>Chromatographia</i> , 2021, 84, 821-832.	0.7	2
463	Sixth International Symposium on chiral discrimination. , 1996, 8, 1.		2
464	Ionic liquids as gas chromatography stationary phases. , 2022, , 171-202.		2
465	Liquid chromatography enantiomeric separation of chiral ethanolamine substituted compounds. <i>Chirality</i> , 2022, , .	1.3	2
466	An Evaluation of the Differential Partitioning and Separation of C60 and C70 Fullerenes in a Biphasic System Using Centrifugal Partition Chromatography (CPC). <i>Journal of Liquid Chromatography and Related Technologies</i> , 1995, 18, 1019-1034.	0.9	1
467	Enhancing the selectivity of polar hydrophilic analytes with a low concentration of barium ions in the mobile phase using geopolymers and silica supports. <i>Talanta</i> , 2020, 207, 120339.	2.9	1
468	Headspace study of chiral interconversion of N-acetyl-homocysteine thiolactones. <i>Journal of Chromatography A</i> , 2021, 1653, 462381.	1.8	1

#	ARTICLE	IF	CITATIONS
469	Analysis of Enantiomeric Compounds Using Multidimensional Liquid Chromatography. , 0, , 319-344.		1
470	Future perspectives for ionic liquids. , 2022, , 369-394.		1
471	Kurt Mislow honorary issue of Chirality. Chirality, 1998, 10, 1-2.	1.3	0
472	The Separation of Hypericin's Enantiomers and Their Photophysics in Chiral Environments^{A¶}. Photochemistry and Photobiology, 2005, 81, 183-186.	1.3	0
473	Multidimensional Separations. Analytical Chemistry, 2014, 86, 11473-11473.	3.2	0
474	Enhanced Performance Separations: Smaller, Faster, More Complex Samples. Analytical Chemistry, 2016, 88, 5561-5561.	3.2	0
475	Extracting More Information Using Less (Sample, Time, Preparation...). Analytical Chemistry, 2018, 90, 6347-6347.	3.2	0