

Antonios Stavropoulos-Kalinoglou

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7462924/publications.pdf>

Version: 2024-02-01

52
papers

3,226
citations

136950

32
h-index

182427

51
g-index

52
all docs

52
docs citations

52
times ranked

3641
citing authors

#	ARTICLE	IF	CITATIONS
1	Smartphone pedometers in adults with asthma: a practical approach to physical activity assessment? A pilot validation study. <i>Journal of Asthma</i> , 2022, 59, 967-975.	1.7	2
2	Associations Between Erythrocyte Membrane Fatty Acid Compositions and Biomarkers of Vascular Health in Adults With Type 1 Diabetes With and Without Insulin Resistance: A Cross-Sectional Analysis. <i>Canadian Journal of Diabetes</i> , 2022, 46, 111-117.	0.8	3
3	Participation in physical activity decreased more in people with rheumatoid arthritis than the general population during the COVID-19 lockdown: a cross-sectional study. <i>Rheumatology International</i> , 2022, 42, 241-250.	3.0	7
4	The effects of training with high-speed interval running on muscle performance are modulated by slope. <i>Physiological Reports</i> , 2021, 9, e14656.	1.7	2
5	Position Statement on Exercise Dosage in Rheumatic and Musculoskeletal Diseases: The Role of the IMPACT-RMD Toolkit. <i>Mediterranean Journal of Rheumatology</i> , 2021, 32, 378.	0.8	10
6	Omega-3 polyunsaturated fatty acid supplementation versus placebo on vascular health, glycaemic control, and metabolic parameters in people with type 1 diabetes: a randomised controlled preliminary trial. <i>Cardiovascular Diabetology</i> , 2020, 19, 127.	6.8	20
7	ACSM Preparticipation Health Screening Guidelines: A UK University Cohort Perspective. <i>Medicine and Science in Sports and Exercise</i> , 2019, 51, 1047-1054.	0.4	7
8	Could IL-6 inhibition prevent exercise-induced fat loss in RA?. <i>Nature Reviews Rheumatology</i> , 2019, 15, 192-194.	8.0	1
9	Comparison of the effects of exercise and anti-TNF treatment on cardiovascular health in rheumatoid arthritis: results from two controlled trials. <i>Rheumatology International</i> , 2019, 39, 219-225.	3.0	19
10	Adiponectin, Resistin, and Visfatin in Childhood Obesity and Exercise. <i>Pediatric Exercise Science</i> , 2015, 27, 454-462.	1.0	24
11	The role of exercise in the management of rheumatoid arthritis. <i>Expert Review of Clinical Immunology</i> , 2015, 11, 1121-1130.	3.0	63
12	Cardiorespiratory fitness levels and their association with cardiovascular profile in patients with rheumatoid arthritis: a cross-sectional study. <i>Rheumatology</i> , 2015, 54, kev035.	1.9	36
13	Prolonged QTc interval predicts all-cause mortality in patients with rheumatoid arthritis: an association driven by high inflammatory burden. <i>Rheumatology</i> , 2014, 53, 131-137.	1.9	73
14	Three months of moderate-intensity exercise reduced plasma 3-nitrotyrosine in rheumatoid arthritis patients. <i>European Journal of Applied Physiology</i> , 2014, 114, 1483-1492.	2.5	34
15	Individualised exercise improves endothelial function in patients with rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2014, 73, 748-751.	0.9	92
16	Muscle wasting in rheumatoid arthritis: The role of oxidative stress. <i>World Journal of Rheumatology</i> , 2014, 4, 44.	0.5	5
17	Predictors of asymmetric dimethylarginine levels in patients with rheumatoid arthritis: the role of insulin resistance. <i>Scandinavian Journal of Rheumatology</i> , 2013, 42, 176-181.	1.1	19
18	Individualised aerobic and resistance exercise training improves cardiorespiratory fitness and reduces cardiovascular risk in patients with rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2013, 72, 1819-1825.	0.9	183

#	ARTICLE	IF	CITATIONS
19	Anti-tumour necrosis factor alpha therapy improves insulin sensitivity in normal-weight but not in obese patients with rheumatoid arthritis. <i>Arthritis Research and Therapy</i> , 2012, 14, R160.	3.5	52
20	Obesity and Arthritis. , 2012, , 355-380.		0
21	Inverted BMI rather than BMI is a better proxy for percentage of body fat. <i>Annals of Human Biology</i> , 2011, 38, 681-684.	1.0	29
22	Obesity in rheumatoid arthritis. <i>Rheumatology</i> , 2011, 50, 450-462.	1.9	173
23	Disease activity and low physical activity associate with number of hospital admissions and length of hospitalisation in patients with rheumatoid arthritis. <i>Arthritis Research and Therapy</i> , 2011, 13, R108.	3.5	45
24	Platelet function in rheumatoid arthritis: arthritic and cardiovascular implications. <i>Rheumatology International</i> , 2011, 31, 153-164.	3.0	134
25	Rheumatoid arthritis susceptibility genes associate with lipid levels in patients with rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2011, 70, 1025-1032.	0.9	51
26	Anti-TNF± therapy may lead to blood pressure reductions through improved endothelium-dependent microvascular function in patients with rheumatoid arthritis. <i>Journal of Human Hypertension</i> , 2011, 25, 699-702.	2.2	36
27	Mean platelet volume in patients with rheumatoid arthritis: the effect of anti-TNF-alpha therapy. <i>Rheumatology International</i> , 2010, 30, 1125-1129.	3.0	155
28	What predicts obesity in patients with rheumatoid arthritis? An investigation of the interactions between lifestyle and inflammation. <i>International Journal of Obesity</i> , 2010, 34, 295-301.	3.4	37
29	The Rationale for Comparative Studies of Accelerated Atherosclerosis in Rheumatic Diseases. <i>Current Vascular Pharmacology</i> , 2010, 8, 437-449.	1.7	90
30	Association of Mean Platelet Volume with Hypertension in Rheumatoid Arthritis. <i>Inflammation and Allergy: Drug Targets</i> , 2010, 9, 45-50.	1.8	66
31	Vascular Function and Inflammation in Rheumatoid Arthritis: the Role of Physical Activity~!2009-11-20~!2009-12-14~!2010-02-22~!. <i>Open Cardiovascular Medicine Journal</i> , 2010, 4, 89-96.	0.3	38
32	Target organ damage in patients with rheumatoid arthritis: The role of blood pressure and heart rate. <i>Atherosclerosis</i> , 2010, 209, 255-260.	0.8	30
33	Rheumatoid cachexia and cardiovascular disease. <i>Nature Reviews Rheumatology</i> , 2010, 6, 445-451.	8.0	133
34	The Rationale for Comparative Studies of Accelerated Atherosclerosis in Rheumatic Diseases. <i>Current Vascular Pharmacology</i> , 2010, 999, 1-13.	1.7	3
35	Association of physical inactivity with increased cardiovascular risk in patients with rheumatoid arthritis. <i>European Journal of Cardiovascular Prevention and Rehabilitation</i> , 2009, 16, 188-194.	2.8	141
36	Associations of obesity with modifiable risk factors for the development of cardiovascular disease in patients with rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2009, 68, 242-245.	0.9	76

#	ARTICLE	IF	CITATIONS
37	Underweight and obese states both associate with worse disease activity and physical function in patients with established rheumatoid arthritis. <i>Clinical Rheumatology</i> , 2009, 28, 439-444.	2.2	81
38	Lack of an Association of GNB3 C825T Polymorphism and Blood Pressure in Patients with Rheumatoid Arthritis. <i>Clinical and Experimental Hypertension</i> , 2009, 31, 428-439.	1.3	3
39	Association of interleukin-6 (IL-6)-174G/C gene polymorphism with cardiovascular disease in patients with rheumatoid arthritis: The role of obesity and smoking. <i>Atherosclerosis</i> , 2009, 204, 178-183.	0.8	85
40	Cigarette smoking associates with body weight and muscle mass of patients with rheumatoid arthritis: a cross-sectional, observational study. <i>Arthritis Research and Therapy</i> , 2008, 10, R59.	3.5	34
41	Cigarette smoking significantly increases basal metabolic rate in patients with rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2008, 67, 70-73.	0.9	67
42	Rheumatoid arthritis, cardiovascular disease and physical exercise: a systematic review. <i>Rheumatology</i> , 2008, 47, 239-248.	1.9	190
43	Polymorphisms of the Endothelin-1 Gene Associate with Hypertension in Patients with Rheumatoid Arthritis. <i>Endothelium: Journal of Endothelial Cell Research</i> , 2008, 15, 203-212.	1.7	53
44	Transforming growth factor- β 1 869T/C, but not interleukin-6 -174G/C, polymorphism associates with hypertension in rheumatoid arthritis. <i>Rheumatology</i> , 2008, 48, 113-118.	1.9	42
45	Serum uric acid is independently associated with hypertension in patients with rheumatoid arthritis. <i>Journal of Human Hypertension</i> , 2008, 22, 177-182.	2.2	35
46	Metabolism in Patients with Rheumatoid Arthritis: Resting Energy Expenditure, Physical Activity and Diet-Induced Thermogenesis. Invited Review. <i>Recent Patents on Endocrine, Metabolic & Immune Drug Discovery</i> , 2008, 2, 97-102.	0.6	3
47	Long-term exposure to medium-dose glucocorticoid therapy associates with hypertension in patients with rheumatoid arthritis. <i>Rheumatology</i> , 2008, 47, 72-75.	1.9	170
48	“Cardiovascular” Drugs in Rheumatoid Arthritis: Killing Two Birds with One Stone?. <i>Immunology, Endocrine and Metabolic Agents in Medicinal Chemistry</i> , 2008, 8, 259-274.	0.5	1
49	Prevalence and associations of hypertension and its control in patients with rheumatoid arthritis. <i>Rheumatology</i> , 2007, 46, 1477-1482.	1.9	250
50	Redefining overweight and obesity in rheumatoid arthritis patients. <i>Annals of the Rheumatic Diseases</i> , 2007, 66, 1316-1321.	0.9	141
51	New resting energy expenditure prediction equations for patients with rheumatoid arthritis. <i>Rheumatology</i> , 2007, 47, 500-506.	1.9	42
52	Blockade of tumour necrosis factor- α in rheumatoid arthritis: effects on components of rheumatoid cachexia. <i>Rheumatology</i> , 2007, 46, 1824-1827.	1.9	140