

# Tamer Akar

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7479832/publications.pdf>

Version: 2024-02-01

76  
papers

4,018  
citations

117571

34  
h-index

118793

62  
g-index

77  
all docs

77  
docs citations

77  
times ranked

3534  
citing authors

#	ARTICLE	IF	CITATIONS
1	Removal of lead and copper ions from aqueous solutions by bacterial strain isolated from soil. <i>Chemical Engineering Journal</i> , 2006, 115, 203-211.	6.6	315
2	Determination of the equilibrium, kinetic and thermodynamic parameters of adsorption of copper(II) ions onto seeds of. <i>Journal of Hazardous Materials</i> , 2005, 124, 200-208.	6.5	259
3	Equilibrium and kinetics of biosorption of lead(II) from aqueous solutions by <i>Cephalosporium aphidicola</i> . <i>Separation and Purification Technology</i> , 2006, 47, 105-112.	3.9	238
4	Biosorption characteristics of <i>Aspergillus flavus</i> biomass for removal of Pb(II) and Cu(II) ions from an aqueous solution. <i>Bioresource Technology</i> , 2006, 97, 1780-1787.	4.8	195
5	Biosorption performance of <i>Botrytis cinerea</i> fungal by-products for removal of Cd(II) and Cu(II) ions from aqueous solutions. <i>Minerals Engineering</i> , 2005, 18, 1099-1109.	1.8	139
6	Biosorption of a reactive textile dye from aqueous solutions utilizing an agro-waste. <i>Desalination</i> , 2009, 249, 757-761.	4.0	135
7	Biosorption kinetics and isotherm studies of Acid Red 57 by dried <i>Cephalosporium aphidicola</i> cells from aqueous solutions. <i>Biochemical Engineering Journal</i> , 2006, 31, 197-203.	1.8	131
8	Chromium(VI) biosorption characteristics of <i>Neurospora crassa</i> fungal biomass. <i>Minerals Engineering</i> , 2005, 18, 681-689.	1.8	130
9	Biosorption of a textile dye (Acid Blue 40) by cone biomass of <i>Thuja orientalis</i> : Estimation of equilibrium, thermodynamic and kinetic parameters. <i>Bioresource Technology</i> , 2008, 99, 3057-3065.	4.8	127
10	<i>Botrytis cinerea</i> as a new fungal biosorbent for removal of Pb(II) from aqueous solutions. <i>Biochemical Engineering Journal</i> , 2005, 25, 227-235.	1.8	126
11	Biosorption of Pb(II) and Cu(II) from aqueous solutions by pretreated biomass of <i>Neurospora crassa</i> . <i>Process Biochemistry</i> , 2005, 40, 3550-3558.	1.8	119
12	Biosorption of Pb(II) by industrial strain of <i>Saccharomyces cerevisiae</i> immobilized on the biomatrix of cone biomass of <i>Pinus nigra</i> : Equilibrium and mechanism analysis. <i>Chemical Engineering Journal</i> , 2007, 131, 293-300.	6.6	111
13	Enhanced biosorption of nickel(II) ions by silica-gel-immobilized waste biomass: Biosorption characteristics in batch and dynamic flow mode. <i>Journal of Hazardous Materials</i> , 2009, 163, 1134-1141.	6.5	107
14	Zn(II) biosorption properties of <i>Botrytis cinerea</i> biomass. <i>Journal of Hazardous Materials</i> , 2006, 131, 137-145.	6.5	101
15	An attractive agro-industrial by-product in environmental cleanup: Dye biosorption potential of untreated olive pomace. <i>Journal of Hazardous Materials</i> , 2009, 166, 1217-1225.	6.5	101
16	Removal of copper(II) ions from synthetic solution and real wastewater by the combined action of dried <i>Trametes versicolor</i> cells and montmorillonite. <i>Hydrometallurgy</i> , 2009, 97, 98-104.	1.8	94
17	Biosorption of Reactive Blue 49 dye under batch and continuous mode using a mixed biosorbent of macro-fungus <i>Agaricus bisporus</i> and <i>Thuja orientalis</i> cones. <i>Chemical Engineering Journal</i> , 2009, 148, 26-34.	6.6	88
18	Biosorption characteristics of <i>Bacillus</i> sp. ATS-2 immobilized in silica gel for removal of Pb(II). <i>Journal of Hazardous Materials</i> , 2006, 136, 317-323.	6.5	80

#	ARTICLE	IF	CITATIONS
19	Investigation of the biosorption characteristics of lead(II) ions onto <i>Symphoricarpus albus</i> : Batch and dynamic flow studies. <i>Journal of Hazardous Materials</i> , 2009, 165, 126-133.	6.5	69
20	Assessment of cationic dye biosorption characteristics of untreated and non-conventional biomass: <i>Pyracantha coccinea</i> berries. <i>Journal of Hazardous Materials</i> , 2009, 168, 1302-1309.	6.5	67
21	Study on the characterization of lead (II) biosorption by fungus <i>Aspergillus parasiticus</i> . <i>Applied Biochemistry and Biotechnology</i> , 2007, 136, 389-405.	1.4	65
22	Decolorization of Reactive Blue 49 contaminated solutions by <i>Capsicum annuum</i> seeds: Batch and continuous mode biosorption applications. <i>Chemical Engineering Journal</i> , 2011, 168, 125-133.	6.6	65
23	Biosorption applications of modified fungal biomass for decolorization of Reactive Red 2 contaminated solutions: Batch and dynamic flow mode studies. <i>Bioresource Technology</i> , 2010, 101, 7271-7277.	4.8	63
24	Biosorption potential of <i>Neurospora crassa</i> cells for decolorization of Acid Red 57 (AR57) dye. <i>Journal of Chemical Technology and Biotechnology</i> , 2006, 81, 1100-1106.	1.6	60
25	Assessment of the biosorption characteristics of a macro-fungus for the decolorization of Acid Red 44 (AR44) dye. <i>Journal of Hazardous Materials</i> , 2009, 171, 865-871.	6.5	55
26	Biosorption of lead(II) ions onto waste biomass of <i>Phaseolus vulgaris</i> L.: estimation of the equilibrium, kinetic and thermodynamic parameters. <i>Desalination</i> , 2009, 244, 188-198.	4.0	54
27	Biosorption performance of surface modified biomass obtained from <i>Pyracantha coccinea</i> for the decolorization of dye contaminated solutions. <i>Chemical Engineering Journal</i> , 2010, 160, 466-472.	6.6	52
28	Biosorption potential of the waste biomaterial obtained from <i>Cucumis melo</i> for the removal of Pb <sup>2+</sup> ions from aqueous media: Equilibrium, kinetic, thermodynamic and mechanism analysis. <i>Chemical Engineering Journal</i> , 2012, 185-186, 82-90.	6.6	52
29	Cadmium exposure from the cement dust emissions: A field study in a rural residence. <i>Chemosphere</i> , 2006, 63, 1546-1552.	4.2	49
30	Effects of chromium exposure from a cement factory. <i>Environmental Research</i> , 2003, 91, 113-118.	3.7	46
31	Chitosan-alunite composite: An effective dye remover with high sorption, regeneration and application potential. <i>Carbohydrate Polymers</i> , 2016, 143, 318-326.	5.1	46
32	Utilization of <i>Thamnidium elegans</i> fungal culture in environmental cleanup: A reactive dye biosorption study. <i>Ecological Engineering</i> , 2013, 58, 363-370.	1.6	41
33	Decolorization of a textile dye, reactive red 198 (rr198), by <i>Aspergillus parasiticus</i> fungal biosorbent. <i>Brazilian Journal of Chemical Engineering</i> , 2009, 26, 399-405.	0.7	40
34	On the utilization of a lignocellulosic waste as an excellent dye remover: Modification, characterization and mechanism analysis. <i>Chemical Engineering Journal</i> , 2013, 229, 257-266.	6.6	39
35	Nickel removal characteristics of an immobilized macro fungus: equilibrium, kinetic and mechanism analysis of the biosorption. <i>Journal of Chemical Technology and Biotechnology</i> , 2013, 88, 680-689.	1.6	36
36	Utilization of the <i>Phaseolus vulgaris</i> L. Waste biomass for decolorization of the textile dye Acid Red 57: determination of equilibrium, kinetic and thermodynamic parameters. <i>Journal of Environmental Science and Health - Part A Toxic/Hazardous Substances and Environmental Engineering</i> , 2007, 42, 591-600.	0.9	34

#	ARTICLE	IF	CITATIONS
37	Adsorption Potential of Lead(II) Ions from Aqueous Solutions onto Capsicum annum Seeds. Separation Science and Technology, 2007, 42, 137-151.	1.3	26
38	Microbial cells immobilized on natural biomatrix as a new potential ecofriendly biosorbent for the biotreatment of reactive dye contamination. Journal of Water Process Engineering, 2021, 39, 101731.	2.6	25
39	Improved biosorption potential of Thuja orientalis cone powder for the biosorptive removal of Basic Blue 9. Carbohydrate Polymers, 2013, 94, 400-408.	5.1	24
40	Effective biodecolorization potential of surface modified lignocellulosic industrial waste biomass. Chemical Engineering Journal, 2015, 259, 286-292.	6.6	22
41	Efficacy of green waste-derived biochar for lead removal from aqueous systems: Characterization, equilibrium, kinetic and application. Journal of Environmental Management, 2021, 289, 112490.	3.8	21
42	Nickel Exposure and its Effects. BioMetals, 2005, 18, 7-13.	1.8	20
43	Dithiocarbamated Symphoricarpos albus as a potential biosorbent for a reactive dye. Chemical Engineering Journal, 2012, 211-212, 442-452.	6.6	20
44	Chemical modification of a plant origin biomass using cationic surfactant ABDAC and the biosorptive decolorization of RR45 containing solutions. Colloids and Surfaces B: Biointerfaces, 2013, 101, 307-314.	2.5	20
45	Multivariate optimization of the decolorization process by surface modified biomaterial: Box-Behnken design and mechanism analysis. Environmental Science and Pollution Research, 2014, 21, 13055-13068.	2.7	19
46	Chitosan immobilization and Fe <sub>3</sub> O <sub>4</sub> functionalization of olive pomace: An ecofriendly and recyclable Pb <sup>2+</sup> biosorbent. Carbohydrate Polymers, 2021, 269, 118266.	5.1	19
47	Biosorption Potential of the Macrofungus Ganoderma carnosum for Removal of Lead(II) Ions from Aqueous Solutions. Journal of Environmental Science and Health - Part A Toxic/Hazardous Substances and Environmental Engineering, 2006, 41, 2587-2606.	0.9	18
48	Effective decolorization potential of Thamnidium elegans: Biosorption optimization, modelling, characterization and application studies. Chemical Engineering Journal, 2013, 221, 461-468.	6.6	18
49	Conversion of natural mineral to effective geosorbent by coating MnO <sub>2</sub> and its application potential for dye contaminated wastewaters. Journal of Cleaner Production, 2018, 189, 887-897.	4.6	18
50	Biosorption of Basic Blue 7 by fungal cells immobilized on the green-type biomatrix of Phragmites australis spongy tissue. International Journal of Phytoremediation, 2018, 20, 145-152.	1.7	18
51	From green biowaste to water treatment applications: Utilization of modified new biochar for the efficient removal of ciprofloxacin. Sustainable Chemistry and Pharmacy, 2021, 24, 100522.	1.6	18
52	Efficient biosorption of a reactive dye from contaminated media by <i>Neurospora sitophila</i> cells and <i>Zea mays</i> silk tissue biomass system. Journal of Chemical Technology and Biotechnology, 2011, 86, 1332-1341.	1.6	16
53	Batch and Dynamic Flow Biosorption Potential of <i>Agaricus bisporus</i> / <i>Thuja orientalis</i> Biomass Mixture for Decolorization of RR45 Dye. Industrial & Engineering Chemistry Research, 2008, 47, 9715-9723.	1.8	15
54	Removal of Pb <sup>2+</sup> ions from contaminated solutions by microbial composite: Combined action of a soilborne fungus <i>Mucor plumbeus</i> and alunite matrix. Chemical Engineering Journal, 2013, 215-216, 626-634.	6.6	12

#	ARTICLE	IF	CITATIONS
55	Biosorptive detoxification of zearalenone biotoxin by surface-modified renewable biomass: process dynamics and application. <i>Journal of the Science of Food and Agriculture</i> , 2019, 99, 1850-1861.	1.7	12
56	Biotransformation of racemic diisophorone by <i>Cephalosporium aphidicola</i> and <i>Neurospora crassa</i> . <i>Biotechnology Letters</i> , 2005, 27, 1007-1010.	1.1	10
57	Nickel and Cadmium Concentrations in Plasma and Na <sup>+</sup> /K <sup>+</sup> ATPase Activities in Erythrocyte Membranes of the People Exposed to Cement Dust Emissions. <i>Environmental Monitoring and Assessment</i> , 2005, 104, 437-444.	1.3	10
58	Treatment design and characteristics of a biosorptive decolorization process by a green type sorbent. <i>Journal of Cleaner Production</i> , 2016, 112, 4844-4853.	4.6	10
59	Removal of Cadmium and Manganese by an Ecofriendly Biomass. <i>Clean - Soil, Air, Water</i> , 2016, 44, 202-210.	0.7	9
60	Anionically reinforced hydrogel network entrapped fungal cells for retention of cadmium in the contaminated aquatic media. <i>Journal of Environmental Management</i> , 2017, 204, 583-593.	3.8	9
61	Process design and potential use of a regenerable biomagsorbent for effective decolorization process. <i>Journal of the Taiwan Institute of Chemical Engineers</i> , 2018, 93, 554-565.	2.7	9
62	The feasibility of <i>Thamnidium elegans</i> cells for color removal from real wastewater. <i>Chemical Engineering Research and Design</i> , 2017, 105, 316-325.	2.7	8
63	A passively immobilized novel biomagsorbent for the effective biosorptive treatment of dye contamination. <i>Environmental Science and Pollution Research</i> , 2019, 26, 25834-25843.	2.7	8
64	Parametric optimization of Cu(II) removal process by a metakaolin-based geopolymer: Batch and continuous process design. <i>Journal of Cleaner Production</i> , 2022, 366, 132819.	4.6	8
65	Design and modeling of the decolorization characteristics of a regenerable and eco-friendly geopolymer: Batch and dynamic flow mode treatment aspects. <i>Journal of Environmental Management</i> , 2021, 298, 113548.	3.8	7
66	Characterization and application of plant-based magnetic biomaterial for batch and fixed-bed mode sequestration of lead from synthetic and real effluents. <i>Ecological Engineering</i> , 2013, 61, 251-257.	1.6	6
67	<i>Phlebia gigantea</i> cells immobilized on renewable biomass matrix as potential ecofriendly scavenger for lead contamination. <i>Environmental Science and Pollution Research</i> , 2020, 27, 16177-16188.	2.7	6
68	Immobilized <i>Mucor plumbeus</i> on sepiolite support: A potential decolorization agent suitable for batch and continuous mode water treatment. <i>Journal of Cleaner Production</i> , 2021, 294, 126283.	4.6	6
69	Attached culture of <i>Gibberella fujikuroi</i> for biocomposite sorbent production and ciprofloxacin sequestration applications. <i>Journal of Chemical Technology and Biotechnology</i> , 2021, 96, 2610-2619.	1.6	6
70	Biosorption potential of surface-modified waste sugar beet pulp for the removal of Reactive Yellow 2 (RY2) anionic dye. <i>Turkish Journal of Chemistry</i> , 2016, 40, 1044-1054.	0.5	4
71	Adsorption of Diazo Dye from Aqueous Solutions by Magnetic Montmorillonite Composite. <i>Clean - Soil, Air, Water</i> , 2021, 49, 2000165.	0.7	4
72	Surface-modified scarlet firethorn: an eco-friendly and effective dye remover with excellent regeneration potential. <i>Desalination and Water Treatment</i> , 2016, 57, 5546-5553.	1.0	3

#	ARTICLE	IF	CITATIONS
73	Magnetically functionalized alunite as a recyclable and ecofriendly adsorbent for efficient removal of Pb <sup>2+</sup> . <i>Journal of Water Process Engineering</i> , 2022, 48, 102867.	2.6	3
74	Synthesis and Evaluation of Demethoxyviridin Derivatives as Potential Antimicrobials. <i>Zeitschrift Fur Naturforschung - Section C Journal of Biosciences</i> , 2005, 60, 686-692.	0.6	1
75	Gel-entrapped biomass of <i>Lactarius salmonicolor</i> for the effective treatment of aquatic Co <sup>2+</sup> and Mn <sup>2+</sup> pollution. <i>Biomass Conversion and Biorefinery</i> , 2024, 14, 4257-4271.	2.9	0
76	Highly effective fish scale-based biochar as a recyclable and green material for toxic cadmium ion bioremoval from liquid phase. <i>Biomass Conversion and Biorefinery</i> , 2024, 14, 4869-4882.	2.9	0