

# Yanmin Hu

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7482642/publications.pdf>

Version: 2024-02-01

59  
papers

3,667  
citations

159585

30  
h-index

138484

58  
g-index

61  
all docs

61  
docs citations

61  
times ranked

5510  
citing authors

#	ARTICLE	IF	CITATIONS
1	The Efficacy of Using Combination Therapy against Multi-Drug and Extensively Drug-Resistant <i>Pseudomonas aeruginosa</i> in Clinical Settings. <i>Antibiotics</i> , 2022, 11, 323.	3.7	10
2	Mefloquine enhances the activity of colistin against antibiotic-resistant Enterobacterales in vitro and in an in vivo animal study. <i>International Journal of Antimicrobial Agents</i> , 2021, 57, 106309.	2.5	3
3	Zidovudine enhances activity of carbapenems against NDM-1-producing Enterobacteriaceae. <i>Journal of Antimicrobial Chemotherapy</i> , 2021, 76, 2302-2305.	3.0	5
4	Antibiotic combination therapy against resistant bacterial infections: synergy, rejuvenation and resistance reduction. <i>Expert Review of Anti-Infective Therapy</i> , 2020, 18, 5-15.	4.4	101
5	A model-based analysis identifies differences in phenotypic resistance between in vitro and in vivo: implications for translational medicine within tuberculosis. <i>Journal of Pharmacokinetics and Pharmacodynamics</i> , 2020, 47, 421-430.	1.8	3
6	Effect of Different Media on the Bactericidal Activity of Colistin and on the Synergistic Combination With Azidothymidine Against mcr-1-Positive Colistin-Resistant <i>Escherichia coli</i> . <i>Frontiers in Microbiology</i> , 2020, 11, 54.	3.5	15
7	Translational Model-Informed Approach for Selection of Tuberculosis Drug Combination Regimens in Early Clinical Development. <i>Clinical Pharmacology and Therapeutics</i> , 2020, 108, 274-286.	4.7	12
8	Urinary bactericidal activity of colistin and azidothymidine combinations against mcr-1-positive colistin-resistant <i>Escherichia coli</i> . <i>International Journal of Antimicrobial Agents</i> , 2019, 54, 55-61.	2.5	10
9	Synergistic activity of colistin with azidothymidine against colistin-resistant <i>Klebsiella pneumoniae</i> clinical isolates collected from inpatients in Greek hospitals. <i>International Journal of Antimicrobial Agents</i> , 2019, 53, 855-858.	2.5	17
10	Bedaquiline kills persistent <i>Mycobacterium tuberculosis</i> with no disease relapse: an in vivo model of a potential cure. <i>Journal of Antimicrobial Chemotherapy</i> , 2019, 74, 1627-1633.	3.0	19
11	Azidothymidine Produces Synergistic Activity in Combination with Colistin against Antibiotic-Resistant <i>Enterobacteriaceae</i> . <i>Antimicrobial Agents and Chemotherapy</i> , 2019, 63, .	3.2	35
12	Moxifloxacin Replacement in Contemporary Tuberculosis Drug Regimens Is Ineffective against Persistent <i>Mycobacterium tuberculosis</i> in the Cornell Mouse Model. <i>Antimicrobial Agents and Chemotherapy</i> , 2018, 62, .	3.2	4
13	Optimal doses of rifampicin in the standard drug regimen to shorten tuberculosis treatment duration and reduce relapse by eradicating persistent bacteria. <i>Journal of Antimicrobial Chemotherapy</i> , 2018, 73, 724-731.	3.0	17
14	A Method to Evaluate Persistent <i>Mycobacterium tuberculosis</i> In Vitro and in the Cornell Mouse Model of Tuberculosis. <i>Methods in Molecular Biology</i> , 2018, 1736, 157-166.	0.9	4
15	Forecasting Clinical Dose-Response From Preclinical Studies in Tuberculosis Research: Translational Predictions With Rifampicin. <i>Clinical Pharmacology and Therapeutics</i> , 2018, 104, 1208-1218.	4.7	22
16	Serum bactericidal activity of colistin and azidothymidine combinations against mcr-1-positive colistin-resistant <i>Escherichia coli</i> . <i>International Journal of Antimicrobial Agents</i> , 2018, 52, 783-789.	2.5	20
17	A Novel erm (44) Gene Variant from a Human <i>Staphylococcus saprophyticus</i> Isolate Confers Resistance to Macrolides and Lincosamides but Not Streptogramins. <i>Antimicrobial Agents and Chemotherapy</i> , 2017, 61, .	3.2	7
18	Investigation of Elimination Rate, Persistent Subpopulation Removal, and Relapse Rates of <i>Mycobacterium tuberculosis</i> by Using Combinations of First-Line Drugs in a Modified Cornell Mouse Model. <i>Antimicrobial Agents and Chemotherapy</i> , 2016, 60, 4778-4785.	3.2	19

#	ARTICLE	IF	CITATIONS
19	Defining dormancy in mycobacterial disease. <i>Tuberculosis</i> , 2016, 99, 131-142.	1.9	66
20	A multistate tuberculosis pharmacometric model: a framework for studying anti-tubercular drug effects <i>in vitro</i> . <i>Journal of Antimicrobial Chemotherapy</i> , 2016, 71, 964-974.	3.0	42
21	High-dose rifampicin kills persisters, shortens treatment duration, and reduces relapse rate <i>in vitro</i> and <i>in vivo</i> . <i>Frontiers in Microbiology</i> , 2015, 6, 641.	3.5	95
22	Nordihydroguaiaretic acid enhances the activities of aminoglycosides against methicillin-sensitive and resistant <i>Staphylococcus aureus</i> <i>in vitro</i> and <i>in vivo</i> . <i>Frontiers in Microbiology</i> , 2015, 6, 1195.	3.5	17
23	Same Exposure but Two Radically Different Responses to Antibiotics: Resilience of the Salivary Microbiome versus Long-Term Microbial Shifts in Feces. <i>MBio</i> , 2015, 6, e01693-15.	4.1	333
24	HspX knock-out in <i>Mycobacterium tuberculosis</i> leads to shorter antibiotic treatment and lower relapse rate in a mouse model – A potential novel therapeutic target. <i>Tuberculosis</i> , 2015, 95, 31-36.	1.9	12
25	Antimicrobial Peptide Novicidin Synergizes with Rifampin, Ceftriaxone, and Ceftazidime against Antibiotic-Resistant Enterobacteriaceae <i>In Vitro</i> . <i>Antimicrobial Agents and Chemotherapy</i> , 2015, 59, 6233-6240.	3.2	47
26	Combinations of $\beta$ -Lactam or Aminoglycoside Antibiotics with Plectasin Are Synergistic against Methicillin-Sensitive and Methicillin-Resistant <i>Staphylococcus aureus</i> . <i>PLoS ONE</i> , 2015, 10, e0117664.	2.5	40
27	Antimicrobial resistance characteristics and fitness of Gram-negative fecal bacteria from volunteers treated with minocycline or amoxicillin. <i>Frontiers in Microbiology</i> , 2014, 5, 722.	3.5	31
28	Identification of the monocyte activating motif in <i>Mycobacterium tuberculosis</i> chaperonin 60.1. <i>Tuberculosis</i> , 2013, 93, 442-447.	1.9	8
29	Can We Prevent Antimicrobial Resistance by Using Antimicrobials Better?. <i>Pathogens</i> , 2013, 2, 422-435.	2.8	27
30	Tuberculous Endocarditis. <i>International Journal of Cardiology</i> , 2013, 167, 640-645.	1.7	35
31	Enhancement by novel anti-methicillin-resistant <i>Staphylococcus aureus</i> compound HT61 of the activity of neomycin, gentamicin, mupirocin and chlorhexidine: <i>in vitro</i> and <i>in vivo</i> studies. <i>Journal of Antimicrobial Chemotherapy</i> , 2013, 68, 374-384.	3.0	35
32	Contradictory Results with High-Dosage Rifamycin in Mice and Humans. <i>Antimicrobial Agents and Chemotherapy</i> , 2013, 57, 1103-1103.	3.2	10
33	Nonmultiplying Bacteria are Profoundly Tolerant to Antibiotics. <i>Handbook of Experimental Pharmacology</i> , 2012, , 99-119.	1.8	25
34	Sudden cardiac death and tuberculosis – How much do we know?. <i>Tuberculosis</i> , 2012, 92, 307-313.	1.9	37
35	Novel classes of antibiotics or more of the same?. <i>British Journal of Pharmacology</i> , 2011, 163, 184-194.	5.4	452
36	<i>Mycobacterium tuberculosis</i> acg Gene Is Required for Growth and Virulence <i>In Vivo</i> . <i>PLoS ONE</i> , 2011, 6, e20958.	2.5	30

#	ARTICLE	IF	CITATIONS
37	3-Ketosteroid 9 $\alpha$ -hydroxylase is an essential factor in the pathogenesis of <i>Mycobacterium tuberculosis</i> . <i>Molecular Microbiology</i> , 2010, 75, 107-121.	2.5	113
38	A New Approach for the Discovery of Antibiotics by Targeting Non-Multiplying Bacteria: A Novel Topical Antibiotic for Staphylococcal Infections. <i>PLoS ONE</i> , 2010, 5, e11818.	2.5	63
39	Comparison of the Moonlighting Actions of the Two Highly Homologous Chaperonin 60 Proteins of <i>Mycobacterium tuberculosis</i> . <i>Infection and Immunity</i> , 2010, 78, 3196-3206.	2.2	50
40	Acute and Persistent <i>Mycobacterium tuberculosis</i> Infections Depend on the Thiol Peroxidase TPX. <i>PLoS ONE</i> , 2009, 4, e5150.	2.5	62
41	<i>Mycobacterial Heat Shock Protein 60s in the Induction and Regulation of Infectious Disease</i> . <i>Heat Shock Proteins</i> , 2009, , 121-133.	0.2	0
42	A model of catheter-associated urinary tract infection initiated by bacterial contamination of the catheter tip. <i>BJU International</i> , 2008, 102, 67-74.	2.5	37
43	Targeting non-multiplying organisms as a way to develop novel antimicrobials. <i>Trends in Pharmacological Sciences</i> , 2008, 29, 143-150.	8.7	69
44	A Biphasic Response From Bladder Epithelial Cells Induced by Catheter Material and Bacteria: An In Vitro Study of the Pathophysiology of Catheter Related Urinary Tract Infection. <i>Journal of Urology</i> , 2008, 180, 1522-1526.	0.4	11
45	<i>Mycobacterium tuberculosis</i> Mutant Lacking the <i>groEL</i> Homologue <i>cpn60.1</i> Is Viable but Fails To Induce an Inflammatory Response in Animal Models of Infection. <i>Infection and Immunity</i> , 2008, 76, 1535-1546.	2.2	100
46	Chaperonin 60 and Macrophage Activation. <i>Novartis Foundation Symposium</i> , 2008, 291, 160-172.	1.1	2
47	New Strategies for Antibacterial Drug Design. <i>Drugs in R and D</i> , 2006, 7, 133-151.	2.2	28
48	Deletion of the <i>Mycobacterium tuberculosis</i> $\hat{1}$ -Crystallin-Like <i>hspX</i> Gene Causes Increased Bacterial Growth In Vivo. <i>Infection and Immunity</i> , 2006, 74, 861-868.	2.2	127
49	Transposon mutagenesis identifies genes which control antimicrobial drug tolerance in stationary-phase <i>Escherichia coli</i> . <i>FEMS Microbiology Letters</i> , 2005, 243, 117-124.	1.8	64
50	The <i>Mycobacterium tuberculosis sigJ</i> gene controls sensitivity of the bacterium to hydrogen peroxide. <i>FEMS Microbiology Letters</i> , 2004, 237, 415-423.	1.8	45
51	The gene controls sensitivity of the bacterium to hydrogen peroxide. <i>FEMS Microbiology Letters</i> , 2004, 237, 415-423.	1.8	39
52	Sterilizing Activities of Fluoroquinolones against Rifampin-Tolerant Populations of <i>Mycobacterium tuberculosis</i> . <i>Antimicrobial Agents and Chemotherapy</i> , 2003, 47, 653-657.	3.2	196
53	The future challenges facing the development of new antimicrobial drugs. <i>Nature Reviews Drug Discovery</i> , 2002, 1, 895-910.	46.4	525
54	Increased levels of <i>sigJ</i> mRNA in late stationary phase cultures of <i>Mycobacterium tuberculosis</i> detected by DNA array hybridisation. <i>FEMS Microbiology Letters</i> , 2001, 202, 59-65.	1.8	57

#	ARTICLE	IF	CITATIONS
55	Increased levels of sigJ mRNA in late stationary phase cultures of Mycobacterium tuberculosis detected by DNA array hybridisation. FEMS Microbiology Letters, 2001, 202, 59-65.	1.8	3
56	Detection of mRNA Transcripts and Active Transcription in Persistent Mycobacterium tuberculosis Induced by Exposure to Rifampin or Pyrazinamide. Journal of Bacteriology, 2000, 182, 6358-6365.	2.2	168
57	Regulation of <i>hmp</i> Gene Transcription in <i>Mycobacterium tuberculosis</i> : Effects of Oxygen Limitation and Nitrosative and Oxidative Stress. Journal of Bacteriology, 1999, 181, 3486-3493.	2.2	79
58	Transcription of Two Sigma 70 Homologue Genes, <i>sigA</i> and <i>sigB</i> , in Stationary-Phase <i>Mycobacterium tuberculosis</i> . Journal of Bacteriology, 1999, 181, 469-476.	2.2	104
59	Transcription of the Stationary-Phase-Associated hspX Gene of Mycobacterium tuberculosis Is Inversely Related to Synthesis of the 16-Kilodalton Protein. Journal of Bacteriology, 1999, 181, 1380-1387.	2.2	56