

Rolf Kiessling

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7498413/publications.pdf>

Version: 2024-02-01

93
papers

8,299
citations

57758

44
h-index

49909

87
g-index

96
all docs

96
docs citations

96
times ranked

10605
citing authors

#	ARTICLE	IF	CITATIONS
1	Generation of Tumor-Specific Cytotoxic T Cells From Blood via InÂVtro Expansion Using Autologous Dendritic Cells Pulsed With Neoantigen-Coupled Microbeads. <i>Frontiers in Oncology</i> , 2022, 12, 866763.	2.8	2
2	Precision radiation of immune checkpoint therapy resistant melanoma metastases (PROMMEL study): study protocol for a phase II open-label multicenter trial. <i>Acta OncolÃ³gica</i> , 2022, 61, 869-873.	1.8	1
3	Targeting of Nrf2 improves antitumoral responses by human NK cells, TIL and CAR T cells during oxidative stress. , 2022, 10, e004458.		18
4	Trogocytosis and fratricide killing impede MSLN-directed CAR T cell functionality. <i>Oncolimmunology</i> , 2022, 11, .	4.6	9
5	Counteracting CAR T cell dysfunction. <i>Oncogene</i> , 2021, 40, 421-435.	5.9	76
6	Cisplatin inhibits frequency and suppressive activity of monocytic myeloid-derived suppressor cells in cancer patients. <i>Oncolimmunology</i> , 2021, 10, 1935557.	4.6	17
7	Predicting anti-PD-1 responders in malignant melanoma from the frequency of S100A9+ monocytes in the blood. , 2021, 9, e002171.		12
8	Interleukinâ€³3 is a Novel Immunosuppressor that Protects Cancer Cells from TIL Killing by a Macrophageâ€³Mediated Shedding Mechanism. <i>Advanced Science</i> , 2021, 8, 2101029.	11.2	20
9	Complete and long-lasting clinical responses in immune checkpoint inhibitor-resistant, metastasized melanoma treated with adoptive T cell transfer combined with DC vaccination. <i>Oncolimmunology</i> , 2020, 9, 1792058.	4.6	30
10	Visualization of human T lymphocyte-mediated eradication of cancer cells in vivo. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2020, 117, 22910-22919.	7.1	32
11	PD-1 checkpoint blockade in advanced melanoma patients: NK cells, monocytic subsets and host PD-L1 expression as predictive biomarker candidates. <i>Oncolimmunology</i> , 2020, 9, 1786888.	4.6	29
12	Caveolin-1-Mediated Tumor Suppression Is Linked to Reduced HIF1Î± S-Nitrosylation and Transcriptional Activity in Hypoxia. <i>Cancers</i> , 2020, 12, 2349.	3.7	11
13	Genetically modified immune cells targeting tumor antigens. , 2020, 214, 107603.		17
14	High expression of ID1 in monocytes is strongly associated with phenotypic and functional MDSC markers in advanced melanoma. <i>Cancer Immunology, Immunotherapy</i> , 2020, 69, 513-522.	4.2	6
15	The Outcome of <i>Ex Vivo</i> TIL Expansion Is Highly Influenced by Spatial Heterogeneity of the Tumor T-Cell Repertoire and Differences in Intrinsic <i>In Vitro</i> Growth Capacity between T-Cell Clones. <i>Clinical Cancer Research</i> , 2020, 26, 4289-4301.	7.0	46
16	Targeting a scavenger receptor on tumor-associated macrophages activates tumor cell killing by natural killer cells. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2020, 117, 32005-32016.	7.1	89
17	Establishment of Melanoma Tumor Xenograft Using Single Cell Line Suspension and Co-injection of Patient-Derived T Cells in Immune-Deficient NSG Mice. <i>Methods in Molecular Biology</i> , 2019, 1913, 207-215.	0.9	1
18	Assessment of Antitumor T-Cell Responses by Flow Cytometry After Coculture of Tumor Cells with Autologous Tumor-Infiltrating Lymphocytes. <i>Methods in Molecular Biology</i> , 2019, 1913, 133-140.	0.9	3

#	ARTICLE	IF	CITATIONS
19	Evaluating Antibody-Dependent Cell-Mediated Cytotoxicity by Chromium Release Assay. <i>Methods in Molecular Biology</i> , 2019, 1913, 167-179.	0.9	1
20	Evaluating Antibody-Dependent Cell-Mediated Cytotoxicity by Flow Cytometry. <i>Methods in Molecular Biology</i> , 2019, 1913, 181-194.	0.9	5
21	Cancer Neopeptides for Immunotherapy: Discordance Between Tumor-Infiltrating T Cell Reactivity and Tumor MHC Peptidome Display. <i>Frontiers in Immunology</i> , 2019, 10, 2766.	4.8	23
22	Self-Delivering RNAi Targeting PD-1 Improves Tumor-Specific T Cell Functionality for Adoptive Cell Therapy of Malignant Melanoma. <i>Molecular Therapy</i> , 2018, 26, 1482-1493.	8.2	38
23	Cripto-1 Plasmid DNA Vaccination Targets Metastasis and Cancer Stem Cells in Murine Mammary Carcinoma. <i>Cancer Immunology Research</i> , 2018, 6, 1417-1425.	3.4	25
24	Enhanced stimulation of human tumor-specific T cells by dendritic cells matured in the presence of interferon- β and multiple toll-like receptor agonists. <i>Cancer Immunology, Immunotherapy</i> , 2017, 66, 1333-1344.	4.2	31
25	IL-15, TIM-3 and NK cells subsets predict responsiveness to anti-CTLA-4 treatment in melanoma patients. <i>Oncolmmunology</i> , 2017, 6, e1261242.	4.6	59
26	Intratumorally injected pro-inflammatory allogeneic dendritic cells as immune enhancers: a first-in-human study in unfavourable risk patients with metastatic renal cell carcinoma. , 2017, 5, 52.		42
27	Ipilimumab treatment decreases monocytic MDSCs and increases CD8 effector memory T cells in long-term survivors with advanced melanoma. <i>Oncotarget</i> , 2017, 8, 21539-21553.	1.8	103
28	Regulation of myeloid cells by activated T cells determines the efficacy of PD-1 blockade. <i>Oncolmmunology</i> , 2016, 5, e1232222.	4.6	48
29	IL-15 activates mTOR and primes stress-activated gene expression leading to prolonged antitumor capacity of NK cells. <i>Blood</i> , 2016, 128, 1475-1489.	1.4	136
30	Coexpressed Catalase Protects Chimeric Antigen Receptor-Redirected T Cells as well as Bystander Cells from Oxidative Stress-Induced Loss of Antitumor Activity. <i>Journal of Immunology</i> , 2016, 196, 759-766.	0.8	164
31	Targeting Suppressive Myeloid Cells Potentiates Checkpoint Inhibitors to Control Spontaneous Neuroblastoma. <i>Clinical Cancer Research</i> , 2016, 22, 3849-3859.	7.0	109
32	Non-classical HLA-class I expression in serous ovarian carcinoma: Correlation with the HLA-genotype, tumor infiltrating immune cells and prognosis. <i>Oncolmmunology</i> , 2016, 5, e1052213.	4.6	51
33	Hypoxia-mediated alterations and their role in the HER-2/neuregulated CREB status and localization. <i>Oncotarget</i> , 2016, 7, 52061-52084.	1.8	11
34	Dendritic cell regulation of NK cell responses involves lymphotoxin-1, IL-12, and TGF- β 2. <i>European Journal of Immunology</i> , 2015, 45, 1783-1793.	2.9	34
35	Methylcholanthrene-Induced Sarcomas Develop Independently from NOX2-Derived ROS. <i>PLoS ONE</i> , 2015, 10, e0129786.	2.5	11
36	Contrasting Effects of the Cytotoxic Anticancer Drug Gemcitabine and the EGFR Tyrosine Kinase Inhibitor Gefitinib on NK Cell-Mediated Cytotoxicity via Regulation of NKG2D Ligand in Non-Small-Cell Lung Cancer Cells. <i>PLoS ONE</i> , 2015, 10, e0139809.	2.5	26

#	ARTICLE	IF	CITATIONS
37	Consensus nomenclature for CD8 ⁺ T cell phenotypes in cancer. <i>Oncolmmunology</i> , 2015, 4, e998538.	4.6	119
38	Checkpoint blockade for cancer therapy: revitalizing a suppressed immune system. <i>Trends in Molecular Medicine</i> , 2015, 21, 482-491.	6.7	146
39	T Cell Blockade Immunotherapy Against Cancer and Abscopal Effect in Combination Therapy. <i>Cancer Drug Discovery and Development</i> , 2015, , 211-229.	0.4	0
40	Classification of current anticancer immunotherapies. <i>Oncotarget</i> , 2014, 5, 12472-12508.	1.8	395
41	Laminins 411 and 421 differentially promote tumor cell migration via $\alpha 6 \beta 1$ integrin and MCAM (CD146). <i>Matrix Biology</i> , 2014, 38, 69-83.	3.6	53
42	A phase I clinical trial combining dendritic cell vaccination with adoptive T cell transfer in patients with stage IV melanoma. <i>Cancer Immunology, Immunotherapy</i> , 2014, 63, 1061-1071.	4.2	68
43	Myeloid-derived suppressor cells and their role in CTLA-4 blockade therapy. <i>Cancer Immunology, Immunotherapy</i> , 2014, 63, 977-983.	4.2	31
44	Myeloid Suppressors Decrease Melanoma Survival by Abating Tumor-Fighting T Cells. <i>Clinical Cancer Research</i> , 2014, 20, 1401-1403.	7.0	3
45	Inhibition of Tumor-Derived Prostaglandin-E2 Blocks the Induction of Myeloid-Derived Suppressor Cells and Recovers Natural Killer Cell Activity. <i>Clinical Cancer Research</i> , 2014, 20, 4096-4106.	7.0	230
46	Intratumoral vaccination with activated allogeneic dendritic cells in patients with newly diagnosed metastatic renal cell carcinoma (mRCC).. <i>Journal of Clinical Oncology</i> , 2014, 32, 3085-3085.	1.6	5
47	The two sides of HER2/neu: immune escape versus surveillance. <i>Trends in Molecular Medicine</i> , 2013, 19, 677-684.	6.7	17
48	Cyclooxygenase-2. <i>Oncolmmunology</i> , 2013, 2, e25157.	4.6	1
49	Ipilimumab Treatment Results in an Early Decrease in the Frequency of Circulating Granulocytic Myeloid-Derived Suppressor Cells as well as Their Arginase1 Production. <i>Cancer Immunology Research</i> , 2013, 1, 158-162.	3.4	112
50	The MAPK Pathway Is a Predominant Regulator of HLA-A Expression in Esophageal and Gastric Cancer. <i>Journal of Immunology</i> , 2013, 191, 6261-6272.	0.8	79
51	Melanoma-Educated CD14 ⁺ Cells Acquire a Myeloid-Derived Suppressor Cell Phenotype through COX-2-Dependent Mechanisms. <i>Cancer Research</i> , 2013, 73, 3877-3887.	0.9	160
52	NF- κ B activation during intradermal DNA vaccination is essential for eliciting tumor protective antigen-specific CTL responses. <i>Human Vaccines and Immunotherapeutics</i> , 2013, 9, 2189-2195.	3.3	15
53	Expression of MHC Class I on breast cancer cells correlates inversely with HER2 expression. <i>Oncolmmunology</i> , 2012, 1, 1104-1110.	4.6	64
54	Opposing consequences of signaling through EGF family members. <i>Oncolmmunology</i> , 2012, 1, 1200-1201.	4.6	2

#	ARTICLE	IF	CITATIONS
55	HER-2/neu-mediated Down-regulation of Biglycan Associated with Altered Growth Properties. <i>Journal of Biological Chemistry</i> , 2012, 287, 24320-24329.	3.4	28
56	On the armament and appearances of human myeloid-derived suppressor cells. <i>Clinical Immunology</i> , 2012, 144, 250-268.	3.2	168
57	HER2/HER3 Signaling Regulates NK Cell-Mediated Cytotoxicity via MHC Class I Chain-Related Molecule A and B Expression in Human Breast Cancer Cell Lines. <i>Journal of Immunology</i> , 2012, 188, 2136-2145.	0.8	51
58	Camouflage and sabotage: tumor escape from the immune system. <i>Cancer Immunology, Immunotherapy</i> , 2011, 60, 1161-1171.	4.2	150
59	T cell recognition of HLA-A2 restricted tumor antigens is impaired by the oncogene HER2. <i>International Journal of Cancer</i> , 2011, 128, 390-401.	5.1	53
60	Designer lymphocytes to fight cancer: a helping hand from modern molecular biology. <i>Journal of Molecular Medicine</i> , 2010, 88, 1081-1084.	3.9	1
61	Expression and prognostic significance of iNOS in uveal melanoma. <i>International Journal of Cancer</i> , 2010, 126, 2682-2689.	5.1	28
62	Immature Immunosuppressive CD14+HLA-DR ^{low} Cells in Melanoma Patients Are Stat3hi and Overexpress CD80, CD83, and DC-Sign. <i>Cancer Research</i> , 2010, 70, 4335-4345.	0.9	366
63	Antibody-Dependent Natural Killer Cell-Mediated Cytotoxicity Engendered by a Kinase-Inactive Human HER2 Adenovirus-Based Vaccination Mediates Resistance to Breast Tumors. <i>Cancer Research</i> , 2010, 70, 7431-7441.	0.9	24
64	Vaccination with a plasmid DNA encoding HER-2/neu together with low doses of GM-CSF and IL-2 in patients with metastatic breast carcinoma: a pilot clinical trial. <i>Journal of Translational Medicine</i> , 2010, 8, 53.	4.4	104
65	HER-2/neu mediated down-regulation of MHC class I antigen processing prevents CTL-mediated tumor recognition upon DNA vaccination in HLA-A2 transgenic mice. <i>Cancer Immunology, Immunotherapy</i> , 2009, 58, 653-664.	4.2	48
66	Prognostic significance of tumor iNOS and COX-2 in stage-III malignant cutaneous melanoma. <i>Cancer Immunology, Immunotherapy</i> , 2009, 58, 1085-1094.	4.2	44
67	Response: Resistance of naturally occurring regulatory T cells toward oxidative stress: possible link with intracellular catecholamine content and implications for cancer therapy. <i>Blood</i> , 2009, 114, 488-489.	1.4	1
68	Transduction with the Antioxidant Enzyme Catalase Protects Human T Cells against Oxidative Stress. <i>Journal of Immunology</i> , 2008, 181, 8382-8390.	0.8	81
69	The CD16 ⁺ CD56 ^{bright} NK Cell Subset Is Resistant to Reactive Oxygen Species Produced by Activated Granulocytes and Has Higher Antioxidative Capacity Than the CD16 ⁺ CD56 ^{dim} Subset. <i>Journal of Immunology</i> , 2007, 179, 4513-4519.	0.8	73
70	Interferon- β renders tumors that express low levels of Her-2/neu sensitive to cytotoxic T cells. <i>Cancer Immunology, Immunotherapy</i> , 2006, 55, 653-662.	4.2	8
71	Small interfering RNA (siRNA) inhibits the expression of the Her2/neu gene, upregulates HLA class I and induces apoptosis of Her2/neu positive tumor cell lines. <i>International Journal of Cancer</i> , 2004, 108, 71-77.	5.1	138
72	Cellular immunity to the Her-2/neu protooncogene. <i>Advances in Cancer Research</i> , 2002, 85, 101-144.	5.0	72

#	ARTICLE	IF	CITATIONS
73	Tumor necrosis factor- α induces coordinated changes in major histocompatibility class I presentation pathway, resulting in increased stability of class I complexes at the cell surface. <i>Blood</i> , 2001, 98, 1108-1115.	1.4	102
74	The identification of a common pathogen-specific HLA class II A*0201-restricted cytotoxic T cell epitope encoded within the heat shock protein 65. <i>European Journal of Immunology</i> , 2001, 31, 3602-3611.	2.9	26
75	Inhibition of Activated/Memory (CD45RO+) T Cells by Oxidative Stress Associated with Block of NF- κ B Activation. <i>Journal of Immunology</i> , 2001, 167, 2595-2601.	0.8	121
76	DNA Immunization of HLA Transgenic Mice with a Plasmid Expressing Mycobacterial Heat Shock Protein 65 Results in HLA Class I- and II-Restricted T Cell Responses That Can Be Augmented by Cytokines. <i>Human Gene Therapy</i> , 2001, 12, 1797-1804.	2.7	19
77	HER-2/neu is expressed in human renal cell carcinoma at heterogeneous levels independently of tumor grading and staging and can be recognized by HLA-A2.1-restricted cytotoxic T lymphocytes. <i>International Journal of Cancer</i> , 2000, 87, 349-359.	5.1	57
78	Tumor-induced immune dysfunction. <i>Cancer Immunology, Immunotherapy</i> , 1999, 48, 353-362.	4.2	208
79	Identification of HER2/neu-derived peptide epitopes recognized by gastric cancer-specific cytotoxic T lymphocytes. <i>International Journal of Cancer</i> , 1998, 78, 202-208.	5.1	75
80	CD28 is not required for rejection of unmanipulated syngeneic and autologous tumors. <i>European Journal of Immunology</i> , 1997, 27, 1988-1993.	2.9	17
81	Mechanisms of escape from CD8+ T-cell clones specific for the HER-2/NEU proto-oncogene expressed in ovarian carcinomas: Related and unrelated to decreased MHC class I expression. , 1997, 70, 112-119.		27
82	Hydrogen peroxide secreted by tumor-derived macrophages downmodulates signal-transducing zeta molecules and inhibits tumor-specific T cell and natural killer cell-mediated cytotoxicity. <i>European Journal of Immunology</i> , 1996, 26, 1308-1313.	2.9	321
83	Immunosuppression in human tumor-host interaction: role of cytokines and alterations in signal-transducing molecules. <i>Seminars in Immunopathology</i> , 1996, 18, 227-242.	4.0	36
84	Alterations in the signal-transducing molecules of T cells and nk cells in colorectal tumor-infiltrating, gut mucosal and peripheral lymphocytes: Correlation with the stage of the disease. <i>International Journal of Cancer</i> , 1995, 61, 765-772.	5.1	191
85	Lack of interleukin-2 (IL-2) expression and selective expression of IL-10 mRNA in human renal cell carcinoma. <i>International Journal of Cancer</i> , 1995, 63, 366-371.	5.1	125
86	The epstein-barr virus latent membrane protein-1 (LMP1) induces interleukin-10 production in burkitt lymphoma lines. <i>International Journal of Cancer</i> , 1994, 57, 240-244.	5.1	132
87	T cell receptor diversity and activation markers in the V β 1 subset of rheumatoid synovial fluid and peripheral blood T lymphocytes. <i>European Journal of Immunology</i> , 1992, 22, 567-574.	2.9	51
88	Role of hsp60 during Autoimmune and Bacterial Inflammation. <i>Immunological Reviews</i> , 1991, 121, 91-111.	6.0	110
89	Increased susceptibility of ifn- γ -treated neuroblastoma cells to lysis by lymphokine-activated killer cells: Participation of ICAM-1 induction on target cells. <i>International Journal of Cancer</i> , 1991, 47, 527-532.	5.1	58
90	Effect of IFN- γ treatment and in vivo passage of murine tumor cell lines on their sensitivity to lymphokine-activated killer (LAK) cell lysis in vitro; association with H-2 expression on the target cells. <i>International Journal of Cancer</i> , 1989, 44, 669-674.	5.1	17

#	ARTICLE	IF	CITATIONS
91	Selective rejection of H ² -deficient lymphoma variants suggests alternative immune defence strategy. Nature, 1986, 319, 675-678.	27.8	1,914
92	Gamma-interferon (IFN- γ) produced during effector and target interactions renders target cells less susceptible to NK-cell-mediated lysis. International Journal of Cancer, 1983, 32, 609-616.	5.1	30
93	An Analysis of the Murine NK Cell as to Structure, Function and Biological Relevance. Immunological Reviews, 1979, 44, 165-208.	6.0	349