
Walter Glen Thomas

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/7508851/publications.pdf Version: 2024-02-01



| # | Article | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | Akt acts as a switch for GPCR transactivation of the TGFâ $\in \hat{I}^2$ receptor type 1. FEBS Journal, 2022, 289, 2642-2656. | 4.7 | 6 |
| 2 | Stimulation of the four isoforms of receptor tyrosine kinase ErbB4, but not ErbB1, confers cardiomyocyte hypertrophy. Journal of Cellular Physiology, 2021, 236, 8160-8170. | 4.1 | 4 |
| 3 | Complex interactions between the angiotensin II type 1 receptor, the epidermal growth factor receptor and TRIO-dependent signaling partners. Biochemical Pharmacology, 2021, 188, 114521. | 4.4 | 2 |
| 4 | Type I Diabetes Mellitus Increases the Cardiovascular Complications of Influenza Virus Infection. Frontiers in Cellular and Infection Microbiology, 2021, 11, 714440. | 3.9 | 3 |
| 5 | Modular transient nanoclustering of activated β2-adrenergic receptors revealed by single-molecule tracking of conformation-specific nanobodies. Proceedings of the National Academy of Sciences of the United States of America, 2020, 117, 30476-30487. | 7.1 | 29 |
| 6 | A Bitter Taste in Your Heart. Frontiers in Physiology, 2020, 11, 431. | 2.8 | 31 |
| 7 | A High-Fat Diet Increases Influenza A Virus-Associated Cardiovascular Damage. Journal of Infectious Diseases, 2020, 222, 820-831. | 4.0 | 21 |
| 8 | Mutations in the NPxxY motif stabilize pharmacologically distinct conformational states of the α _{1B} - and β ₂ -adrenoceptors. Science Signaling, 2019, 12, . | 3.6 | 14 |
| 9 | BRET-based assay to monitor EGFR transactivation by the AT1R reveals Gq/11 protein-independent activation and AT1R-EGFR complexes. Biochemical Pharmacology, 2018, 158, 232-242. | 4.4 | 19 |
| 10 | CRIM1 is necessary for coronary vascular endothelial cell development and homeostasis. Journal of Molecular Histology, 2017, 48, 53-61. | 2.2 | 10 |
| 11 | Cavin-1 deficiency modifies myocardial and coronary function, stretch responses and ischaemic tolerance: roles of NOS over-activity. Basic Research in Cardiology, 2017, 112, 24. | 5.9 | 15 |
| 12 | Transactivation of the epidermal growth factor receptor in responses to myocardial stress and cardioprotection. International Journal of Biochemistry and Cell Biology, 2017, 83, 97-110. | 2.8 | 24 |
| 13 | Functional screening in human cardiac organoids reveals a metabolic mechanism for cardiomyocyte cell cycle arrest. Proceedings of the National Academy of Sciences of the United States of America, 2017, 114, E8372-E8381. | 7.1 | 361 |
| 14 | Gaq proteins: molecular pharmacology and therapeutic potential. Cellular and Molecular Life Sciences, 2017, 74, 1379-1390. | 5.4 | 43 |
| 15 | Crim1 has cell-autonomous and paracrine roles during embryonic heart development. Scientific Reports, 2016, 6, 19832. | 3.3 | 6 |
| 16 | Epidermal Growth Factor Receptor Transactivation: Mechanisms, Pathophysiology, and Potential Therapies in the Cardiovascular System. Annual Review of Pharmacology and Toxicology, 2016, 56, 627-653. | 9.4 | 125 |
| 17 | Taste and Hypertension in Humans: Targeting Cardiovascular Disease. Current Pharmaceutical Design, 2016, 22, 2290-2305. | 1.9 | 15 |
| 18 | Helix 8 of the angiotensin- II type 1A receptor interacts with phosphatidylinositol phosphates and modulates membrane insertion. Scientific Reports, 2015, 5, 9972. | 3.3 | 12 |

| # | Article | IF | CITATIONS |
|----|---|------|-----------|
| 19 | Cardiac gene expression data and in silico analysis provide novel insights into human and mouse taste receptor gene regulation. Naunyn-Schmiedeberg's Archives of Pharmacology, 2015, 388, 1009-1027. | 3.0 | 23 |
| 20 | Variability in Human Bitter Taste Sensitivity to Chemically Diverse Compounds Can Be Accounted for by Differential TAS2R Activation. Chemical Senses, 2015, 40, 427-435. | 2.0 | 38 |
| 21 | Extracellular Surface Residues of the <i>α</i> _{1B} -Adrenoceptor Critical for G Protein–Coupled Receptor Function. Molecular Pharmacology, 2015, 87, 121-129. | 2.3 | 9 |
| 22 | Structural determinants for binding to angiotensin converting enzyme 2 (ACE2) and angiotensin receptors 1 and 2. Frontiers in Pharmacology, 2015, 6, 5. | 3.5 | 17 |
| 23 | G protein-coupled receptors in cardiac biology: old and new receptors. Biophysical Reviews, 2015, 7, 77-89. | 3.2 | 18 |
| 24 | International Union of Basic and Clinical Pharmacology. XCIX. Angiotensin Receptors: Interpreters of Pathophysiological Angiotensinergic Stimuli. Pharmacological Reviews, 2015, 67, 754-819. | 16.0 | 245 |
| 25 | Bitter taste receptor agonists elicit Gâ€proteinâ€dependent negative inotropy in the murine heart. FASEB Journal, 2014, 28, 4497-4508. | 0.5 | 72 |
| 26 | Extrasensory perception: Odorant and taste receptors beyond the nose and mouth. , 2014, 142, 41-61. | | 98 |
| 27 | A functional siRNA screen identifies genes modulating angiotensin II-mediated EGFR transactivation. Journal of Cell Science, 2013, 126, 5377-90. | 2.0 | 30 |
| 28 | Unravelling the molecular complexity of <scp>GPCR</scp> â€mediated <scp>EGFR</scp> transactivation using functional genomics approaches. FEBS Journal, 2013, 280, 5258-5268. | 4.7 | 53 |
| 29 | PAQR3 Modulates Insulin Signaling by Shunting Phosphoinositide 3-Kinase p110α to the Golgi Apparatus. Diabetes, 2013, 62, 444-456. | 0.6 | 52 |
| 30 | Expression, Regulation and Putative Nutrient-Sensing Function of Taste GPCRs in the Heart. PLoS ONE, 2013, 8, e64579. | 2.5 | 121 |
| 31 | PAQR10 and PAQR11 mediate Ras signaling in the Golgi apparatus. Cell Research, 2012, 22, 661-676. | 12.0 | 37 |
| 32 | Angiotensin Type 1A Receptors in C1 Neurons of the Rostral Ventrolateral Medulla Modulate the Pressor Response to Aversive Stress. Journal of Neuroscience, 2012, 32, 2051-2061. | 3.6 | 41 |
| 33 | Angiotensin 1A receptors transfected into caudal ventrolateral medulla inhibit baroreflex gain and stress responses. Cardiovascular Research, 2012, 96, 330-339. | 3.8 | 10 |
| 34 | Real-Time Measurement of F-Actin Remodelling during Exocytosis Using Lifeact-EGFP Transgenic Animals. PLoS ONE, 2012, 7, e39815. | 2.5 | 22 |
| 35 | Silencing Relaxin-3 in Nucleus Incertus of Adult Rodents: A Viral Vector-based Approach to Investigate Neuropeptide Function. PLoS ONE, 2012, 7, e42300. | 2.5 | 20 |
| 36 | Efferent projections of C3 adrenergic neurons in the rat central nervous system. Journal of Comparative Neurology, 2012, 520, 2352-2368. | 1.6 | 24 |

| # | Article | IF | CITATIONS |
|----|--|------|-----------|
| 37 | Regulation of angiotensinogen by angiotensin II in mouse primary astrocyte cultures. Journal of Neurochemistry, 2011, 119, 18-26. | 3.9 | 25 |
| 38 | Heteromerization of angiotensin receptors changes trafficking and arrestin recruitment profiles. Cellular Signalling, 2011, 23, 1767-1776. | 3.6 | 63 |
| 39 | A Single β-Amino Acid Substitution to Angiotensin II Confers AT ₂ Receptor Selectivity and Vascular Function. Hypertension, 2011, 57, 570-576. | 2.7 | 51 |
| 40 | Relative affinity of angiotensin peptides and novel ligands at AT1 and AT2 receptors. Clinical Science, 2011, 121, 297-303. | 4.3 | 241 |
| 41 | Determination of the Exact Molecular Requirements for Type 1 Angiotensin Receptor Epidermal Growth Factor Receptor Transactivation and Cardiomyocyte Hypertrophy. Hypertension, 2011, 57, 973-980. | 2.7 | 27 |
| 42 | The renin–angiotensin system and cancer: old dog, new tricks. Nature Reviews Cancer, 2010, 10, 745-759. | 28.4 | 438 |
| 43 | Expression of Angiotensin Type 1A Receptors in C1 Neurons Restores the Sympathoexcitation to Angiotensin in the Rostral Ventrolateral Medulla of Angiotensin Type 1A Knockout Mice. Hypertension, 2010, 56, 143-150. | 2.7 | 34 |
| 44 | Ligand-Supported Purification of the Urotensin-II Receptor. Molecular Pharmacology, 2010, 78, 639-647. | 2.3 | 5 |
| 45 | Glucocorticoids Suppress Growth in Neonatal Cardiomyocytes Co-Expressing AT ₂ and AT ₁ Angiotensin Receptors. Neonatology, 2010, 97, 257-265. | 2.0 | 6 |
| 46 | Differential Participation of Angiotensin II Type 1 and 2 Receptors in the Regulation of Cardiac Cell Death Triggered by Angiotensin II. American Journal of Hypertension, 2009, 22, 569-576. | 2.0 | 15 |
| 47 | High-Density Lipoprotein Modulates Glucose Metabolism in Patients With Type 2 Diabetes Mellitus. Circulation, 2009, 119, 2103-2111. | 1.6 | 363 |
| 48 | Prolonged RXFP1 and RXFP2 signaling can be explained by poor internalization and a lack of β-arrestin recruitment. American Journal of Physiology - Cell Physiology, 2009, 296, C1058-C1066. | 4.6 | 44 |
| 49 | Heritable pathologic cardiac hypertrophy in adulthood is preceded by neonatal cardiac growth restriction. American Journal of Physiology - Regulatory Integrative and Comparative Physiology, 2009, 296, R672-R680. | 1.8 | 31 |
| 50 | Angiotensin II Type 2 Receptor Antagonizes Angiotensin II Type 1 Receptor–Mediated Cardiomyocyte Autophagy. Hypertension, 2009, 53, 1032-1040. | 2.7 | 100 |
| 51 | Development and Optimization of MicroRNA against Relaxinâ€3. Annals of the New York Academy of Sciences, 2009, 1160, 261-264. | 3.8 | 5 |
| 52 | Role of helix 8 in G protein-coupled receptors based on structure–function studies on the type 1 angiotensin receptor. Molecular and Cellular Endocrinology, 2009, 302, 118-127. | 3.2 | 54 |
| 53 | Beta-arrestin 2 is required for complement C1q expression in macrophages and constrains factor-independent survival. Molecular Immunology, 2009, 47, 340-347. | 2.2 | 19 |
| 54 | Endothelin-1 activates ETA receptors on human vascular smooth muscle cells to yield proteoglycans with increased binding to LDL. Atherosclerosis, 2009, 205, 451-457. | 0.8 | 29 |

| # | Article | IF | CITATIONS |
|----|---|------|-----------|
| 55 | Immunoprecipitation and Phosphorylation of G Protein-Coupled Receptors. Methods in Molecular Biology, 2009, 552, 359-371. | 0.9 | 2 |
| 56 | The angiotensin II type 2 (AT2) receptor: an enigmatic seven transmembrane receptor. Frontiers in Bioscience - Landmark, 2009, Volume, 958. | 3.0 | 99 |
| 57 | Adenovirus-mediated delivery of relaxin reverses cardiac fibrosis. Molecular and Cellular Endocrinology, 2008, 280, 30-38. | 3.2 | 48 |
| 58 | Phospholipase C/Protein Kinase C Pathway Mediates Angiotensin II-Dependent Apoptosis in Neonatal Rat Cardiac Fibroblasts Expressing AT1 Receptor. Journal of Cardiovascular Pharmacology, 2008, 52, 184-190. | 1.9 | 27 |
| 59 | UBF levels determine the number of active ribosomal RNA genes in mammals. Journal of Cell Biology, 2008, 183, 1259-1274. | 5.2 | 171 |
| 60 | Type 1 angiotensin receptor pharmacology: Signaling beyond G proteins. , 2007, 113, 210-226. | | 76 |
| 61 | Is helix VIII of G proteinâ€coupled receptors (GPCRs) a lipidâ€activated signalling sensor?. FASEB Journal, 2007, 21, A614. | 0.5 | 0 |
| 62 | Effect of Dominant-Negative Epidermal Growth Factor Receptors on Cardiomyocyte Hypertrophy. Journal of Receptor and Signal Transduction Research, 2006, 26, 659-677. | 2.5 | 14 |
| 63 | Baroreceptor reflex stimulation does not induce cytomegalovirus promoter-driven transgene expression in the ventrolateral medulla in vivo. Autonomic Neuroscience: Basic and Clinical, 2006, 126-127, 150-155. | 2.8 | 0 |
| 64 | Fine mapping of Lvm1: a quantitative trait locus controlling heart size independently of blood pressure. Pulmonary Pharmacology and Therapeutics, 2006, 19, 70-73. | 2.6 | 5 |
| 65 | Tackling the EGFR in pathological tissue remodelling. Pulmonary Pharmacology and Therapeutics, 2006, 19, 74-78. | 2.6 | 25 |
| 66 | CNTF reverses obesity-induced insulin resistance by activating skeletal muscle AMPK. Nature Medicine, 2006, 12, 541-548. | 30.7 | 250 |
| 67 | Protein Kinase C Regulates the Cell Surface Activity of Endothelin-Converting Enzyme-1. International Journal of Peptide Research and Therapeutics, 2006, 12, 291-295. | 1.9 | 17 |
| 68 | Extended bioluminescence resonance energy transfer (eBRET) for monitoring prolonged protein–protein interactions in live cells. Cellular Signalling, 2006, 18, 1664-1670. | 3.6 | 98 |
| 69 | Interleukin-6 Increases Insulin-Stimulated Glucose Disposal in Humans and Glucose Uptake and Fatty Acid Oxidation In Vitro via AMP-Activated Protein Kinase. Diabetes, 2006, 55, 2688-2697. | 0.6 | 699 |
| 70 | Expression of Constitutively Active Angiotensin Receptors in the Rostral Ventrolateral Medulla Increases Blood Pressure. Hypertension, 2006, 47, 1054-1061. | 2.7 | 57 |
| 71 | Role of Angiotensin II Type 1A Receptor Phosphorylation, Phospholipase D, and Extracellular Calcium in Isoform-specific Protein Kinase C Membrane Translocation Responses. Journal of Biological Chemistry, 2006, 281, 26340-26349. | 3.4 | 15 |
| 72 | Evaluation of the Membrane-binding Properties of the Proximal Region of the Angiotensin II Receptor (AT1A) Carboxyl Terminus by Surface Plasmon Resonance. Analytical Sciences, 2005, 21, 171-174. | 1.6 | 19 |

| # | Article | IF | CITATIONS |
|----|---|------|-----------|
| 73 | G Protein Coupling and Second Messenger Generation Are Indispensable for Metalloprotease-dependent, Heparin-binding Epidermal Growth Factor Shedding through Angiotensin II Type-1 Receptor. Journal of Biological Chemistry, 2005, 280, 26592-26599. | 3.4 | 115 |
| 74 | Double Trouble for Type 1 Angiotensin Receptors in Atherosclerosis. New England Journal of Medicine, 2005, 352, 506-508. | 27.0 | 8 |
| 75 | Effect of Intrauterine Growth Restriction on the Number of Cardiomyocytes in Rat Hearts. Pediatric Research, 2005, 57, 796-800. | 2.3 | 151 |
| 76 | Helix I of β-Arrestin Is Involved in Postendocytic Trafficking but Is Not Required for Membrane Translocation, Receptor Binding, and Internalization. Molecular Pharmacology, 2005, 67, 375-382. | 2.3 | 10 |
| 77 | Dual Pathways for Nuclear Factor κB Activation by Angiotensin II in Vascular Smooth Muscle. Circulation Research, 2005, 97, 975-982. | 4.5 | 58 |
| 78 | The Angiotensin II Type 2 Receptor Causes Constitutive Growth of Cardiomyocytes and Does Not Antagonize Angiotensin II Type 1 Receptor–Mediated Hypertrophy. Hypertension, 2005, 46, 1347-1354. | 2.7 | 123 |
| 79 | The Angiotensin II Type 2 Receptor Causes Constitutive Growth of Cardiomyocytes and Does Not Antagonize Angiotensin II Type 1 Receptor–Mediated Hypertrophy. Hypertension, 2005, 46, 1347-1354. | 2.7 | 4 |
| 80 | Urotensin II Promotes Hypertrophy of Cardiac Myocytes via Mitogen-Activated Protein Kinases. Molecular Endocrinology, 2004, 18, 2344-2354. | 3.7 | 84 |
| 81 | What?s new in the renin-angiotensin system?. Cellular and Molecular Life Sciences, 2004, 61, 2695-2703. | 5.4 | 37 |
| 82 | What?s new in the renin-angiotensin system?. Cellular and Molecular Life Sciences, 2004, 61, 2687-2694. | 5.4 | 14 |
| 83 | p38 mitogen-activated protein kinase inhibition improves cardiac function and attenuates left ventricular remodeling following myocardial infarction in the rat. Journal of the American College of Cardiology, 2004, 44, 1679-1689. | 2.8 | 157 |
| 84 | Urotensin II: the old kid in town. Trends in Endocrinology and Metabolism, 2004, 15, 175-182. | 7.1 | 64 |
| 85 | Cardiovascular role of urotensin II: effect of chronic infusion in the rat. Peptides, 2004, 25, 1783-1788. | 2.4 | 34 |
| 86 | Agonist-dependent internalization of the angiotensin II type one receptor (AT1): role of C-terminus phosphorylation in recruitment of β-arrestins. Regulatory Peptides, 2004, 120, 141-148. | 1.9 | 20 |
| 87 | Angiotensinll mediates cardiomyocyte hypertrophic growth pathways via MMP-dependent HB-EGF liberation. International Journal of Peptide Research and Therapeutics, 2003, 10, 431-435. | 0.1 | 1 |
| 88 | Surface plasmon resonance spectroscopy in the study of membrane-mediated cell signalling. Journal of Peptide Science, 2003, 9, 77-89. | 1.4 | 52 |
| 89 | AngiotensinII mediates cardiomyocyte hypertrophic growth pathways via MMP-dependent HB-EGF liberation. International Journal of Peptide Research and Therapeutics, 2003, 10, 431-435. | 1.9 | 0 |
| 90 | Adrenomedullin inhibits angiotensin AT1A receptor expression and function in cardiac fibroblasts. Regulatory Peptides, 2003, 112, 131-137. | 1.9 | 15 |

| # | Article | IF | CITATIONS |
|-----|--|-----|-----------|
| 91 | Angiotensin receptors: form and function and distribution. International Journal of Biochemistry and Cell Biology, 2003, 35, 774-779. | 2.8 | 82 |
| 92 | Arresting angiotensin type 1 receptors. Trends in Endocrinology and Metabolism, 2003, 14, 130-136. | 7.1 | 36 |
| 93 | Direct Actions of Urotensin II on the Heart. Circulation Research, 2003, 93, 246-253. | 4.5 | 196 |
| 94 | Caveolin Interacts with the Angiotensin II Type 1 Receptor during Exocytic Transport but Not at the Plasma Membrane. Journal of Biological Chemistry, 2003, 278, 23738-23746. | 3.4 | 110 |
| 95 | Emerging Role of the Urotensin II System in Cardiovascular Disease. Cardiology, 2003, 3, 153-158. | 0.3 | 2 |
| 96 | Side-Chain Substitutions within Angiotensin II Reveal Different Requirements for Signaling, Internalization, and Phosphorylation of Type 1A Angiotensin Receptors. Molecular Pharmacology, 2002, 61, 768-777. | 2.3 | 227 |
| 97 | Adenoviral-Directed Expression of the Type 1A Angiotensin Receptor Promotes Cardiomyocyte Hypertrophy via Transactivation of the Epidermal Growth Factor Receptor. Circulation Research, 2002, 90, 135-142. | 4.5 | 159 |
| 98 | Electrostatic and Hydrophobic Forces Tether the Proximal Region of the Angiotensin II Receptor (AT1A) Carboxyl Terminus to Anionic Lipidsâ€. Biochemistry, 2002, 41, 7830-7840. | 2.5 | 42 |
| 99 | Casein Kinase II Sites in the Intracellular C-terminal Domain of the Thyrotropin-releasing Hormone Receptor and Chimeric Conadotropin-releasing Hormone Receptors Contribute to β-Arrestin-dependent Internalization. Journal of Biological Chemistry, 2001, 276, 18066-18074. | 3.4 | 63 |
| 100 | Angiotensin II enhances noradrenaline release from sympathetic nerves of the rat prostate via a novel angiotensin receptor: implications for the pathophysiology of benign prostatic hyperplasia. Journal of Endocrinology, 2001, 171, 97-108. | 2.6 | 43 |
| 101 | Association of β-Arrestin 1 with the Type 1A Angiotensin II Receptor Involves Phosphorylation of the Receptor Carboxyl Terminus and Correlates with Receptor Internalization. Molecular Endocrinology, 2001, 15, 1706-1719. | 3.7 | 74 |
| 102 | Association of Â-Arrestin 1 with the Type 1A Angiotensin II Receptor Involves Phosphorylation of the Receptor Carboxyl Terminus and Correlates with Receptor Internalization. Molecular Endocrinology, 2001, 15, 1706-1719. | 3.7 | 58 |
| 103 | Tethering of the Proximal Region of the Angiotensin II Receptor (AT1A) C-Terminus to the Cell Membrane. , 2001, , 293-294. | | 0 |
| 104 | Agonist-induced Phosphorylation of the Angiotensin II (AT1A) Receptor Requires Generation of a Conformation That Is Distinct from the Inositol Phosphate-signaling State. Journal of Biological Chemistry, 2000, 275, 2893-2900. | 3.4 | 95 |
| 105 | Regulation of angiotensin II type 1 (AT1) receptor function. Regulatory Peptides, 1999, 79, 9-23. | 1.9 | 90 |
| 106 | Identification of a Ca2+/calmodulin-binding domain within the carboxyl-terminus of the angiotensin II (AT1A) receptor. FEBS Letters, 1999, 455, 367-371. | 2.8 | 28 |
| 107 | Identification of protein kinase C phosphorylation sites in the angiotensin II (AT1A) receptor. Biochemical Journal, 1999, 343, 637-644. | 3.7 | 31 |
| 108 | Identification of protein kinase C phosphorylation sites in the angiotensin II (AT1A) receptor. Biochemical Journal, 1999, 343, 637. | 3.7 | 12 |

| # | Article | IF | CITATIONS |
|-----|---|-----|-----------|
| 109 | Phosphorylation of the Angiotensin II (AT1A) Receptor Carboxyl Terminus: A Role in Receptor Endocytosis. Molecular Endocrinology, 1998, 12, 1513-1524. | 3.7 | 81 |
| 110 | Phosphorylation of the Angiotensin II (AT1A) Receptor Carboxyl Terminus: A Role in Receptor Endocytosis. Molecular Endocrinology, 1998, 12, 1513-1524. | 3.7 | 31 |
| 111 | Evidence against a role for protein kinase C in the regulation of the angiotensin II (AT1A) receptor. European Journal of Pharmacology, 1996, 295, 119-122. | 3.5 | 7 |
| 112 | Activation of the STAT Pathway by Angiotensin II in T3CHO/AT1A Cells. Journal of Biological Chemistry, 1995, 270, 19059-19065. | 3.4 | 68 |
| 113 | Stable expression of a functional rat angiotensin II (AT1A) receptor in CHO-K1 cells: Rapid desensitization by angiotensin II. Molecular and Cellular Biochemistry, 1995, 146, 79-89. | 3.1 | 46 |
| 114 | Angiotensin II Receptor Endocytosis Involves Two Distinct Regions of the Cytoplasmic Tail. Journal of Biological Chemistry, 1995, 270, 22153-22159. | 3.4 | 106 |
| 115 | Stable Expression of a Truncated AT1A Receptor in CHO-K1 Cells. Journal of Biological Chemistry, 1995, 270, 207-213. | 3.4 | 121 |
| 116 | Molecular forms of rat angiotensinogen in plasma and brain: identification by isoelectric focusing and immunoblot analysis. Regulatory Peptides, 1995, 59, 31-41. | 1.9 | 3 |
| 117 | A Novel Inhibitory Role for Glucocorticoids in the Secretion of Angiotensinogen by C6 Glioma Cells. Journal of Neurochemistry, 1994, 62, 1296-1301. | 3.9 | 7 |
| 118 | Angiotensinogen Secretion by Single Rat Pituitary Cells: Detection by a Reverse Haemolytic Plaque Assay and Cell Identification by Immunocytochemistry. Neuroendocrinology, 1992, 55, 308-316. | 2.5 | 24 |
| 119 | Angiotensinogen is secreted by pure rat neuronal cell cultures. Brain Research, 1992, 588, 191-200. | 2.2 | 48 |
| 120 | Immunccytochemical Localization of Angiotensinogen in Rat Brain: Dependence of Neuronal Immunoreactivity on Method of Tissue Processing. Journal of Neuroendocrinology, 1991, 3, 653-660. | 2.6 | 18 |
| 121 | Oxytocin Receptors in the Mammary Gland and Reproductive Tract of a Marsupial, the Brushtail Possum (Trichosurus Vulpecula)1. Biology of Reproduction, 1991, 45, 673-679. | 2.7 | 18 |
| 122 | Effect of intra-ovarian infusion of oxytocin on plasma progesterone concentrations in pregnant ewes. Reproduction, 1991, 92, 453-460. | 2.6 | 4 |
| 123 | Immunocytochemical Localization of Angiotensinogen and Angiotensin II in the Rat Pituitary. Journal of Neuroendocrinology, 1990, 2, 297-304. | 2.6 | 9 |
| 124 | Angiotensin receptors in an Australian marsupial, the brushtail possum Trichosurus vulpecula. General and Comparative Endocrinology, 1990, 77, 116-126. | 1.8 | 3 |
| 125 | The immunocytochemical localization of angiotensinogen in the rat ovary. Cell and Tissue Research, 1990, 261, 367-373. | 2.9 | 36 |
| 126 | Uterine oxytocin receptors in an australian marsupial, the brushtail possum, Trichosurus vulpecula. Comparative Biochemistry and Physiology A, Comparative Physiology, 1990, 95, 135-138. | 0.6 | 6 |

| # | Article | IF | CITATIONS |
|-----|---|-----|-----------|
| 127 | Oxytocin receptors in the ovine corpus luteum. Journal of Endocrinology, 1989, 121, 117-123. | 2.6 | 38 |
| 128 | Immunocytochemical localization of angiotensinogen in the rat brain. Neuroscience, 1988, 25, 319-341. | 2.3 | 98 |
| 129 | Purification of Rat Angiotensinogen. Preparative Biochemistry and Biotechnology, 1986, 16, 45-59. | 0.5 | 11 |
| 130 | Liver angiotensin II receptors in the rat: binding properties and regulation by dietary Na+ and angiotensin II. Journal of Endocrinology, 1985, 106, 103-111. | 2.6 | 17 |
| 131 | Regulation of rat brain angiotensin II (AII) receptors by intravenous AII and low dietary Na+. Brain Research, 1985, 345, 54-61. | 2.2 | 22 |