

# Christina Pamporaki

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7550397/publications.pdf>

Version: 2024-02-01

37  
papers

960  
citations

623734

14  
h-index

454955

30  
g-index

38  
all docs

38  
docs citations

38  
times ranked

1012  
citing authors

#	ARTICLE	IF	CITATIONS
1	Pre- versus post-operative untargeted plasma nuclear magnetic resonance spectroscopy metabolomics of pheochromocytoma and paraganglioma. <i>Endocrine</i> , 2022, 75, 254-265.	2.3	3
2	Differences in clinical presentation and management between pre- and postsurgical diagnoses of urinary bladder paraganglioma: is there clinical relevance? A systematic review. <i>World Journal of Urology</i> , 2022, 40, 385-390.	2.2	8
3	Clonidine suppression test for a reliable diagnosis of pheochromocytoma: When to use. <i>Clinical Endocrinology</i> , 2022, 97, 541-550.	2.4	6
4	Head/neck paragangliomas: focus on tumor location, mutational status and plasma methoxytyramine. <i>Endocrine-Related Cancer</i> , 2022, 29, 213-224.	3.1	12
5	Targeting 11-Beta Hydroxylase With [131I]IMAZA: A Novel Approach for the Treatment of Advanced Adrenocortical Carcinoma. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2022, 107, e1348-e1355.	3.6	5
6	Improved Diagnostic Accuracy of Clonidine Suppression Testing Using an Age-Related Cutoff for Plasma Normetanephrine. <i>Hypertension</i> , 2022, 79, 1257-1264.	2.7	8
7	Determinants of disease-specific survival in patients with and without metastatic pheochromocytoma and paraganglioma. <i>European Journal of Cancer</i> , 2022, 169, 32-41.	2.8	18
8	Preanalytical Considerations and Outpatient Versus Inpatient Tests of Plasma Metanephrines to Diagnose Pheochromocytoma. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2022, 107, e3689-e3698.	3.6	4
9	Metastatic pheochromocytoma and paraganglioma: signs and symptoms related to catecholamine secretion. <i>Discover Oncology</i> , 2021, 12, 9.	2.1	5
10	Pseudohypoxic pheochromocytomas and paragangliomas dominate in children. <i>Pediatric Blood and Cancer</i> , 2021, 68, e28981.	1.5	14
11	Optimized procedures for testing plasma metanephrines in patients on hemodialysis. <i>Scientific Reports</i> , 2021, 11, 14706.	3.3	5
12	Targeted Metabolomics as a Tool in Discriminating Endocrine From Primary Hypertension. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2021, 106, e1111-e1128.	3.6	19
13	Plasma metanephrines and prospective prediction of tumor location, size and mutation type in patients with pheochromocytoma and paraganglioma. <i>Clinical Chemistry and Laboratory Medicine</i> , 2021, 59, 353-363.	2.3	32
14	Overnight/first-morning urine free metanephrines and methoxytyramine for diagnosis of pheochromocytoma and paraganglioma: is this an option?. <i>European Journal of Endocrinology</i> , 2020, 182, 499-509.	3.7	13
15	Blood sampling for metanephrines: to stick or stick and wait?. <i>Clinical Chemistry and Laboratory Medicine</i> , 2020, 58, 1609-1610.	2.3	0
16	Development of a Function-Integrative Sleeve for Medical Applications. <i>Sensors</i> , 2019, 19, 2588.	3.8	3
17	Reference intervals for LC-MS/MS measurements of plasma free, urinary free and urinary acid-hydrolyzed deconjugated normetanephrine, metanephrine and methoxytyramine. <i>Clinica Chimica Acta</i> , 2019, 490, 46-54.	1.1	50
18	Pheochromocytoma and paraganglioma: clinical feature-based disease probability in relation to catecholamine biochemistry and reason for disease suspicion. <i>European Journal of Endocrinology</i> , 2019, 181, 409-420.	3.7	58

#	ARTICLE	IF	CITATIONS
19	A high rate of modestly elevated plasma normetanephrine in a population referred for suspected PPGL when measured in a seated position. <i>European Journal of Endocrinology</i> , 2019, 181, 301-309.	3.7	25
20	Biochemical Diagnosis of Chromaffin Cell Tumors in Patients at High and Low Risk of Disease: Plasma versus Urinary Free or Deconjugated O-Methylated Catecholamine Metabolites. <i>Clinical Chemistry</i> , 2018, 64, 1646-1656.	3.2	121
21	Optimized Reference Intervals for Plasma Free Metanephrines in Patients With CKD. <i>American Journal of Kidney Diseases</i> , 2018, 72, 907-909.	1.9	19
22	Plasma methoxytyramine: clinical utility with metanephrines for diagnosis of pheochromocytoma and paraganglioma. <i>European Journal of Endocrinology</i> , 2017, 177, 103-113.	3.7	82
23	Reference intervals for plasma concentrations of adrenal steroids measured by LC-MS/MS: Impact of gender, age, oral contraceptives, body mass index and blood pressure status. <i>Clinica Chimica Acta</i> , 2017, 470, 115-124.	1.1	116
24	Characteristics of Pediatric vs Adult Pheochromocytomas and Paragangliomas. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2017, 102, 1122-1132.	3.6	120
25	The ovarian response to standard gonadotropin stimulation is influenced by AMHRII genotypes. <i>Gynecological Endocrinology</i> , 2016, 32, 641-645.	1.7	12
26	Dipping in Ambulatory Blood Pressure Monitoring Correlates With Overnight Urinary Excretion of Catecholamines and Sodium. <i>Journal of Clinical Hypertension</i> , 2016, 18, 921-926.	2.0	12
27	Endocrine Hypertension and Chronic Kidney Disease. , 2015, , 185-231.		0
28	Supine or Sitting? Economic and other considerations for use of plasma metanephrines for diagnosis of phaeochromocytoma. <i>Clinical Endocrinology</i> , 2015, 82, 463-464.	2.4	11
29	Biochemical diagnosis of phaeochromocytoma using plasma free normetanephrine, metanephrine and methoxytyramine: importance of supine sampling under fasting conditions. <i>Clinical Endocrinology</i> , 2014, 80, 478-486.	2.4	96
30	Seasonal variation in plasma free normetanephrine concentrations: implications for biochemical diagnosis of pheochromocytoma. <i>European Journal of Endocrinology</i> , 2014, 170, 349-357.	3.7	25
31	Seasonal variations of plasma normetanephrine levels: the authors reply. <i>European Journal of Endocrinology</i> , 2014, 170, L3.	3.7	2
32	The follicular outcome after standard gonadotropin stimulation is associated with ER1± and ER1² genotypes. <i>Endocrine</i> , 2014, 47, 930-935.	2.3	6
33	Association of the (TAAAA)n repeat polymorphism of SHBG gene with the age at menopause in Greek postmenopausal women. <i>Maturitas</i> , 2014, 78, 113-116.	2.4	2
34	Plasma free deconjugated metanephrines for diagnosis of phaeochromocytoma. <i>Clinical Endocrinology</i> , 2013, 79, 476-483.	2.4	15
35	Sperm flow cytometric parameters are associated with ICSI outcome. <i>Reproductive BioMedicine Online</i> , 2013, 26, 611-618.	2.4	10
36	The ovarian response to standard gonadotrophin stimulation depends on FSHR, SHBG and CYP19 gene synergism. <i>Journal of Assisted Reproduction and Genetics</i> , 2012, 29, 1185-1191.	2.5	19

#	ARTICLE	IF	CITATIONS
37	Adrenocortical Tumors and Pheochromocytoma/Paraganglioma Initially Mistaken as Neuroblastoma—Experiences From the GPOH-MET Registry. <i>Frontiers in Endocrinology</i> , 0, 13, .	3.5	4