Qingyuan Zhang

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/7575067/publications.pdf

Version: 2024-02-01

361413 552781 2,610 38 20 26 citations g-index h-index papers 39 39 39 2906 docs citations times ranked citing authors all docs

#	Article	IF	Citations
1	NASA's surface biology and geology designated observable: A perspective on surface imaging algorithms. Remote Sensing of Environment, 2021, 257, 112349.	11.0	148
2	Characterization of a seasonally snow-covered evergreen forest ecosystem. International Journal of Applied Earth Observation and Geoinformation, 2021, 103, 102464.	2.8	2
3	Evaluating impacts of snow, surface water, soil and vegetation on empirical vegetation and snow indices for the Utqiaġvik tundra ecosystem in Alaska with the LVS3 model. Remote Sensing of Environment, 2020, 240, 111677.	11.0	12
4	The 2013 FLEXâ€"US Airborne Campaign at the Parker Tract Loblolly Pine Plantation in North Carolina, USA. Remote Sensing, 2017, 9, 612.	4.0	27
5	EO-1 Data Quality and Sensor Stability with Changing Orbital Precession at the End of a 16 Year Mission. Remote Sensing, 2017, 9, 412.	4.0	17
6	Monitoring interannual variation in global crop yield using long-term AVHRR and MODIS observations. ISPRS Journal of Photogrammetry and Remote Sensing, 2016, 114, 191-205.	11.1	75
7	Integrating chlorophyll fAPAR and nadir photochemical reflectance index from EO-1/Hyperion to predict cornfield daily gross primary production. Remote Sensing of Environment, 2016, 186, 311-321.	11.0	22
8	Assessment of terrestrial vegetation dynamics from MODIS fAPAR <inf>chl</inf> product and land surface model. , 2016, , .		0
9	Preliminary fAPAR <inf>chl</inf> products from MODIS and hyperion. , 2016, , .		0
10	Estimation of crop gross primary production (GPP): II. Do scaled MODIS vegetation indices improve performance?. Agricultural and Forest Meteorology, 2015, 200, 1-8.	4.8	31
11	Estimation of crop gross primary production (GPP): I. impact of MODIS observation footprint and impact of vegetation BRDF characteristics. Agricultural and Forest Meteorology, 2014, 191, 51-63.	4.8	35
12	Estimation of crop gross primary production (GPP): fAPARchl versus MOD15A2 FPAR. Remote Sensing of Environment, 2014, 153, 1-6.	11.0	58
13	Impacts of light use efficiency and fPAR parameterization on gross primary production modeling. Agricultural and Forest Meteorology, 2014, 189-190, 187-197.	4.8	58
14	Variations of Foliage Chlorophyll fAPAR and Foliage Non-Chlorophyll fAPAR (fAPAR\$_{m chl}\$,) Tj ETQq0 0 0 rgBT Applied Earth Observations and Remote Sensing, 2013, 6, 2254-2264.	/Overlock 4.9	10 Tf 50 22 27
15	Integrating Solar Induced Fluorescence and the Photochemical Reflectance Index for Estimating Gross Primary Production in a Cornfield. Remote Sensing, 2013, 5, 6857-6879.	4.0	85
16	Using EO-1 Hyperion to Simulate HyspIRI Products for a Coniferous Forest: The Fraction of PAR Absorbed by Chlorophyll \$(hbox{fAPAR}_{m chl})\$ and Leaf Water Content (LWC). IEEE Transactions on Geoscience and Remote Sensing, 2012, 50, 1844-1852.	6.3	36
17	Canopy level Chlorophyll Fluorescence and the PRI in a cornfield. , 2012, , .		3
18	The photochemical reflectance index from directional cornfield reflectances: Observations and simulations. Remote Sensing of Environment, 2012, 124, 444-453.	11.0	29

#	Article	IF	CITATIONS
19	Spectral bio-indicator simulations for tracking photosynthetic activities in a corn field. Proceedings of SPIE, $2011, , .$	0.8	2
20	Applications using EO-1 hyperion AT-sensor and surface reflectance: comparisons and case studies. , 2010, , .		0
21	Using EO-1 Hyperion images to prototype environmental products for HysplRI. , 2010, , .		4
22	Introduction to fraction of absorbed par by canopy chlorophyll (fAPAR <inf>chl</inf>) and canopy leaf water content derived from hyperion, simulated HyspIRI and MODIS images., 2010,,.		1
23	Utilizing in situ directional hyperspectral measurements to validate bio-indicator simulations for a corn crop canopy. Ecological Informatics, 2010, 5, 330-338.	5. 2	19
24	Can a satellite-derived estimate of the fraction of PAR absorbed by chlorophyll (FAPARchl) improve predictions of light-use efficiency and ecosystem photosynthesis for a boreal aspen forest?. Remote Sensing of Environment, 2009, 113, 880-888.	11.0	102
25	Land cover classification with MODIS data in China. , 2009, , .		O
26	Comparisons between in Situ Anisotropic Reflectance Measurements and Simulations for Vegetation Canopies: Validation and Sensitivity Analysis. , 2008, , .		0
27	Detecting leaf phenology of seasonally moist tropical forests in South America with multi-temporal MODIS images. Remote Sensing of Environment, 2006, 103, 465-473.	11.0	179
28	Characterization of seasonal variation of forest canopy in a temperate deciduous broadleaf forest, using daily MODIS data. Remote Sensing of Environment, 2006, 105, 189-203.	11.0	69
29	Study on method of extracting winter wheat area planted based on spectral features using Terra/MODIS., 2005, 6043, 14.		0
30	Satellite-based modeling of gross primary production in a seasonally moist tropical evergreen forest. Remote Sensing of Environment, 2005, 94, 105-122.	11.0	242
31	Estimating light absorption by chlorophyll, leaf and canopy in a deciduous broadleaf forest using MODIS data and a radiative transfer model. Remote Sensing of Environment, 2005, 99, 357-371.	11.0	189
32	MODELING GROSS PRIMARY PRODUCTION OF AN EVERGREEN NEEDLELEAF FOREST USING MODIS AND CLIMATE DATA. , 2005, 15, 954-969.		177
33	Satellite-based modeling of gross primary production in an evergreen needleleaf forest. Remote Sensing of Environment, 2004, 89, 519-534.	11.0	682
34	Land cover characterization of Temperate East Asia using multi-temporal VEGETATION sensor data. Remote Sensing of Environment, 2004, 90, 477-489.	11.0	125
35	Sensitivity of vegetation indices to atmospheric aerosols: continental-scale observations in Northern Asia. Remote Sensing of Environment, 2003, 84, 385-392.	11.0	153
36	<title>High-resolution image to correct evapotranspiration of NOAA image</title> ., 2000, 4203, 21.		0

#	Article	IF	CITATIONS
37	<title>Analysis of vegetation green wave change in China using NOAA AVHRR data sets</title> ., 2000, , .		O
38	<title>Cereal production assessment for North Korea</title> ., 2000, 4203, 143.		0