

# Carlos L Arteaga

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7598168/publications.pdf>

Version: 2024-02-01

201  
papers

27,451  
citations

5876

81  
h-index

5965

160  
g-index

215  
all docs

215  
docs citations

215  
times ranked

30401  
citing authors

| #  | ARTICLE  | IF   | CITATIONS |
|----|--|------|-----------|
| 1  | Ribociclib as First-Line Therapy for HR-Positive, Advanced Breast Cancer. <i>New England Journal of Medicine</i> , 2016, 375, 1738-1748.   | 13.9 | 1,390     |
| 2  | Transforming Growth Factor- $\beta$ 1 Mediates Epithelial to Mesenchymal Transdifferentiation through a RhoA-dependent Mechanism. <i>Molecular Biology of the Cell</i> , 2001, 12, 27-36.  | 0.9  | 962       |
| 3  | ERBB Receptors: From Oncogene Discovery to Basic Science to Mechanism-Based Cancer Therapeutics. <i>Cancer Cell</i> , 2014, 25, 282-303.   | 7.7  | 817       |
| 4  | Treatment of HER2-positive breast cancer: current status and future perspectives. <i>Nature Reviews Clinical Oncology</i> , 2012, 9, 16-32.  | 12.5 | 735       |
| 5  | PKB/Akt mediates cell-cycle progression by phosphorylation of p27Kip1 at threonine 157 and modulation of its cellular localization. <i>Nature Medicine</i> , 2002, 8, 1145-1152.   | 15.2 | 729       |
| 6  | The PI3K/AKT Pathway as a Target for Cancer Treatment. <i>Annual Review of Medicine</i> , 2016, 67, 11-28.   | 5.0  | 631       |
| 7  | HER kinase inhibition in patients with HER2- and HER3-mutant cancers. <i>Nature</i> , 2018, 554, 189-194.  | 13.7 | 572       |
| 8  | Emergence of Constitutively Active Estrogen Receptor- $\beta$ Mutations in Pretreated Advanced Estrogen Receptor-Positive Breast Cancer. <i>Clinical Cancer Research</i> , 2014, 20, 1757-1767.  | 3.2  | 529       |
| 9  | TGF- $\beta$ 2 inhibition enhances chemotherapy action against triple-negative breast cancer. <i>Journal of Clinical Investigation</i> , 2013, 123, 1348-1358.   | 3.9  | 495       |
| 10 | Herceptin-induced inhibition of phosphatidylinositol-3 kinase and Akt is required for antibody-mediated effects on p27, cyclin D1, and antitumor action. <i>Cancer Research</i> , 2002, 62, 4132-41.   | 0.4  | 471       |
| 11 | Human Breast Cancer Cells Selected for Resistance to Trastuzumab <i>in vivo</i> Overexpress Epidermal Growth Factor Receptor and ErbB Ligands and Remain Dependent on the ErbB Receptor Network. <i>Clinical Cancer Research</i> , 2007, 13, 4909-4919.                    | 3.2  | 463       |
| 12 | Loss of PTEN/MMAC1/TEP in EGF receptor-expressing tumor cells counteracts the antitumor action of EGFR tyrosine kinase inhibitors. <i>Oncogene</i> , 2003, 22, 2812-2822.  | 2.6  | 449       |
| 13 | MYC and MCL1 Cooperatively Promote Chemotherapy-Resistant Breast Cancer Stem Cells via Regulation of Mitochondrial Oxidative Phosphorylation. <i>Cell Metabolism</i> , 2017, 26, 633-647.e7.   | 7.2  | 449       |
| 14 | Hyperactivation of phosphatidylinositol-3 kinase promotes escape from hormone dependence in estrogen receptor-positive human breast cancer. <i>Journal of Clinical Investigation</i> , 2010, 120, 2406-2413.   | 3.9  | 447       |
| 15 | Acquired resistance to EGFR tyrosine kinase inhibitors in cancer cells is mediated by loss of IGF-binding proteins. <i>Journal of Clinical Investigation</i> , 2008, 118, 2609-19.   | 3.9  | 443       |
| 16 | RAS/MAPK Activation Is Associated with Reduced Tumor-Infiltrating Lymphocytes in Triple-Negative Breast Cancer: Therapeutic Cooperation Between MEK and PD-1/PD-L1 Immune Checkpoint Inhibitors. <i>Clinical Cancer Research</i> , 2016, 22, 1499-1509.                    | 3.2  | 428       |
| 17 | Buparlisib plus fulvestrant versus placebo plus fulvestrant in postmenopausal, hormone receptor-positive, HER2-negative, advanced breast cancer (BELLE-2): a randomised, double-blind, placebo-controlled, phase 3 trial. <i>Lancet Oncology</i> , The, 2017, 18, 904-916. | 5.1  | 427       |
| 18 | HER2 kinase domain mutation results in constitutive phosphorylation and activation of HER2 and EGFR and resistance to EGFR tyrosine kinase inhibitors. <i>Cancer Cell</i> , 2006, 10, 25-38.   | 7.7  | 426       |

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 19 | Epidermal Growth Factor Receptor Dependence in Human Tumors: More Than Just Expression?. <i>Oncologist</i> , 2002, 7, 31-39.   | 1.9 | 424       |
| 20 | Molecular Profiling of the Residual Disease of Triple-Negative Breast Cancers after Neoadjuvant Chemotherapy Identifies Actionable Therapeutic Targets. <i>Cancer Discovery</i> , 2014, 4, 232-245.  | 7.7 | 413       |
| 21 | Overcoming Endocrine Resistance in Breast Cancer. <i>Cancer Cell</i> , 2020, 37, 496-513.  | 7.7 | 411       |
| 22 | Transcriptional and posttranslational up-regulation of HER3 (ErbB3) compensates for inhibition of the HER2 tyrosine kinase. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2011, 108, 5021-5026. | 3.3 | 403       |
| 23 | Blockade of TGF- $\beta$ 2 inhibits mammary tumor cell viability, migration, and metastases. <i>Journal of Clinical Investigation</i> , 2002, 109, 1551-1559.  | 3.9 | 402       |
| 24 | Mutations in the phosphatidylinositol 3-kinase pathway: role in tumor progression and therapeutic implications in breast cancer. <i>Breast Cancer Research</i> , 2011, 13, 224.  | 2.2 | 365       |
| 25 | Growth retardation and tumour inhibition by BRCA1. <i>Nature Genetics</i> , 1996, 12, 298-302.   | 9.4 | 359       |
| 26 | Phosphatidylinositol 3-Kinase and Antiestrogen Resistance in Breast Cancer. <i>Journal of Clinical Oncology</i> , 2011, 29, 4452-4461.   | 0.8 | 346       |
| 27 | Feedback upregulation of HER3 (ErbB3) expression and activity attenuates antitumor effect of PI3K inhibitors. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2012, 109, 2718-2723.               | 3.3 | 313       |
| 28 | Inhibition of TGF- $\beta$ 2 with neutralizing antibodies prevents radiation-induced acceleration of metastatic cancer progression. <i>Journal of Clinical Investigation</i> , 2007, 117, 1305-1313.                                       | 3.9 | 307       |
| 29 | PI3K/AKT/mTOR: role in breast cancer progression, drug resistance, and treatment. <i>Cancer and Metastasis Reviews</i> , 2016, 35, 515-524.  | 2.7 | 300       |
| 30 | MEK Inhibition Leads to PI3K/AKT Activation by Relieving a Negative Feedback on ERBB Receptors. <i>Cancer Research</i> , 2012, 72, 3228-3237.  | 0.4 | 287       |
| 31 | ER-Dependent E2F Transcription Can Mediate Resistance to Estrogen Deprivation in Human Breast Cancer. <i>Cancer Discovery</i> , 2011, 1, 338-351.  | 7.7 | 284       |
| 32 | BIM Expression in Treatment-Naïve Cancers Predicts Responsiveness to Kinase Inhibitors. <i>Cancer Discovery</i> , 2011, 1, 352-365.  | 7.7 | 268       |
| 33 | A Phase Ib Study of Alpelisib (BYL719), a PI3K- $\alpha$ -Specific Inhibitor, with Letrozole in ER+/HER2- Metastatic Breast Cancer. <i>Clinical Cancer Research</i> , 2017, 23, 26-34.   | 3.2 | 268       |
| 34 | Aberrant FGFR signaling mediates resistance to CDK4/6 inhibitors in ER+ breast cancer. <i>Nature Communications</i> , 2019, 10, 1373.  | 5.8 | 252       |
| 35 | Quantitative Optical Imaging of Primary Tumor Organoid Metabolism Predicts Drug Response in Breast Cancer. <i>Cancer Research</i> , 2014, 74, 5184-5194.   | 0.4 | 251       |
| 36 | Targeting the TGF- $\beta$ 2 signaling network in human neoplasia. <i>Cancer Cell</i> , 2003, 3, 531-536.  | 7.7 | 240       |

| #  | ARTICLE   | IF   | CITATIONS |
|----|---|------|-----------|
| 37 | Loss of Phosphatase and Tensin Homolog or Phosphoinositol-3 Kinase Activation and Response to Trastuzumab or Lapatinib in Human Epidermal Growth Factor Receptor 2â€œOverexpressing Locally Advanced Breast Cancers. <i>Journal of Clinical Oncology</i> , 2011, 29, 166-173. | 0.8  | 235       |
| 38 | Overview of epidermal growth factor receptor biology and its role as a therapeutic target in human neoplasia. <i>Seminars in Oncology</i> , 2002, 29, 3-9.  | 0.8  | 232       |
| 39 | Overall Survival with Ribociclib plus Letrozole in Advanced Breast Cancer. <i>New England Journal of Medicine</i> , 2022, 386, 942-950.   | 13.9 | 220       |
| 40 | Profiling of residual breast cancers after neoadjuvant chemotherapy identifies DUSP4 deficiency as a mechanism of drug resistance. <i>Nature Medicine</i> , 2012, 18, 1052-1059.  | 15.2 | 219       |
| 41 | ErbB-targeted therapeutic approaches in human cancer. <i>Experimental Cell Research</i> , 2003, 284, 122-130.   | 1.2  | 206       |
| 42 | Type I Transforming Growth Factor Î² Receptor Binds to and Activates Phosphatidylinositol 3-Kinase. <i>Journal of Biological Chemistry</i> , 2005, 280, 10870-10876.  | 1.6  | 201       |
| 43 | Increased Malignancy of Neu-Induced Mammary Tumors Overexpressing Active Transforming Growth Factor Î²1. <i>Molecular and Cellular Biology</i> , 2003, 23, 8691-8703.   | 1.1  | 190       |
| 44 | Trastuzumab Has Preferential Activity against Breast Cancers Driven by HER2 Homodimers. <i>Cancer Research</i> , 2011, 71, 1871-1882.   | 0.4  | 185       |
| 45 | Elevation of Receptor Tyrosine Kinase EphA2 Mediates Resistance to Trastuzumab Therapy. <i>Cancer Research</i> , 2010, 70, 299-308.   | 0.4  | 182       |
| 46 | Kinome-Wide RNA Interference Screen Reveals a Role for PDK1 in Acquired Resistance to CDK4/6 Inhibition in ER-Positive Breast Cancer. <i>Cancer Research</i> , 2017, 77, 2488-2499.   | 0.4  | 178       |
| 47 | Transforming Growth Factor Î² Enhances Epithelial Cell Survival via Akt-dependent Regulation of FKHL1. <i>Molecular Biology of the Cell</i> , 2001, 12, 3328-3339.  | 0.9  | 175       |
| 48 | Mutant <i>PIK3CA</i> accelerates HER2-driven transgenic mammary tumors and induces resistance to combinations of anti-HER2 therapies. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2013, 110, 14372-14377.                        | 3.3  | 168       |
| 49 | Conditional Overexpression of Active Transforming Growth Factor Î²1 In vivo Accelerates Metastases of Transgenic Mammary Tumors. <i>Cancer Research</i> , 2004, 64, 9002-9011.  | 0.4  | 164       |
| 50 | Stand Up to Cancer Phase Ib Study of Pan-Phosphoinositide-3-Kinase Inhibitor Buparlisib With Letrozole in Estrogen Receptor-Positive/Human Epidermal Growth Factor Receptor 2-Negative Metastatic Breast Cancer. <i>Journal of Clinical Oncology</i> , 2014, 32, 1202-1209.   | 0.8  | 159       |
| 51 | Resistance to HER2-directed antibodies and tyrosine kinase inhibitors. <i>Cancer Biology and Therapy</i> , 2011, 11, 793-800.   | 1.5  | 156       |
| 52 | Transforming Growth Factor Î² Engages TACE and ErbB3 To Activate Phosphatidylinositol-3 Kinase/Akt in ErbB2-Overexpressing Breast Cancer and Desensitizes Cells to Trastuzumab. <i>Molecular and Cellular Biology</i> , 2008, 28, 5605-5620.                                  | 1.1  | 153       |
| 53 | Invasion and metastasis of a mammary tumor involves TGF-? signaling. <i>International Journal of Cancer</i> , 2001, 91, 76-82.  | 2.3  | 148       |
| 54 | Autocrine Transforming Growth Factor-Î² Signaling Mediates Smad-independent Motility in Human Cancer Cells. <i>Journal of Biological Chemistry</i> , 2003, 278, 3275-3285.  | 1.6  | 148       |

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 55 | Dabrafenib and Trametinib in Patients With Tumors With <i>BRAF</i> <sup>V600E</sup> Mutations: Results of the NCI-MATCH Trial Subprotocol H. <i>Journal of Clinical Oncology</i> , 2020, 38, 3895-3904.             | 0.8 | 145       |
| 56 | Overexpression of HER2 (erbB2) in Human Breast Epithelial Cells Unmasks Transforming Growth Factor $\beta$ -induced Cell Motility. <i>Journal of Biological Chemistry</i> , 2004, 279, 24505-24513.                 | 1.6 | 144       |
| 57 | Challenges for the Clinical Development of PI3K Inhibitors: Strategies to Improve Their Impact in Solid Tumors. <i>Cancer Discovery</i> , 2019, 9, 482-491.   | 7.7 | 141       |
| 58 | In situ single-cell analysis identifies heterogeneity for PIK3CA mutation and HER2 amplification in HER2-positive breast cancer. <i>Nature Genetics</i> , 2015, 47, 1212-1219.                                      | 9.4 | 139       |
| 59 | A Kinome-Wide Screen Identifies the Insulin/IGF-I Receptor Pathway as a Mechanism of Escape from Hormone Dependence in Breast Cancer. <i>Cancer Research</i> , 2011, 71, 6773-6784.                                 | 0.4 | 138       |
| 60 | Targeting HER1/EGFR: a molecular approach to cancer therapy. <i>Seminars in Oncology</i> , 2003, 30, 3-14.  | 0.8 | 134       |
| 61 | Transforming Growth Factor $\beta$ Induces Clustering of HER2 and Integrins by Activating Src-Focal Adhesion Kinase and Receptor Association to the Cytoskeleton. <i>Cancer Research</i> , 2009, 69, 475-482.       | 0.4 | 126       |
| 62 | Inhibition of Mammalian Target of Rapamycin Is Required for Optimal Antitumor Effect of HER2 Inhibitors against HER2-Overexpressing Cancer Cells. <i>Clinical Cancer Research</i> , 2009, 15, 7266-7276.            | 3.2 | 124       |
| 63 | A versatile oblique plane microscope for large-scale and high-resolution imaging of subcellular dynamics. <i>ELife</i> , 2020, 9, .   | 2.8 | 120       |
| 64 | ErbB2/Neu-Induced, Cyclin D1-Dependent Transformation Is Accelerated in p27-Haploinsufficient Mammary Epithelial Cells but Impaired in p27-Null Cells. <i>Molecular and Cellular Biology</i> , 2002, 22, 2204-2219. | 1.1 | 113       |
| 65 | An Antibody That Locks HER3 in the Inactive Conformation Inhibits Tumor Growth Driven by HER2 or Neuregulin. <i>Cancer Research</i> , 2013, 73, 6024-6035.  | 0.4 | 109       |
| 66 | Inhibition of TGF $\beta$ signaling in cancer therapy. <i>Current Opinion in Genetics and Development</i> , 2006, 16, 30-37.  | 1.5 | 107       |
| 67 | HER3 Is Required for HER2-Induced Preneoplastic Changes to the Breast Epithelium and Tumor Formation. <i>Cancer Research</i> , 2012, 72, 2672-2682.   | 0.4 | 106       |
| 68 | Nivolumab Is Effective in Mismatch Repair-Deficient Noncolorectal Cancers: Results From Arm Z1D-A Subprotocol of the NCI-MATCH (EAY131) Study. <i>Journal of Clinical Oncology</i> , 2020, 38, 214-222.             | 0.8 | 106       |
| 69 | Triple-negative breast cancers with amplification of JAK2 at the 9p24 locus demonstrate JAK2-specific dependence. <i>Science Translational Medicine</i> , 2016, 8, 334ra53.   | 5.8 | 105       |
| 70 | Phase II Study of AZD4547 in Patients With Tumors Harboring Aberrations in the FGFR Pathway: Results From the NCI-MATCH Trial (EAY131) Subprotocol W. <i>Journal of Clinical Oncology</i> , 2020, 38, 2407-2417.    | 0.8 | 102       |
| 71 | The selective estrogen receptor downregulator GDC-0810 is efficacious in diverse models of ER+ breast cancer. <i>ELife</i> , 2016, 5, .   | 2.8 | 100       |
| 72 | Cardio-Oncology. <i>Circulation</i> , 2015, 132, 2248-2258.   | 1.6 | 99        |

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 73 | Trastuzumab-Resistant Cells Rely on a HER2-PI3K-FoxO-Survivin Axis and Are Sensitive to PI3K Inhibitors. <i>Cancer Research</i> , 2013, 73, 1190-1200.   | 0.4 | 98        |
| 74 | Enabling a Genetically Informed Approach to Cancer Medicine: A Retrospective Evaluation of the Impact of Comprehensive Tumor Profiling Using a Targeted Next-Generation Sequencing Panel. <i>Oncologist</i> , 2014, 19, 616-622. | 1.9 | 94        |
| 75 | Association of FGFR1 with ER $\pm$ Maintains Ligand-Independent ER Transcription and Mediates Resistance to Estrogen Deprivation in ER+ Breast Cancer. <i>Clinical Cancer Research</i> , 2017, 23, 6138-6150.                    | 3.2 | 94        |
| 76 | Treatment of Triple-Negative Breast Cancer with TORC1/2 Inhibitors Sustains a Drug-Resistant and Notch-Dependent Cancer Stem Cell Population. <i>Cancer Research</i> , 2016, 76, 440-452.  | 0.4 | 93        |
| 77 | Dual Blockade of HER2 in HER2-Overexpressing Tumor Cells Does Not Completely Eliminate HER3 Function. <i>Clinical Cancer Research</i> , 2013, 19, 610-619.   | 3.2 | 91        |
| 78 | Genomic profiling of ER <sup>+</sup> breast cancers after short-term estrogen suppression reveals alterations associated with endocrine resistance. <i>Science Translational Medicine</i> , 2017, 9, .                           | 5.8 | 91        |
| 79 | TBCRC 032 IB/II Multicenter Study: Molecular Insights to AR Antagonist and PI3K Inhibitor Efficacy in Patients with AR+ Metastatic Triple-Negative Breast Cancer. <i>Clinical Cancer Research</i> , 2020, 26, 2111-2123.         | 3.2 | 91        |
| 80 | Ribociclib plus letrozole versus letrozole alone in patients with de novo HR+, HER2 <sup>+</sup> advanced breast cancer in the randomized MONALEESA-2 trial. <i>Breast Cancer Research and Treatment</i> , 2018, 168, 127-134.   | 1.1 | 90        |
| 81 | Tyrosine kinase inhibitors. <i>Cancer Cell</i> , 2004, 5, 525-531.   | 7.7 | 89        |
| 82 | The brain microenvironment mediates resistance in luminal breast cancer to PI3K inhibition through HER3 activation. <i>Science Translational Medicine</i> , 2017, 9, .   | 5.8 | 89        |
| 83 | A Gene Expression Signature from Human Breast Cancer Cells with Acquired Hormone Independence Identifies MYC as a Mediator of Antiestrogen Resistance. <i>Clinical Cancer Research</i> , 2011, 17, 2024-2034.                    | 3.2 | 88        |
| 84 | Epidermal Growth Factor Receptor (EGFR) Antibody Down-regulates Mutant Receptors and Inhibits Tumors Expressing EGFR Mutations. <i>Journal of Biological Chemistry</i> , 2006, 281, 40183-40192.                                 | 1.6 | 85        |
| 85 | An Acquired HER2 <sup>T798I</sup> Gatekeeper Mutation Induces Resistance to Neratinib in a Patient with HER2 Mutant-Driven Breast Cancer. <i>Cancer Discovery</i> , 2017, 7, 575-585.  | 7.7 | 85        |
| 86 | Efficacy and Determinants of Response to HER Kinase Inhibition in HER2-Mutant Metastatic Breast Cancer. <i>Cancer Discovery</i> , 2020, 10, 198-213.   | 7.7 | 83        |
| 87 | Drug response in organoids generated from frozen primary tumor tissues. <i>Scientific Reports</i> , 2016, 6, 18889.  | 1.6 | 81        |
| 88 | Combination of Antibody That Inhibits Ligand-Independent HER3 Dimerization and a p110 $\alpha$ Inhibitor Potently Blocks PI3K Signaling and Growth of HER2+ Breast Cancers. <i>Cancer Research</i> , 2013, 73, 6013-6023.        | 0.4 | 79        |
| 89 | Autocrine IGF-I/insulin receptor axis compensates for inhibition of AKT in ER-positive breast cancer cells with resistance to estrogen deprivation. <i>Breast Cancer Research</i> , 2013, 15, R55.                               | 2.2 | 79        |
| 90 | Association with HSP90 Inhibits Cbl-Mediated Down-regulation of Mutant Epidermal Growth Factor Receptors. <i>Cancer Research</i> , 2006, 66, 6990-6997.  | 0.4 | 76        |

| #   | ARTICLE  | IF   | CITATIONS |
|-----|--|------|-----------|
| 91  | Extracellular Matrix/Integrin Signaling Promotes Resistance to Combined Inhibition of HER2 and PI3K in HER2+ Breast Cancer. <i>Cancer Research</i> , 2017, 77, 3280-3292.  | 0.4  | 76        |
| 92  | A Phase II Randomized Study of Neoadjuvant Letrozole Plus Alpelisib for Hormone Receptor-Positive, Human Epidermal Growth Factor Receptor 2-Negative Breast Cancer (NEO-ORB). <i>Clinical Cancer Research</i> , 2019, 25, 2975-2987.   | 3.2  | 76        |
| 93  | Cyclin-Dependent Kinase Inhibitor P27Kip1 Is Required for Mouse Mammary Gland Morphogenesis and Function. <i>Journal of Cell Biology</i> , 2001, 153, 917-932.   | 2.3  | 75        |
| 94  | Combined Blockade of Activating <i>ERBB2</i> Mutations and ER Results in Synthetic Lethality of ER+/HER2 Mutant Breast Cancer. <i>Clinical Cancer Research</i> , 2019, 25, 277-289.  | 3.2  | 74        |
| 95  | A Phase I-II Study of Combined Blockade of the ErbB Receptor Network with Trastuzumab and Gefitinib in Patients with HER2 (ErbB2)-Overexpressing Metastatic Breast Cancer. <i>Clinical Cancer Research</i> , 2008, 14, 6277-6283.  | 3.2  | 69        |
| 96  | ErbB3 Ablation Impairs PI3K/Akt-Dependent Mammary Tumorigenesis. <i>Cancer Research</i> , 2011, 71, 3941-3951.   | 0.4  | 69        |
| 97  | Direct inhibition of PI3K in combination with dual HER2 inhibitors is required for optimal antitumor activity in HER2+ breast cancer cells. <i>Breast Cancer Research</i> , 2014, 16, R9.  | 2.2  | 69        |
| 98  | <i>HER2</i> missense mutations have distinct effects on oncogenic signaling and migration. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2015, 112, E6205-14.   | 3.3  | 69        |
| 99  | Inhibition of Transforming Growth Factor- $\beta$ Signaling in Human Cancer: Targeting a Tumor Suppressor Network as a Therapeutic Strategy: Fig. 1.. <i>Clinical Cancer Research</i> , 2006, 12, 4142-4146.   | 3.2  | 68        |
| 100 | Human Breast Cancer Cells Harboring a Gatekeeper T798M Mutation in HER2 Overexpress EGFR Ligands and Are Sensitive to Dual Inhibition of EGFR and HER2. <i>Clinical Cancer Research</i> , 2013, 19, 5390-5401.   | 3.2  | 67        |
| 101 | $^{18}\text{F}$ -Fluoroestradiol PET/CT Measurement of Estrogen Receptor Suppression during a Phase I Trial of the Novel Estrogen Receptor-Targeted Therapeutic GDC-0810: Using an Imaging Biomarker to Guide Drug Dosage in Subsequent Trials. <i>Clinical Cancer Research</i> , 2017, 23, 3053-3060. | 3.2  | 66        |
| 102 | New Strategies in HER2-Overexpressing Breast Cancer: Many Combinations of Targeted Drugs Available. <i>Clinical Cancer Research</i> , 2011, 17, 952-958.   | 3.2  | 65        |
| 103 | HER3 and mutant EGFR meet MET. <i>Nature Medicine</i> , 2007, 13, 675-677.   | 15.2 | 64        |
| 104 | HER2-Overexpressing Breast Cancers Amplify FGFR Signaling upon Acquisition of Resistance to Dual Therapeutic Blockade of HER2. <i>Clinical Cancer Research</i> , 2017, 23, 4323-4334.  | 3.2  | 64        |
| 105 | RNA interference (RNAi) screening approach identifies agents that enhance paclitaxel activity in breast cancer cells. <i>Breast Cancer Research</i> , 2010, 12, R41.   | 2.2  | 63        |
| 106 | Clinical trial design and end points for epidermal growth factor receptor-targeted therapies: implications for drug development and practice. <i>Clinical Cancer Research</i> , 2003, 9, 1579-89.  | 3.2  | 63        |
| 107 | TROPiCS-02: A Phase III study investigating sacituzumab govitecan in the treatment of HR+/HER2-metastatic breast cancer. <i>Future Oncology</i> , 2020, 16, 705-715.   | 1.1  | 62        |
| 108 | Phase 2 study of buparlisib (BKM120), a pan-class I PI3K inhibitor, in patients with metastatic triple-negative breast cancer. <i>Breast Cancer Research</i> , 2020, 22, 120.  | 2.2  | 60        |

| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 109 | TGF- $\beta$ signaling promotes tumor vasculature by enhancing the pericyte-endothelium association. <i>BMC Cancer</i> , 2018, 18, 670.  | 1.1 | 58        |
| 110 | Melanoma response to anti-PD-L1 immunotherapy requires JAK1 signaling, but not JAK2. <i>OncImmunology</i> , 2018, 7, e1438106.   | 2.1 | 54        |
| 111 | FGFR1 Amplification Mediates Endocrine Resistance but Retains TORC Sensitivity in Metastatic Hormone Receptor-Positive (HR+) Breast Cancer. <i>Clinical Cancer Research</i> , 2019, 25, 6443-6451.   | 3.2 | 54        |
| 112 | HER (erbB) tyrosine kinase inhibitors in the treatment of breast cancer. <i>Seminars in Oncology</i> , 2002, 29, 4-10.   | 0.8 | 54        |
| 113 | Kinome-wide Functional Screen Identifies Role of PLK1 in Hormone-Independent, ER-Positive Breast Cancer. <i>Cancer Research</i> , 2015, 75, 405-414.   | 0.4 | 53        |
| 114 | Impact of Genomics on Personalized Cancer Medicine. <i>Clinical Cancer Research</i> , 2012, 18, 612-618.   | 3.2 | 52        |
| 115 | Buparlisib plus fulvestrant versus placebo plus fulvestrant for postmenopausal, hormone receptor-positive, human epidermal growth factor receptor 2-negative, advanced breast cancer: Overall survival results from BELLE-2. <i>European Journal of Cancer</i> , 2018, 103, 147-154.   | 1.3 | 52        |
| 116 | Elacestrant (RAD1901) exhibits anti-tumor activity in multiple ER+ breast cancer models resistant to CDK4/6 inhibitors. <i>Breast Cancer Research</i> , 2019, 21, 146.   | 2.2 | 52        |
| 117 | When Tumor Suppressor TGF- $\beta$ Meets the HER2 (ERBB2) Oncogene. <i>Journal of Mammary Gland Biology and Neoplasia</i> , 2011, 16, 81-88.   | 1.0 | 50        |
| 118 | Modeling the cancer patient with genetically engineered mice. <i>Cancer Cell</i> , 2004, 5, 115-120.   | 7.7 | 49        |
| 119 | Will PI3K pathway inhibitors be effective as single agents in patients with cancer?. <i>Oncotarget</i> , 2011, 2, 1314-1321.   | 0.8 | 49        |
| 120 | Optimal Targeting of HER2-PI3K Signaling in Breast Cancer: Mechanistic Insights and Clinical Implications. <i>Cancer Research</i> , 2013, 73, 3817-3820.   | 0.4 | 49        |
| 121 | Phase II Study of Lapatinib in Combination With Trastuzumab in Patients With Human Epidermal Growth Factor Receptor 2-Positive Metastatic Breast Cancer: Clinical Outcomes and Predictive Value of Early [ <sup>18</sup> F]Fluorodeoxyglucose Positron Emission Tomography Imaging (TBCRC 003). <i>Journal of Clinical Oncology</i> , 2015, 33, 2623-2631. | 0.8 | 49        |
| 122 | Trastuzumab, an appropriate first-line single-agent therapy for HER2-overexpressing metastatic breast cancer. <i>Breast Cancer Research</i> , 2003, 5, 96-100.   | 2.2 | 48        |
| 123 | EGF Receptor As a Therapeutic Target: Patient Selection and Mechanisms of Resistance to Receptor-Targeted Drugs. <i>Journal of Clinical Oncology</i> , 2003, 21, 289s-291.   | 0.8 | 48        |
| 124 | EGF receptor mutations in lung cancer: From humans to mice and maybe back to humans. <i>Cancer Cell</i> , 2006, 9, 421-423.  | 7.7 | 47        |
| 125 | Co-occurring gain-of-function mutations in HER2 and HER3 modulate HER2/HER3 activation, oncogenesis, and HER2 inhibitor sensitivity. <i>Cancer Cell</i> , 2021, 39, 1099-1114.e8.  | 7.7 | 45        |
| 126 | Activating PIK3CA Mutations Induce an Epidermal Growth Factor Receptor (EGFR)/Extracellular Signal-regulated Kinase (ERK) Paracrine Signaling Axis in Basal-like Breast Cancer*. <i>Molecular and Cellular Proteomics</i> , 2015, 14, 1959-1976.   | 2.5 | 44        |



| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 127 | Discovery of Potent Myeloid Cell Leukemia-1 (Mcl-1) Inhibitors That Demonstrate in Vivo Activity in Mouse Xenograft Models of Human Cancer. <i>Journal of Medicinal Chemistry</i> , 2019, 62, 3971-3988.                                       | 2.9 | 44        |
| 128 | Metabolic modulation by CDK4/6 inhibitor promotes chemokine-mediated recruitment of T <sub>H</sub> 1 cells into mammary tumors. <i>Cell Reports</i> , 2021, 35, 108944.  | 2.9 | 44        |
| 129 | ERBB1 and ERBB2 Have Distinct Functions in Tumor Cell Invasion and Intravasation. <i>Clinical Cancer Research</i> , 2009, 15, 3733-3739.   | 3.2 | 43        |
| 130 | Systematic Prioritization of Druggable Mutations in ~45000 Genomes Across 16 Cancer Types Using a Structural Genomics-based Approach. <i>Molecular and Cellular Proteomics</i> , 2016, 15, 642-656.  | 2.5 | 43        |
| 131 | The multifunctional role of transforming growth factor (TGF)- $\beta$ s on mammary epithelial cell biology. <i>Breast Cancer Research and Treatment</i> , 1996, 38, 49-56.   | 1.1 | 41        |
| 132 | Challenges in the development of anti-epidermal growth factor receptor therapies in breast cancer. <i>Seminars in Oncology</i> , 2004, 31, 3-8.  | 0.8 | 38        |
| 133 | Is There a Future for AKT Inhibitors in the Treatment of Cancer?. <i>Clinical Cancer Research</i> , 2016, 22, 2599-2601.   | 3.2 | 38        |
| 134 | Convergence of p53 and Transforming Growth Factor $\beta$ 2 (TGF $\beta$ 2) Signaling on Activating Expression of the Tumor Suppressor Gene maspin in Mammary Epithelial Cells. <i>Journal of Biological Chemistry</i> , 2007, 282, 5661-5669. | 1.6 | 37        |
| 135 | In vivo hyperspectral imaging of microvessel response to trastuzumab treatment in breast cancer xenografts. <i>Biomedical Optics Express</i> , 2014, 5, 2247.  | 1.5 | 37        |
| 136 | Epidermal Growth Factor Receptor Dependence in Human Tumors: More Than Just Expression?. <i>Oncologist</i> , 2002, 7, 31-39.   | 1.9 | 37        |
| 137 | Why Is This Effective HSP90 Inhibitor Not Being Developed in HER2+ Breast Cancer?. <i>Clinical Cancer Research</i> , 2011, 17, 4919-4921.  | 3.2 | 35        |
| 138 | Pooled ctDNA analysis of the MONALEESA (ML) phase III advanced breast cancer (ABC) trials.. <i>Journal of Clinical Oncology</i> , 2020, 38, 1009-1009.   | 0.8 | 34        |
| 139 | Neoadjuvant Trials in ER+ Breast Cancer: A Tool for Acceleration of Drug Development and Discovery. <i>Cancer Discovery</i> , 2017, 7, 561-574.  | 7.7 | 33        |
| 140 | Hyperactivation of TORC1 Drives Resistance to the Pan-HER Tyrosine Kinase Inhibitor Neratinib in HER2-Mutant Cancers. <i>Cancer Cell</i> , 2020, 37, 183-199.e5.   | 7.7 | 33        |
| 141 | Metabolic diversity within breast cancer brain-tropic cells determines metastatic fitness. <i>Cell Metabolism</i> , 2022, 34, 90-105.e7.   | 7.2 | 33        |
| 142 | PIK3CA Activating Mutations: A Discordant Role in Early Versus Advanced Hormone-Dependent Estrogen Receptor-Positive Breast Cancer?. <i>Journal of Clinical Oncology</i> , 2014, 32, 2932-2934.  | 0.8 | 32        |
| 143 | Collagen density and alignment in responsive and resistant trastuzumab-treated breast cancer xenografts. <i>Journal of Biomedical Optics</i> , 2015, 20, 026004.   | 1.4 | 32        |
| 144 | Overview of rationale and clinical trials with signal transduction inhibitors in lung cancer. <i>Seminars in Oncology</i> , 2002, 29, 15-26.   | 0.8 | 31        |

| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 145 | PIK3CA and MAP3K1 alterations imply luminal A status and are associated with clinical benefit from pan-PI3K inhibitor buparlisib and letrozole in ER+ metastatic breast cancer. <i>Npj Breast Cancer</i> , 2019, 5, 31.                  | 2.3 | 31        |
| 146 | Nuclear FGFR1 Regulates Gene Transcription and Promotes Antiestrogen Resistance in ER+ Breast Cancer. <i>Clinical Cancer Research</i> , 2021, 27, 4379-4396.   | 3.2 | 30        |
| 147 | Complex role of tumor cell transforming growth factor (TGF)- $\beta$ 2s on breast carcinoma progression. <i>Journal of Mammary Gland Biology and Neoplasia</i> , 1996, 1, 373-380.   | 1.0 | 29        |
| 148 | An ERBB1-3 Neutralizing Antibody Mixture With High Activity Against Drug-Resistant HER2+ Breast Cancers With ERBB Ligand Overexpression. <i>Journal of the National Cancer Institute</i> , 2017, 109, .                                  | 3.0 | 29        |
| 149 | Extended Adjuvant Therapy with Neratinib Plus Fulvestrant Blocks ER/HER2 Crosstalk and Maintains Complete Responses of ER+/HER2+ Breast Cancers: Implications to the ExteNET Trial. <i>Clinical Cancer Research</i> , 2019, 25, 771-783. | 3.2 | 29        |
| 150 | A kinase-inactive type II TGF $\beta$ 2 receptor impairs BMP signaling in human breast cancer cells. <i>Biochemical and Biophysical Research Communications</i> , 2003, 301, 108-112.  | 1.0 | 28        |
| 151 | PIK3CA C2 Domain Deletions Hyperactivate Phosphoinositide 3-kinase (PI3K), Generate Oncogene Dependence, and Are Exquisitely Sensitive to PI3K Inhibitors. <i>Clinical Cancer Research</i> , 2018, 24, 1426-1435.                        | 3.2 | 27        |
| 152 | ER+ Breast Cancers Resistant to Prolonged Neoadjuvant Letrozole Exhibit an E2F4 Transcriptional Program Sensitive to CDK4/6 Inhibitors. <i>Clinical Cancer Research</i> , 2018, 24, 2517-2529.   | 3.2 | 26        |
| 153 | Phase II Study of Copanlisib in Patients With Tumors With PIK3CA Mutations: Results From the NCI-MATCH ECOG-ACRIN Trial (EAY131) Subprotocol Z1F. <i>Journal of Clinical Oncology</i> , 2022, 40, 1552-1561.                             | 0.8 | 26        |
| 154 | Proline rich 11 (PRR11) overexpression amplifies PI3K signaling and promotes antiestrogen resistance in breast cancer. <i>Nature Communications</i> , 2020, 11, 5488.  | 5.8 | 25        |
| 155 | Tumor p38MAPK signaling enhances breast carcinoma vascularization and growth by promoting expression and deposition of pro-tumorigenic factors. <i>Oncotarget</i> , 2017, 8, 61969-61981.  | 0.8 | 25        |
| 156 | Evidence for a positive role of transforming growth factor- $\beta$ 2 in human breast cancer cell tumorigenesis. <i>Journal of Cellular Biochemistry</i> , 1993, 53, 187-193.  | 1.2 | 24        |
| 157 | Ligand-independent phosphorylation of Y869 (Y845) links mutant EGFR signaling to stat-mediated gene expression. <i>Experimental Cell Research</i> , 2008, 314, 413-419.  | 1.2 | 23        |
| 158 | NCCN Oncology Research Program's Investigator Steering Committee and NCCN Best Practices Committee Molecular Profiling Surveys. <i>Journal of the National Comprehensive Cancer Network: JNCCN</i> , 2015, 13, 1337-1346.                | 2.3 | 23        |
| 159 | Differential Outcomes in Codon 12/13 and Codon 61 NRAS-Mutated Cancers in the Phase II NCI-MATCH Trial of Binimetinib in Patients with NRAS-Mutated Tumors. <i>Clinical Cancer Research</i> , 2021, 27, 2996-3004.                       | 3.2 | 23        |
| 160 | A Mathematical Model Quantifies Proliferation and Motility Effects of TGF- $\beta$ 2 on Cancer Cells. <i>Computational and Mathematical Methods in Medicine</i> , 2009, 10, 71-83.   | 0.7 | 22        |
| 161 | Discordant Cellular Response to Presurgical Letrozole in Bilateral Synchronous ER+ Breast Cancers with a KRAS Mutation or FGFR1 Gene Amplification. <i>Molecular Cancer Therapeutics</i> , 2012, 11, 2301-2305.                          | 1.9 | 22        |
| 162 | Conditional Loss of ErbB3 Delays Mammary Gland Hyperplasia Induced by Mutant PIK3CA without Affecting Mammary Tumor Latency, Gene Expression, or Signaling. <i>Cancer Research</i> , 2013, 73, 4075-4085.                                | 0.4 | 22        |

| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 163 | Functional KRAS mutations and a potential role for PI3K/AKT activation in Wilms tumors. <i>Molecular Oncology</i> , 2017, 11, 405-421.   | 2.1 | 22        |
| 164 | Reply to "p53 and secreted tumour suppressors". <i>Nature Genetics</i> , 1996, 13, 269-272.  | 9.4 | 20        |
| 165 | Characterization of breast cancers with PI3K mutations in an academic practice setting using SNaPshot profiling. <i>Breast Cancer Research and Treatment</i> , 2014, 145, 389-399.   | 1.1 | 20        |
| 166 | Targeting tyrosine-kinases and estrogen receptor abrogates resistance to endocrine therapy in breast cancer. <i>Oncotarget</i> , 2014, 5, 9049-9064.   | 0.8 | 20        |
| 167 | Phase Ib Study of Safety and Pharmacokinetics of the PI3K Inhibitor SAR245408 with the HER3-Neutralizing Human Antibody SAR256212 in Patients with Solid Tumors. <i>Clinical Cancer Research</i> , 2017, 23, 3520-3528.            | 3.2 | 19        |
| 168 | Is Dual mTORC1 and mTORC2 Therapeutic Blockade Clinically Feasible in Cancer?. <i>JAMA Oncology</i> , 2019, 5, 1564.   | 3.4 | 19        |
| 169 | PIK3CA mutations in Peruvian patients with HER2-amplified and triple negative non-metastatic breast cancers. <i>Hematology/ Oncology and Stem Cell Therapy</i> , 2014, 7, 142-148.   | 0.6 | 18        |
| 170 | Cdk Inhibitor p27Kip1 and Hormone Dependence in Breast Cancer. <i>Clinical Cancer Research</i> , 2004, 10, 368s-371s.  | 3.2 | 18        |
| 171 | Dual inhibition of Type I and Type III PI3 kinases increases tumor cell apoptosis in HER2+ breast cancers. <i>Breast Cancer Research</i> , 2015, 17, 148.  | 2.2 | 17        |
| 172 | Clinical Development of Phosphatidylinositol-3 Kinase Pathway Inhibitors. <i>Current Topics in Microbiology and Immunology</i> , 2010, 347, 189-208.   | 0.7 | 15        |
| 173 | First-line ribociclib (RIB) + letrozole (LET) in hormone receptor-positive (HR+), HER2-negative (HER2-) advanced breast cancer (ABC): MONALEESA-2 biomarker analyses.. <i>Journal of Clinical Oncology</i> , 2018, 36, 1022-1022.  | 0.8 | 15        |
| 174 | Phosphatase and Tensin Homolog Deficiency and Resistance to Trastuzumab and Chemotherapy. <i>Journal of Clinical Oncology</i> , 2013, 31, 2073-2075.   | 0.8 | 14        |
| 175 | Estrogen receptor coregulator binding modulator (ERX-11) enhances the activity of CDK4/6 inhibitors against estrogen receptor-positive breast cancers. <i>Breast Cancer Research</i> , 2019, 21, 150.                              | 2.2 | 14        |
| 176 | Transcriptome- and proteome-oriented identification of dysregulated eIF4G, STAT3, and Hippo pathways altered by PIK3CA H1047R in HER2/ER-positive breast cancer. <i>Breast Cancer Research and Treatment</i> , 2016, 160, 457-474. | 1.1 | 13        |
| 177 | FGFR signaling and endocrine resistance in breast cancer: Challenges for the clinical development of FGFR inhibitors. <i>Biochimica Et Biophysica Acta: Reviews on Cancer</i> , 2021, 1876, 188595.                                | 3.3 | 13        |
| 178 | Dead-box or black-box: is DDX1 a potential biomarker in breast cancer?. <i>Breast Cancer Research and Treatment</i> , 2011, 127, 65-67.  | 1.1 | 12        |
| 179 | Epigenetic Repression of STING by MYC Promotes Immune Evasion and Resistance to Immune Checkpoint Inhibitors in Triple-Negative Breast Cancer. <i>Cancer Immunology Research</i> , 2022, 10, 829-843.                              | 1.6 | 12        |
| 180 | Exploring Biomarkers of Phosphoinositide 3-Kinase Pathway Activation in the Treatment of Hormone Receptor Positive, Human Epidermal Growth Receptor 2 Negative Advanced Breast Cancer. <i>Oncologist</i> , 2019, 24, 305-312.      | 1.9 | 11        |

| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 181 | Dabrafenib and trametinib in patients with tumors with BRAF V600E/K mutations: Results from the molecular analysis for therapy choice (MATCH) Arm H.. Journal of Clinical Oncology, 2019, 37, 3002-3002.   | 0.8 | 10        |
| 182 | Phase II Study of Taselisib in <i>PIK3CA</i> -Mutated Solid Tumors Other Than Breast and Squamous Lung Cancer: Results From the NCI-MATCH ECOG-ACRIN Trial (EAY131) Subprotocol I. JCO Precision Oncology, 2022, 6, e2100424.  | 1.5 | 9         |
| 183 | Inhibiting tyrosine kinases: successes and limitations. Cancer Biology and Therapy, 2003, 2, S79-83.   | 1.5 | 9         |
| 184 | Progress in Breast Cancer: Overview. Clinical Cancer Research, 2013, 19, 6353-6359.  | 3.2 | 8         |
| 185 | Motivation for Launching a Cancer Metastasis Inhibition (CMI) Program. Targeted Oncology, 2018, 13, 61-68.   | 1.7 | 8         |
| 186 | Targeting LIPA independent of its lipase activity is a therapeutic strategy in solid tumors via induction of endoplasmic reticulum stress. Nature Cancer, 2022, 3, 866-884.  | 5.7 | 8         |
| 187 | ERBB receptors in cancer: signaling from the inside. Breast Cancer Research, 2011, 13, 304.  | 2.2 | 7         |
| 188 | Neratinib: Inching Up on the Cure Rate of HER2+ Breast Cancer?. Clinical Cancer Research, 2018, 24, 3483-3485.   | 3.2 | 6         |
| 189 | Drug-Resistant Brain Metastases: A Role for Pharmacology, Tumor Evolution, and Too-Late Therapy. Cancer Discovery, 2015, 5, 1124-1126.   | 7.7 | 5         |
| 190 | Unexpected Benefit from Alpelisib and Fulvestrant in a Woman with Highly Pre-treated ER-Positive, HER2-Negative PIK3CA Mutant Metastatic Breast Cancer. Clinical Drug Investigation, 2018, 38, 1071-1075.  | 1.1 | 4         |
| 191 | Reply to T. Shimoi et al and Y. Shimanuki et al. Journal of Clinical Oncology, 2021, 39, JCO.21.01905.   | 0.8 | 3         |
| 192 | Abstract GS2-01: Overall survival subgroup analysis by metastatic site from the phase 3 MONALEESA-2 study of first-line ribociclib + letrozole in postmenopausal patients with advanced HR+/HER2 <sup>+</sup> breast cancer. Cancer Research, 2022, 82, GS2-01-GS2-01. | 0.4 | 2         |
| 193 | Abstract PD2-05: Genomic profiling of PAM50-based intrinsic subtypes in HR+/HER2- advanced breast cancer (ABC) across the MONALEESA (ML) studies. Cancer Research, 2022, 82, PD2-05-PD2-05.  | 0.4 | 2         |
| 194 | Phase 1 pilot study with dose expansion of chemotherapy in combination with CD40 agonist and Flt3 ligand in metastatic triple-negative breast cancer.. Journal of Clinical Oncology, 2022, 40, TPS1126-TPS1126.  | 0.8 | 2         |
| 195 | Abstract P4-01-02: A spectrum of secondary mutations in <i>HER2</i> augment breast cancer cell growth and reduce neratinib sensitivity in <i>HER2</i> -mutant breast cancer. Cancer Research, 2022, 82, P4-01-02-P4-01-02.   | 0.4 | 1         |
| 196 | Abstract GS3-09: Loss of <i>ASXL1</i> tumor suppressor promotes resistance to CDK4/6 inhibitors in ER+ breast cancer. Cancer Research, 2022, 82, GS3-09-GS3-09.  | 0.4 | 1         |
| 197 | Abstract PD2-01: A platform of CDK4/6 inhibitor-resistant patient-derived breast cancer organoids illuminates mechanisms of resistance and therapeutic vulnerabilities. Cancer Research, 2022, 82, PD2-01-PD2-01.  | 0.4 | 1         |
| 198 | Enhancement of Epidermal Growth Factor Receptor-degradation Pathway in Acquired Gefitinib-resistant Human Non-small Cell Lung Cancer Cell Lines. The Showa University Journal of Medical Sciences, 2004, 16, 147-159.  | 0.1 | 0         |

| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 199 | Abstract PR05: P-REX1 creates a positive feedback loop to activate growth factor receptor/PI3K signaling. , 2013, , .  |     | 0         |
| 200 | Hyperactivation of Torc1 Drives Resistance to the Pan-Her Tyrosine Kinase Inhibitor Neratinib in Her2-Mutant Cancers. SSRN Electronic Journal, 0, , .                                    | 0.4 | 0         |
| 201 | Abstract P5-17-09: A genome-wide CRISPR screen identifies PRMT5 as a novel therapeutic target in ER+/ <i>RB1</i> -deficient breast cancer. Cancer Research, 2022, 82, P5-17-09-P5-17-09. | 0.4 | 0         |