

# Thomas Seidl

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7607214/publications.pdf>

Version: 2024-02-01

29  
papers

1,664  
citations

430874

18  
h-index

552781

26  
g-index

30  
all docs

30  
docs citations

30  
times ranked

3097  
citing authors

#	ARTICLE	IF	CITATIONS
1	Deep phenotyping of Tregs identifies an immune signature for idiopathic aplastic anemia and predicts response to treatment. <i>Blood</i> , 2016, 128, 1193-1205.	1.4	117
2	Expansion of myeloid derived suppressor cells correlates with number of T regulatory cells and disease progression in myelodysplastic syndrome. <i>Oncotarget</i> , 2016, 5, e1062208.	4.6	97
3	RPL27A is a target of miR-595 and may contribute to the myelodysplastic phenotype through ribosomal dysgenesis. <i>Oncotarget</i> , 2016, 7, 47875-47890.	1.8	19
4	SF3B1 mutant MDS-initiating cells may arise from the haematopoietic stem cell compartment. <i>Nature Communications</i> , 2015, 6, 10004.	12.8	68
5	JAK inhibition induces silencing of T Helper cytokine secretion and a profound reduction in T regulatory cells. <i>British Journal of Haematology</i> , 2015, 171, 60-73.	2.5	73
6	Spliceosome mutations exhibit specific associations with epigenetic modifiers and proto-oncogenes mutated in myelodysplastic syndrome. <i>Haematologica</i> , 2013, 98, 1058-1066.	3.5	67
7	The effects of 5-azacytidine on the function and number of regulatory T cells and T-effectors in myelodysplastic syndrome. <i>Haematologica</i> , 2013, 98, 1196-1205.	3.5	91
8	JAK Inhibition Reduces CD25 high CD27+ FOXP3+ T Regulatory Cells and Causes a Silencing Of T Effector Cells In Patients With Myeloproliferative Neoplasms Whilst Promoting a TH17 Phenotype. <i>Blood</i> , 2013, 122, 4092-4092.	1.4	8
9	SF3B1 Mutant Clones From Patients With Refractory Anaemia With Ringed Sideroblasts (RARS) Originate From The Early Haematopoietic Stem Cells and Maintain Their Engraftment Potential. <i>Blood</i> , 2013, 122, 262-262.	1.4	0
10	Myeloid Derived Suppressor Cell Expansion Correlates With Increased regulatory T Cells In Myelodysplastic Syndrome. <i>Blood</i> , 2013, 122, 2766-2766.	1.4	0
11	Functional characterization of CD4+ T cells in aplastic anemia. <i>Blood</i> , 2012, 119, 2033-2043.	1.4	140
12	B cell agonists upregulate AID and APOBEC3G deaminases, which induce IgA and IgG class antibodies and anti-viral function. <i>Immunology</i> , 2012, 135, 207-215.	4.4	14
13	Expansion of CD141int Dendritic Cells in Myelodysplastic Syndrome (MDS). <i>Blood</i> , 2012, 120, 3856-3856.	1.4	0
14	Functional Characterization of CD4+ T-Cells in Aplastic Anemia (AA). <i>Blood</i> , 2011, 118, 1340-1340.	1.4	1
15	5-Azacytidine Specifically Depletes Regulatory T Cells (Tregs) in Myelodysplastic Syndrome (MDS) Patients. <i>Blood</i> , 2011, 118, 787-787.	1.4	2
16	Stress-activated dendritic cells interact with CD4 <sup>+</sup> T cells to elicit homeostatic memory. <i>European Journal of Immunology</i> , 2010, 40, 1628-1638.	2.9	33
17	The melanocortin receptor agonist NDP-MSH impairs the allostimulatory function of dendritic cells. <i>Immunology</i> , 2010, 129, 610-619.	4.4	9
18	The effect of allogeneic <i>in vitro</i> stimulation and <i>in vivo</i> immunization on memory CD4 <sup>+</sup> T cell APOBEC3G expression and HIV-1 infectivity. <i>European Journal of Immunology</i> , 2009, 39, 1956-1965.	2.9	18

#	ARTICLE	IF	CITATIONS
19	Mucosal immunization in macaques upregulates the innate APOBEC 3G anti-viral factor in CD4+ memory T cells. <i>Vaccine</i> , 2009, 27, 870-881.	3.8	28
20	Evidence for reduced B-cell progenitors in early (low-risk) myelodysplastic syndrome. <i>Blood</i> , 2005, 106, 2982-2991.	1.4	127
21	Molecular Signatures of Self-Renewal, Differentiation, and Lineage Choice in Multipotential Hemopoietic Progenitor Cells In Vitro. <i>Molecular and Cellular Biology</i> , 2004, 24, 741-756.	2.3	87
22	Rules for Gene Usage Inferred from a Comparison of Large-Scale Gene Expression Profiles of T and B Lymphocyte Development. <i>Journal of Immunology</i> , 2003, 170, 1339-1353.	0.8	52
23	Developmental markers of B cells are superior to those of T cells for identification of stages with distinct gene expression profiles. <i>Journal of Leukocyte Biology</i> , 2003, 74, 602-610.	3.3	2
24	Changes in Gene Expression Profiles in Developing B Cells of Murine Bone Marrow. <i>Genome Research</i> , 2002, 12, 98-111.	5.5	92
25	Profound effect of normalization on detection of differentially expressed genes in oligonucleotide microarray data analysis. <i>Genome Biology</i> , 2002, 3, research0033.1.	9.6	76
26	The VpreB protein of the surrogate light-chain can pair with some $\lambda$ heavy-chains in the absence of the $\lambda$ 5 protein. <i>European Journal of Immunology</i> , 2001, 31, 1999-2006.	2.9	18
27	Mouse pre-immunocytes as non-proliferating multipotent precursors of macrophages, interferon-producing cells, CD8 $\alpha^+$ and CD8 $\alpha^-$ dendritic cells. <i>European Journal of Immunology</i> , 2001, 31, 3403-3412.	2.9	71
28	Repertoire selection by pre-B-cell receptors and B-cell receptors, and genetic control of B-cell development from immature to mature B cells. <i>Immunological Reviews</i> , 2000, 175, 33-46.	6.0	209
29	Activated Murine B Lymphocytes and Dendritic Cells Produce a Novel CC Chemokine which Acts Selectively on Activated T Cells. <i>Journal of Experimental Medicine</i> , 1998, 188, 451-463.	8.5	145