
Gavin C Woodruff

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/7621529/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Biology and genome of a newly discovered sibling species of Caenorhabditis elegans. Nature Communications, 2018, 9, 3216.	12.8	102
2	Insights Into Species Divergence and the Evolution of Hermaphroditism From Fertile Interspecies Hybrids of Caenorhabditis Nematodes. Genetics, 2010, 186, 997-1012.	2.9	96
3	Simplification and Desexualization of Gene Expression in Self-Fertile Nematodes. Current Biology, 2012, 22, 2167-2172.	3.9	68
4	Intense Sperm-Mediated Sexual Conflict Promotes Reproductive Isolation in Caenorhabditis Nematodes. PLoS Biology, 2014, 12, e1001915.	5.6	61
5	Causes and consequences of the evolution of reproductive mode in Caenorhabditis nematodes. Trends in Genetics, 2012, 28, 213-220.	6.7	44
6	Field studies reveal a close relative of C. elegans thrives in the fresh figs of Ficus septica and disperses on its Ceratosolen pollinating wasps. BMC Ecology, 2018, 18, 26.	3.0	21
7	Mating Damages the Cuticle of C. elegans Hermaphrodites. PLoS ONE, 2014, 9, e104456.	2.5	19
8	Degradation of the Repetitive Genomic Landscape in a Close Relative of Caenorhabditis elegans. Molecular Biology and Evolution, 2020, 37, 2549-2567.	8.9	15
9	Dramatic evolution of body length due to postembryonic changes in cell size in a newly discovered close relative of <i>Caenorhabditis elegans </i> . Evolution Letters, 2018, 2, 427-441.	3.3	13
10	A large close relative of C. elegans is slow-developing but not long-lived. BMC Evolutionary Biology, 2019, 19, 74.	3.2	13
11	Teratodiplogaster variegatae n. sp. (Nematoda: Diplogastridae) isolated from the syconia of Ficus variegata Blume on Ishigaki Island, Okinawa, Japan. Nematology, 2014, 16, 1153-1166.	0.6	8
12	Diplogasteroides asiaticus n. sp. is Associated with Monochamus alternatus in Japan. Journal of Nematology, 2015, 47, 105-15.	0.9	7