
Ye-Xiong Li

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/7652864/publications.pdf Version: 2024-02-01



VE-XIONCL

#	Article	IF	CITATIONS
1	Guidelines for the Diagnosis and Treatment of Hepatocellular Carcinoma (2019 Edition). Liver Cancer, 2020, 9, 682-720.	7.7	427
2	Guidelines for Diagnosis and Treatment of Primary Liver Cancer in China (2017 Edition). Liver Cancer, 2018, 7, 235-260.	7.7	426
3	Radiotherapy As Primary Treatment for Stage IE and IIE Nasal Natural Killer/T-Cell Lymphoma. Journal of Clinical Oncology, 2006, 24, 181-189.	1.6	327
4	Modern Radiation Therapy for Extranodal Lymphomas: Field and Dose Guidelines From the International Lymphoma Radiation Oncology Group. International Journal of Radiation Oncology Biology Physics, 2015, 92, 11-31.	0.8	303
5	Hypofractionated versus conventional fractionated postmastectomy radiotherapy for patients with high-risk breast cancer: a randomised, non-inferiority, open-label, phase 3 trial. Lancet Oncology, The, 2019, 20, 352-360.	10.7	258
6	Automatic segmentation of the clinical target volume and organs at risk in the planning <scp>CT</scp> for rectal cancer using deep dilated convolutional neural networks. Medical Physics, 2017, 44, 6377-6389.	3.0	241
7	Deep Deconvolutional Neural Network for Target Segmentation of Nasopharyngeal Cancer in Planning Computed Tomography Images. Frontiers in Oncology, 2017, 7, 315.	2.8	157
8	Risk-adapted therapy for early-stage extranodal nasal-type NK/T-cell lymphoma: analysis from a multicenter study. Blood, 2015, 126, 1424-1432.	1.4	150
9	Multicenter, Randomized, Phase III Trial of Short-Term Radiotherapy Plus Chemotherapy Versus Long-Term Chemoradiotherapy in Locally Advanced Rectal Cancer (STELLAR). Journal of Clinical Oncology, 2022, 40, 1681-1692.	1.6	145
10	A feasibility study on an automated method to generate patientâ€specific dose distributions for radiotherapy using deep learning. Medical Physics, 2019, 46, 56-64.	3.0	124
11	Fully automatic and robust segmentation of the clinical target volume for radiotherapy of breast cancer using big data and deep learning. Physica Medica, 2018, 50, 13-19.	0.7	121
12	PIWI-interacting RNA-54265 is oncogenic and a potential therapeutic target in colorectal adenocarcinoma. Theranostics, 2018, 8, 5213-5230.	10.0	115
13	Primary non-hodgkin's lymphoma of the nasal cavity. , 1998, 83, 449-456.		114
14	Variable Clinical Presentations of Nasal and Waldeyer Ring Natural Killer/T-Cell Lymphoma. Clinical Cancer Research, 2009, 15, 2905-2912.	7.0	105
15	Clinical features and treatment outcome of nasal-type NK/T-cell lymphoma of Waldeyer ring. Blood, 2008, 112, 3057-3064.	1.4	102
16	Clinical implications of plasma Epstein-Barr virus DNA in early-stage extranodal nasal-type NK/T-cell lymphoma patients receiving primary radiotherapy. Blood, 2012, 120, 2003-2010.	1.4	93
17	Primary radiotherapy showed favorable outcome in treating extranodal nasal-type NK/T-cell lymphoma in children and adolescents. Blood, 2009, 114, 4771-4776.	1.4	83
18	Radiotherapy Alone With Curative Intent in Patients With Stage I Extranodal Nasal-Type NK/T-Cell Lymphoma. International Journal of Radiation Oncology Biology Physics, 2012, 82, 1809-1815.	0.8	75

#	Article	IF	CITATIONS
19	Role of Radiation Therapy in Patients With Relapsed/Refractory Diffuse Large B-Cell Lymphoma: Guidelines from the International Lymphoma Radiation Oncology Group. International Journal of Radiation Oncology Biology Physics, 2018, 100, 652-669.	0.8	71
20	Validation of nomogram-revised risk index and comparison with other models for extranodal nasal-type NK/T-cell lymphoma in the modern chemotherapy era: indication for prognostication and clinical decision-making. Leukemia, 2021, 35, 130-142.	7.2	70
21	Expert consensus on multidisciplinary therapy of colorectal cancer with lung metastases (2019) Tj ETQq1 1 0.7	784314 rgB 17.0	T /Qyerlock 1
22	Association of Improved Locoregional Control With Prolonged Survival in Early-Stage Extranodal Nasal-Type Natural Killer/T-Cell Lymphoma. JAMA Oncology, 2017, 3, 83.	7.1	64
23	Mild Toxicity and Favorable Prognosis of High–Dose and Extended Involved-Field Intensity-Modulated Radiotherapy for Patients With Early-Stage Nasal NK/T-Cell Lymphoma. International Journal of Radiation Oncology Biology Physics, 2012, 82, 1115-1121.	0.8	63
24	Failure patterns and clinical implications in early stage nasal natural killer/Tâ€cell lymphoma treated with primary radiotherapy. Cancer, 2011, 117, 5203-5211.	4.1	58
25	Hypofractionated Versus Conventional Fractionated Radiotherapy After Breast-Conserving Surgery in the Modern Treatment Era: A Multicenter, Randomized Controlled Trial From China. Journal of Clinical Oncology, 2020, 38, 3604-3614.	1.6	58
26	High-Dose and Extended-Field Intensity Modulated Radiation Therapy for Early-Stage NK/T-Cell Lymphoma ofÂWaldeyer's Ring: Dosimetric Analysis and Clinical Outcome. International Journal of Radiation Oncology Biology Physics, 2013, 87, 1086-1093.	0.8	51
27	Mapping Patterns of Ipsilateral Supraclavicular Nodal Metastases in Breast Cancer: Rethinking the Clinical Target Volume for High-risk Patients. International Journal of Radiation Oncology Biology Physics, 2015, 93, 268-276.	0.8	51
28	Survival benefit with <scp>IMRT</scp> following narrowâ€margin hepatectomy in patients with hepatocellular carcinoma close to major vessels. Liver International, 2015, 35, 2603-2610.	3.9	49
29	Phase 2 Study of Adjuvant Radiotherapy Following Narrowâ€Margin Hepatectomy in Patients With HCC. Hepatology, 2021, 74, 2595-2604.	7.3	43
30	Circulating serum microRNA-345 correlates with unfavorable pathological response to preoperative chemoradiotherapy in locally advanced rectal cancer. Oncotarget, 2016, 7, 64233-64243.	1.8	39
31	The Impact of Postoperative Conformal RadiotherapyÂafter Radical Surgery on Survival andÂRecurrence in Pathologic T3NOMO Esophageal Carcinoma: AÂPropensity Score-Matched Analysis. Journal of Thoracic Oncology, 2017, 12, 1143-1151.	1.1	35
32	Deep Learning Improved Clinical Target Volume Contouring Quality and Efficiency for Postoperative Radiation Therapy in Non-small Cell Lung Cancer. Frontiers in Oncology, 2019, 9, 1192.	2.8	35
33	First-line non–anthracycline-based chemotherapy for extranodal nasal-type NK/T-cell lymphoma: a retrospective analysis from the CLCG. Blood Advances, 2020, 4, 3141-3153.	5.2	35
34	A proposal for a new staging system for extranodal natural killer T-cell lymphoma: a multicenter study from China and Asia Lymphoma Study Group. Leukemia, 2020, 34, 2243-2248.	7.2	35
35	CNN-Based Quality Assurance for Automatic Segmentation of Breast Cancer in Radiotherapy. Frontiers in Oncology, 2020, 10, 524.	2.8	35
36	Immunophenotypic and Clinical Differences Between the Nasal and Extranasal Subtypes of Upper Aerodigestive Tract Natural Killer/T-Cell Lymphoma. International Journal of Radiation Oncology Biology Physics, 2014, 88, 806-813.	0.8	33

#	Article	IF	CITATIONS
37	Radiation-Induced Lymphopenia Predicts Poorer Prognosis in Patients With Breast Cancer: A Post Hoc Analysis of a Randomized Controlled Trial of Postmastectomy Hypofractionated Radiation Therapy. International Journal of Radiation Oncology Biology Physics, 2020, 108, 277-285.	0.8	33
38	Spread patterns of lymph nodes and the value of elective neck irradiation for esthesioneuroblastoma. Radiotherapy and Oncology, 2015, 117, 328-332.	0.6	32
39	GDP (Gemcitabine, Dexamethasone, and Cisplatin) Is Highly Effective and Well-Tolerated for Newly Diagnosed Stage IV and Relapsed/Refractory Extranodal Natural Killer/T-Cell Lymphoma, Nasal Type. Medicine (United States), 2016, 95, e2787.	1.0	32
40	Prospective evaluation of microscopic extension using whole-mount preparation in patients with hepatocellular carcinoma: Definition of clinical target volume for radiotherapy. Radiation Oncology, 2010, 5, 73.	2.7	31
41	Long-term outcomes of patients with esthesioneuroblastomas: A cohort from a single institution. Oral Oncology, 2016, 53, 48-53.	1.5	31
42	MiRâ€320b/RAD21 axis affects hepatocellular carcinoma radiosensitivity to ionizing radiation treatment through DNA damage repair signaling. Cancer Science, 2021, 112, 575-588.	3.9	31
43	Radiotherapy is essential after complete response to asparaginase-containing chemotherapy in early-stage extranodal nasal-type NK/T-cell lymphoma: A multicenter study from the China Lymphoma Collaborative Group (CLCG). Radiotherapy and Oncology, 2018, 129, 3-9.	0.6	30
44	Genome landscapes of rectal cancer before and after preoperative chemoradiotherapy. Theranostics, 2019, 9, 6856-6866.	10.0	27
45	Modern Radiation Therapy for Extranodal Nasal-Type NK/T-cell Lymphoma: Risk-Adapted Therapy, Target Volume, and Dose Guidelines from the International Lymphoma Radiation Oncology Group. International Journal of Radiation Oncology Biology Physics, 2021, 110, 1064-1081.	0.8	26
46	Response prediction and risk stratification of patients with rectal cancer after neoadjuvant therapy through an analysis of circulating tumour DNA. EBioMedicine, 2022, 78, 103945.	6.1	26
47	Immunophenotypic characteristics and clinical relevance of CD56+ and CD56â^' extranodal nasal-type natural killer/T-cell lymphoma. Leukemia and Lymphoma, 2011, 52, 417-424.	1.3	25
48	Riskâ€based, responseâ€adapted therapy for earlyâ€stage extranodal nasalâ€type <scp>NK</scp> /Tâ€cell lymphoma in the modern chemotherapy era: A China Lymphoma Collaborative Group study. American Journal of Hematology, 2020, 95, 1047-1056.	4.1	25
49	Risk-adapted survival benefit of IMRT in early-stage NKTCL: a multicenter study from the China Lymphoma Collaborative Group. Blood Advances, 2018, 2, 2369-2377.	5.2	24
50	Diffuse large B ell lymphoma. Cancer, 2009, 115, 4980-4989.	4.1	23
51	Dosimetric and Clinical Outcomes of Involved-Field Intensity-Modulated Radiotherapy After Chemotherapy for Early-Stage Hodgkin's Lymphoma With Mediastinal Involvement. International Journal of Radiation Oncology Biology Physics, 2012, 84, 210-216.	0.8	23
52	Risk-Dependent Conditional Survival and Failure Hazard After Radiotherapy for Early-Stage Extranodal Natural Killer/T-Cell Lymphoma. JAMA Network Open, 2019, 2, e190194.	5.9	23
53	Postoperative Radiotherapy in Pathological T2–3NOMO Thoracic Esophageal Squamous Cell Carcinoma: Interim Report of a Prospective, Phase III, Randomized Controlled Study. Oncologist, 2020, 25, e701-e708.	3.7	23
54	Comparison of Treatment Outcomes With Breast-conserving Surgery Plus Radiotherapy Versus Mastectomy for Patients With Stage I Breast Cancer: A Propensity Score-matched Analysis. Clinical Breast Cancer, 2018, 18, e975-e984.	2.4	21

#	Article	IF	CITATIONS
55	<p>Optimal postoperative adjuvant treatment strategy for HBV-related hepatocellular carcinoma with microvascular invasion: a propensity score analysis</p> . OncoTargets and Therapy, 2019, Volume 12, 1237-1247.	2.0	21
56	Benefit of adjuvant radiotherapy following narrow-margin hepatectomy in patients with intrahepatic cholangiocarcinoma that adhere to major vessels. Cancer Management and Research, 2018, Volume 10, 3973-3981.	1.9	20
57	Effect of primary tumor invasion on treatment and survival in extranodal nasal-type NK/T-cell lymphoma in the modern chemotherapy era: a multicenter study from the China Lymphoma Collaborative Group (CLCG). Leukemia and Lymphoma, 2019, 60, 2669-2678.	1.3	20
58	A multicenter, randomized, phase III trial of short-term radiotherapy plus chemotherapy versus long-term chemoradiotherapy in locally advanced rectal cancer (STELLAR): The final reports Journal of Clinical Oncology, 2021, 39, 3510-3510.	1.6	20
59	Prognostic factors and treatment outcomes for patients with stage II extranodal nasal-type natural killer/T-cell lymphoma of the upper aerodigestive tract. Leukemia and Lymphoma, 2014, 55, 1832-1837.	1.3	19
60	Patterns of Primary Tumor Invasion and Regional Lymph Node Spread Based on Magnetic Resonance Imaging in Early-Stage Nasal NK/T-cell Lymphoma: Implications for Clinical Target Volume Definition and Prognostic Significance. International Journal of Radiation Oncology Biology Physics, 2017, 97, 50-59.	0.8	19
61	Favorable outcome with doxorubicinâ€based chemotherapy and radiotherapy for adult patients with early stage primary systemic anaplastic largeâ€cell lymphoma. European Journal of Haematology, 2013, 90, 195-201.	2.2	18
62	Gemcitabine, dexamethasone, and cisplatin (GDP) as salvage chemotherapy for patients with relapsed or refractory peripheral T cell lymphoma—not otherwise specified. Annals of Hematology, 2017, 96, 245-251.	1.8	18
63	LncRNA and mRNA signatures associated with neoadjuvant chemoradiotherapy downstaging effects in rectal cancer. Journal of Cellular Biochemistry, 2019, 120, 5207-5217.	2.6	18
64	Association of progression-free or event-free survival with overall survival in diffuse large B-cell lymphoma after immunochemotherapy: a systematic review. Leukemia, 2020, 34, 2576-2591.	7.2	18
65	Dosimetric Evaluation and Treatment Outcome of Intensity Modulated Radiation Therapy After Doxorubicin-Based Chemotherapy for Primary Mediastinal Large B-Cell Lymphoma. International Journal of Radiation Oncology Biology Physics, 2013, 85, 1289-1295.	0.8	17
66	Efficacy and Safety of Intensity-Modulated Radiotherapy Following Transarterial Chemoembolization in Patients With Unresectable Hepatocellular Carcinoma. Medicine (United States), 2016, 95, e3789.	1.0	17
67	A propensity-score matching analysis comparing long-term survival of surgery alone and postoperative treatment for patients in node positive or stage III esophageal squamous cell carcinoma after R0 esophagectomy. Radiotherapy and Oncology, 2019, 140, 159-166.	0.6	16
68	Survival Advantage With the Addition of Radiation Therapy to Chemotherapy in Early Stage Peripheral T-Cell Lymphoma, Not Otherwise Specified. International Journal of Radiation Oncology Biology Physics, 2013, 85, 1051-1056.	0.8	15
69	Phase 2 Study of First-line Intensity Modulated Radiation Therapy Followed by Gemcitabine, Dexamethasone, and Cisplatin for High-Risk, Early Stage Extranodal Nasal-Type NK/T-Cell Lymphoma: The GREEN Study. International Journal of Radiation Oncology Biology Physics, 2018, 102, 61-70.	0.8	15
70	Use of sequential endorectal US to predict the tumor response of preoperative chemoradiotherapy in rectal cancer. Gastrointestinal Endoscopy, 2017, 85, 669-674.	1.0	14
71	Riskâ€dependent curability of radiotherapy for elderly patients with earlyâ€stage extranodal nasalâ€type NK/Tâ€cell lymphoma: A multicenter study from the China Lymphoma Collaborative Group (CLCG). Cancer Medicine, 2018, 7, 5952-5961.	2.8	14
	Upfront radiation is essential for high-risk early-stage extranodal NK/T-cell lymphoma, nasal type:		

comparison of two sequential treatment modalities combining radiotherapy and GDP (gemcitabine,) Tj ETQq0 0 0 rgBT /Overlack 10 Tf !

#	Article	IF	CITATIONS
73	Progression-free survival at 24 months and subsequent survival of patients with extranodal NK/T-cell lymphoma: a China Lymphoma Collaborative Group (CLCG) study. Leukemia, 2021, 35, 1671-1682.	7.2	14
74	Nomogram and recursive partitioning analysis to predict overall survival in patients with stage IIB-III thoracic esophageal squamous cell carcinoma after esophagectomy. Oncotarget, 2016, 7, 55211-55221.	1.8	13
75	Clinical Disparity and Favorable Prognoses for Patients With Waldeyer Ring Extranodal Nasal-type NK/T-cell Lymphoma and Diffuse Large B-cell Lymphoma. American Journal of Clinical Oncology: Cancer Clinical Trials, 2014, 37, 41-46.	1.3	12
76	Experts' consensus on intraoperative radiotherapy for pancreatic cancer. Cancer Letters, 2019, 449, 1-7.	7.2	12
77	Effectiveness of the AJCC 8th edition staging system for selecting patients with T1–2N1 breast cancer for post-mastectomy radiotherapy: a joint analysis of 1986 patients from two institutions. BMC Cancer, 2020, 20, 792.	2.6	12
78	The prognostic value of MRI-detected extramural vascular invasion (mrEMVI) for rectal cancer patients treated with neoadjuvant therapy: a meta-analysis. European Radiology, 2021, 31, 8827-8837.	4.5	12
79	Postmastectomy Radiation Therapy Based on Pathologic Nodal Status in Clinical Node-Positive Stage II to III Breast Cancer Treated with Neoadjuvant Chemotherapy. International Journal of Radiation Oncology Biology Physics, 2020, 108, 1030-1039.	0.8	11
80	Effect of age as a continuous variable on survival outcomes and treatment selection in patients with extranodal nasal-type NK/T-cell lymphoma from the China Lymphoma Collaborative Group (CLCG). Aging, 2019, 11, 8463-8473.	3.1	11
81	Phaseâ€lâ€study of postoperative radiotherapy combined with capecitabine for gastric cancer. World Journal of Gastroenterology, 2014, 20, 1067.	3.3	10
82	A prognostic nomogram for overall survival after neoadjuvant radiotherapy or chemoradiotherapy in thoracic esophageal squamous cell carcinoma: a retrospective analysis. Oncotarget, 2017, 8, 41102-41112.	1.8	10
83	Impact of Magnetic Field on Dose Distribution in MR-Guided Radiotherapy of Head and Neck Cancer. Frontiers in Oncology, 2020, 10, 1739.	2.8	10
84	Nomogram predicting survival as a selection criterion for postmastectomy radiotherapy in patients with T1 to T2 breast cancer with 1 to 3 positive lymph nodes. Cancer, 2020, 126, 3857-3866.	4.1	10
85	Accurate method for evaluating the duration of the entire radiotherapy process. Journal of Applied Clinical Medical Physics, 2020, 21, 252-258.	1.9	10
86	Neoadjuvant Chemotherapy–Guided Bladder-Sparing Treatment for Muscle-Invasive Bladder Cancer: Results of a Pilot Phase II Study. Cancer Research and Treatment, 2021, 53, 1156-1165.	3.0	10
87	Automatic segmentation of three clinical target volumes in radiotherapy using lifelong learning. Radiotherapy and Oncology, 2021, 157, 1-7.	0.6	10
88	Stageâ€dependent conditional survival and failure hazard of nonâ€metastatic nasopharyngeal carcinoma after intensityâ€modulated radiation therapy: Clinical implications for treatment strategies and surveillance. Cancer Medicine, 2021, 10, 3613-3621.	2.8	10
89	Interim analysis of postoperative chemoradiotherapy with capecitabine and oxaliplatin versus capecitabine alone for pathological stage II and III rectal cancer: a randomized multicenter phase III trial. Oncotarget, 2016, 7, 25576-25584.	1.8	10
90	Associations of Genetic Variations in Mismatch Repair Genes MSH3 and PMS1 with Acute Adverse Events and Survival in Patients with Rectal Cancer Receiving Postoperative Chemoradiotherapy. Cancer Research and Treatment, 2019, 51, 1198-1206.	3.0	10

#	Article	IF	CITATIONS
91	Association of the Cumulative Dose of Radioactive Iodine Therapy With Overall Survival in Patients With Differentiated Thyroid Cancer and Pulmonary Metastases. Frontiers in Oncology, 2019, 9, 558.	2.8	9
92	Development and external validation of a nomogram to predict four or more positive nodes in breast cancer patients with one to three positive sentinel lymph nodes. Breast, 2020, 53, 143-151.	2.2	9
93	Survival benefit with salvage radiotherapy for patients with locoregionally recurrent extranodal NK/T cell lymphoma, nasal type. Annals of Hematology, 2013, 92, 325-332.	1.8	8
94	Postoperative intensity-modulated radiation therapy provides favorable local control and low toxicities in patients with soft tissue sarcomas in the extremities and trunk wall. OncoTargets and Therapy, 2015, 8, 2843.	2.0	8
95	Clinical and prognostic differences between ALK-negative anaplastic large cell lymphoma and peripheral T cell lymphoma, not otherwise specified: a single institution experience. Annals of Hematology, 2016, 95, 1271-1280.	1.8	8
96	Intensity Modulated Radiation Therapy for Early-Stage Primary Gastric Diffuse Large B-Cell Lymphoma: Dosimetric Analysis, Clinical Outcome, and Quality of Life. International Journal of Radiation Oncology Biology Physics, 2016, 95, 712-720.	0.8	8
97	Observation of different tumor motion magnitude within liver and estimate of internal motion margins in postoperative patients with hepatocellular carcinoma. Cancer Management and Research, 2017, Volume 9, 839-848.	1.9	8
98	A feasible study on using multiplexed sensitivity-encoding to reduce geometric distortion in diffusion-weighted echo planar imaging. Magnetic Resonance Imaging, 2018, 54, 153-159.	1.8	8
99	Cerebral functional abnormalities in patients with nasopharyngeal carcinoma after radiotherapy. Chinese Medical Journal, 2019, 132, 1563-1571.	2.3	8
100	Radiomics Analysis of Fat-Saturated T2-Weighted MRI Sequences for the Prediction of Prognosis in Soft Tissue Sarcoma of the Extremities and Trunk Treated With Neoadjuvant Radiotherapy. Frontiers in Oncology, 2021, 11, 710649.	2.8	8
101	Pretreatment nutritional risk as a prognostic factor in head and neck cancer patients receiving radiotherapy or chemoradiotherapy. Asia Pacific Journal of Clinical Nutrition, 2019, 28, 223-229.	0.4	8
102	Guidelines for radiotherapy of prostate cancer (2020 edition). Precision Radiation Oncology, 2021, 5, 160-182.	1.1	8
103	Evaluation of MLC leaf transmission on IMRT treatment plan quality of patients with advanced lung cancer. Medical Dosimetry, 2018, 43, 313-318.	0.9	7
104	Trastuzumab Provides a Comparable Prognosis in Patients With HER2-Positive Breast Cancer to Those With HER2-Negative Breast Cancer: Post Hoc Analyses of a Randomized Controlled Trial of Post-Mastectomy Hypofractionated Radiotherapy. Frontiers in Oncology, 2020, 10, 605750.	2.8	7
105	The status of medical physics in radiotherapy in China. Physica Medica, 2021, 85, 147-157.	0.7	7
106	POstmastectomy radioThErapy in Node-posiTive breast cancer with or without Internal mAmmary nodaL irradiation (POTENTIAL): a study protocol for a multicenter prospective phase III randomized controlled trial. BMC Cancer, 2021, 21, 1185.	2.6	7
107	Dosimetric and Clinical Outcomes With Intensity Modulated Radiation Therapy After Chemotherapy for Patients With Early-Stage Diffuse Large B-cell Lymphoma of Waldeyer Ring. International Journal of Radiation Oncology Biology Physics, 2016, 96, 379-386.	0.8	6
108	A fast optimization approach for treatment planning of volumetric modulated arc therapy. Radiation Oncology, 2018, 13, 101.	2.7	6

#	Article	IF	CITATIONS
109	Radiotherapy plays an important role in improving the survival outcome in patients with T1–2N1M0 breast cancer – a joint analysis of 4262 real world cases from two institutions. BMC Cancer, 2020, 20, 1155.	2.6	6
110	Managing a radiotherapy center safely and efficiently using risk-adaptive strategies during coronavirus disease pandemic: Experience from national cancer center of China. Radiotherapy and Oncology, 2020, 148, 243-244.	0.6	6
111	Inhibition of EPS8L3 suppresses liver cancer progression and enhances efficacy of sorafenib treatment. Biomedicine and Pharmacotherapy, 2020, 128, 110284.	5.6	6
112	Safety and efficacy of preoperative chemoradiotherapy in fit older patients with intermediate or locally advanced rectal cancer evaluated by comprehensive geriatric assessment: A planned interim analysis of a multicenter, phase II trial. Journal of Geriatric Oncology, 2021, 12, 572-577.	1.0	6
113	A novel nomogram for predicting locoregional recurrence risk in breast cancer patients treated with neoadjuvant chemotherapy and mastectomy. Radiotherapy and Oncology, 2021, 161, 191-197.	0.6	6
114	Cost-effectiveness of postmastectomy hypofractionated radiation therapy vs conventional fractionated radiation therapy for high-risk breast cancer. Breast, 2021, 58, 72-79.	2.2	6
115	Down-staging depth score to predict outcomes in locally advanced rectal cancer achieving ypl stage after neoadjuvant chemo-radiotherapy versus de novo stage pl cohort: A propensity score-matched analysis. Chinese Journal of Cancer Research: Official Journal of China Anti-Cancer Association, Beijing Institute for Cancer Research. 2018. 30. 373-381.	2.2	6
116	First-Line Chemoradiation With or Without Chidamide (Tucidinostat) in Patients With Intermediate- and High-Risk Early-Stage Extranodal Nasal-Type Natural Killer/T-Cell Lymphoma: A Randomized Phase 2 Study in China. International Journal of Radiation Oncology Biology Physics, 2022, 113, 833-844.	0.8	6
117	Epidemiologic Study of Radiotherapy Use in China in Patients With Breast Cancer Between 1999 and 2008. Clinical Breast Cancer, 2013, 13, 47-52.	2.4	5
118	A prospective phase I study of hypo-fractionated neoadjuvant radiotherapy for locally advanced gastric cancer. BMC Cancer, 2018, 18, 803.	2.6	5
119	Preoperative versus postoperative chemo-radiotherapy for locally advanced gastric cancer: a multicenter propensity score-matched analysis. BMC Cancer, 2022, 22, 212.	2.6	5
120	Epidemiologic study of compliance to postmastectomy radiation therapy guidelines in breast cancer patients in China between 1999 and 2008. Practical Radiation Oncology, 2013, 3, 209-215.	2.1	4
121	Long-term outcomes of moderately hypofractionated radiotherapy (67.5ÂGy in 25 fractions) for prostate cancer confined to the pelvis: a single center retrospective analysis. Radiation Oncology, 2020, 15, 231.	2.7	4
122	Locoregional recurrence patterns in women with breast cancer who have not undergone post-mastectomy radiotherapy. Radiation Oncology, 2020, 15, 212.	2.7	4
123	Prognosis and Prophylactic Regional Nodal Irradiation in Breast Cancer Patients With the First Isolated Chest Wall Recurrence After Mastectomy. Frontiers in Oncology, 2020, 10, 600525.	2.8	4
124	Adjuvant treatment may benefit patients with high-risk upper rectal cancer: A nomogram and recursive partitioning analysis of 547 patients. Oncotarget, 2016, 7, 66160-66169.	1.8	4
125	Tomotherapy as an adjuvant treatment for gastroesophageal junction and stomach cancer may reduce bowel and bone marrow toxicity compared to intensity-modulated radiotherapy and volumetric-modulated arc therapy. Oncotarget, 2017, 8, 39727-39735.	1.8	4
126	Chemoradiotherapy is an alternative choice for patients with primary mediastinal seminoma. Radiation Oncology, 2022, 17, 58.	2.7	4

#	Article	IF	CITATIONS
127	Preoperative Concurrent Chemoradiotherapy Versus Neoadjuvant Chemotherapy for Locally Advanced Gastric Cancer: Phase II Randomized Study. Frontiers in Oncology, 2022, 12, 870741.	2.8	4
128	Postmastectomy chest wall radiotherapy with single low-energy electron beam: An assessment of outcome and prognostic factors. Practical Radiation Oncology, 2012, 2, 106-113.	2.1	3
129	Postoperative Capecitabine with Concurrent Intensity-Modulated Radiotherapy or Three-Dimensional Conformal Radiotherapy for Patients with Stage II and III Rectal Cancer. PLoS ONE, 2015, 10, e0124601.	2.5	3
130	Hypofractionated intensity-modulated radiation therapy for prostate cancer confined to the pelvis: analysis of efficacy and late toxicity. Journal of Radiation Oncology, 2015, 4, 95-101.	0.7	3
131	Associations of Genetic Variations in MicroRNA Seed Regions With Acute Adverse Events and Survival in Patients With Rectal Cancer Receiving Postoperative Chemoradiation Therapy. International Journal of Radiation Oncology Biology Physics, 2018, 100, 1026-1033.	0.8	3
132	A study of the dosimetric characteristics between different fixed-field IMRT and VMAT in early-stage primary mediastinal B-cell lymphoma. Medical Dosimetry, 2018, 43, 91-99.	0.9	3
133	A Phase II Trial of Concurrent Temozolomide and Hypofractionated Stereotactic Radiotherapy for Complex Brain Metastases. Oncologist, 2019, 24, e914-e920.	3.7	3
134	Outcomes after hypofractionated stereotactic radiotherapy for colorectal cancer oligometastases. Journal of Surgical Oncology, 2019, 119, 532-538.	1.7	3
135	Risk stratification for prediction of locoregional recurrence in patients with pathologic T1–2N0 breast cancer after mastectomy. BMC Cancer, 2020, 20, 1132.	2.6	3
136	Therapeutic response and long-term outcome of differentiated thyroid cancer with pulmonary metastases treated by radioiodine therapy. Oncotarget, 2017, 8, 92715-92726.	1.8	3
137	The Sequence of Intracranial Radiotherapy and Systemic Treatment With Tyrosine Kinase Inhibitors for Gene-Driven Non-Small Cell Lung Cancer Brain Metastases in the Targeted Treatment Era: A 10-Year Single-Center Experience. Frontiers in Oncology, 2021, 11, 732883.	2.8	3
138	Survey on the Use of Radiotherapy to Treat Early Breast Cancer following Breast-conserving Surgery in China. Tumori, 2014, 100, 512-517.	1.1	2
139	Clinicopathological characteristics and treatment outcomes of Chinese patients with genitourinary embryonal rhabdomyosarcoma. World Journal of Surgical Oncology, 2015, 13, 190.	1.9	2
140	Influence of tumor location on the intensity-modulated radiation therapy plan of helical tomotherapy. Medical Dosimetry, 2017, 42, 334-340.	0.9	2
141	Exploring correlation information for image compression of four-dimensional computed tomography. Quantitative Imaging in Medicine and Surgery, 2019, 9, 1270-1277.	2.0	2
142	<p>Hypofractionated Radiotherapy for 35 Patients with Adrenal Metastases: A Single-Institution Experience</p> . Cancer Management and Research, 2020, Volume 12, 11563-11571.	1.9	2
143	Timing of Chemotherapy and Radiotherapy Following Breast-Conserving Surgery for Early-Stage Breast Cancer: A Retrospective Analysis. Frontiers in Oncology, 2020, 10, 571390.	2.8	2
144	Quality of Life After Partial or Whole-Breast Irradiation in Breast-Conserving Therapy for Low-Risk Breast Cancer: 1-Year Results of a Phase 2 Randomized Controlled Trial. Frontiers in Oncology, 2021, 11, 738318.	2.8	2

#	Article	IF	CITATIONS
145	Addition of rituximab is not associated with survival benefit compared with CHOP alone for patients with stage I diffuse large B-cell lymphoma. Chinese Journal of Cancer Research: Official Journal of China Anti-Cancer Association, Beijing Institute for Cancer Research, 2015, 27, 516-23.	2.2	2
146	Postoperative Chemoradiotherapy With Capecitabine and Oxaliplatin vs Capecitabine for Stage II to III Rectal Cancer. JAMA Network Open, 2021, 4, e2136116.	5.9	2
147	Effect of postmastectomy radiotherapy on pT1-2N1 breast cancer patients with different molecular subtypes. Breast, 2022, 61, 108-117.	2.2	2
148	Long-Term Outcome of Centrally Located Hepatocellular Carcinomas Treated by Radical Resection Combined With Intraoperative Electron Radiotherapy (IOERT). Frontiers in Oncology, 2022, 12, 773301.	2.8	2
149	Variability of Target Volumes and Organs at Risk Delineation in Breast Cancer Radiation Therapy: Quality Assurance Results of the Pretrial Benchmark Case for the POTENTIAL Trial. Practical Radiation Oncology, 2022, , .	2.1	2
150	Comparison analysis of first-line asparaginase- versus non-asparaginase-based regimens for early-stage extranodal NK/T-cell lymphoma. Annals of Hematology, 2022, 101, 2021-2034.	1.8	2
151	Patients with pathological stage N2 rectal cancer treated with early adjuvant chemotherapy have a lower treatment failure rate. BMC Cancer, 2017, 17, 182.	2.6	1
152	Comparison of 2 methods for prediction of liver dosimetric indices in hepatocellular cancer IMRT planning. Medical Dosimetry, 2019, 44, e80-e85.	0.9	1
153	Abnormal pretreatment coagulation factor levels correlate with poor prognosis in patients with early-stage extranodal nasal-type natural/killer T cell lymphoma. Annals of Hematology, 2020, 99, 1303-1309.	1.8	1
154	Down-staging depth score could be a survival predictor for locally advanced gastric cancer patients after preoperative chemoradiotherapy. Chinese Journal of Cancer Research: Official Journal of China Anti-Cancer Association, Beijing Institute for Cancer Research, 2021, 33, 447-456.	2.2	1
155	Efficacy and toxicity of capecitabine combined with intensity-modulated radiotherapy after D1/D2 lymph node dissection in patients with gastric cancer. World Journal of Gastrointestinal Oncology, 2021, 13, 1532-1543.	2.0	1
156	Possible contribution of IMRT in postoperative radiochemotherapy for rectal cancer: analysis on 1798 patients by prediction model. Oncotarget, 2016, 7, 46536-46544.	1.8	1
157	Mapping of Level I Axillary Lymph Nodes in Patients with Newly Diagnosed Breast Cancer: Optimal Target Delineation and Treatment Techniques for Breast and Level I Axilla Irradiation. Practical Radiation Oncology, 2022, 12, 487-495.	2.1	1
158	Changes in postmastectomy radiotherapy targets in breast cancer in China between 1999 and 2008: an epidemiological study. Journal of Radiation Oncology, 2013, 2, 293-301.	0.7	0
159	Moderately hypofractionated breast radiation therapy: is more evidence needed? – Authors' reply. Lancet Oncology, The, 2019, 20, e227.	10.7	0
160	Four-Dimensional Cone-Beam Computed Tomography Image Compression Using Video Encoder for Radiotherapy. Journal of Digital Imaging, 2020, 33, 1292-1300.	2.9	0
161	A proposal for a new staging system for extranodal natural killer T-cell lymphoma, nasal type, to predict the treatment strategy: A multicentre study from the Chinese Southwest Oncology Group and Asia Lymphoma Study Group Journal of Clinical Oncology, 2018, 36, 7552-7552.	1.6	0
162	Risk-adapted therapy for advanced-stage natural killer/T-cell lymphoma: An analysis from the China Lymphoma Collaborative Group Study Journal of Clinical Oncology, 2020, 38, e20041-e20041.	1.6	0

#	Article	IF	CITATIONS
163	Development and Validation of an MRI-Based Nomogram Model for Predicting Disease-Free Survival in Locally Advanced Rectal Cancer Treated With Neoadjuvant Radiotherapy. Frontiers in Oncology, 2021, 11, 784156.	2.8	0
164	Progression-Free Survival and Time to Progression as Potential Surrogate Endpoints for Overall Survival in Chemoradiotherapy Trials in Limited-Stage Small-Cell Lung Cancer: A Systematic Review and Meta-Analysis. Frontiers in Oncology, 2022, 12, 810580.	2.8	0