

Stuart E Parsons

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7672204/publications.pdf>

Version: 2024-02-01

68
papers

2,187
citations

218677

26
h-index

254184

43
g-index

72
all docs

72
docs citations

72
times ranked

2240
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 1 | Thermal energetics of male courtship song in a lek-breeding bat. <i>Behavioral Ecology and Sociobiology</i> , 2022, 76, 1. | 1.4 | 2 |
| 2 | Antennal morphology and microsensory architecture of the New Zealand magpie moth, <i>Nyctemera annulata</i> (Lepidoptera: Erebidae): diversity, distribution and dimorphism. <i>Austral Entomology</i> , 2018, 57, 303-323. | 1.4 | 7 |
| 3 | Spatiotemporal and demographic variation in the diet of New Zealand lesser short-tailed bats (<i>Mystacina tuberculata</i>). <i>Ecology and Evolution</i> , 2018, 8, 7599-7610. | 1.9 | 17 |
| 4 | The high-output singing displays of a lekking bat encode information on body size and individual identity. <i>Behavioral Ecology and Sociobiology</i> , 2018, 72, 1. | 1.4 | 59 |
| 5 | Bat detective” Deep learning tools for bat acoustic signal detection. <i>PLoS Computational Biology</i> , 2018, 14, e1005995. | 3.2 | 128 |
| 6 | Geographic patterns of song variation reveal timing of song acquisition in a wild avian population. <i>Behavioral Ecology</i> , 2017, 28, 1085-1092. | 2.2 | 10 |
| 7 | Positive emotional contagion in a New Zealand parrot. <i>Current Biology</i> , 2017, 27, R213-R214. | 3.9 | 47 |
| 8 | Cold and alone? Roost choice and season affect torpor patterns in lesser short-tailed bats. <i>Oecologia</i> , 2017, 183, 1-8. | 2.0 | 24 |
| 9 | Audiogram of the kea parrot, <i>Nestor notabilis</i> . <i>Journal of the Acoustical Society of America</i> , 2016, 140, 3739-3744. | 1.1 | 7 |
| 10 | Integration over song classification replicates: Song variant analysis in the hihi. <i>Journal of the Acoustical Society of America</i> , 2015, 137, 2542-2551. | 1.1 | 18 |
| 11 | Adoption of alternative habitats by a threatened, obligate forest-dwelling bat in a fragmented landscape. <i>Journal of Mammalogy</i> , 2015, 96, 927-937. | 1.3 | 9 |
| 12 | Females as mobile resources: communal roosts promote the adoption of lek breeding in a temperate bat. <i>Behavioral Ecology</i> , 2015, 26, 1156-1163. | 2.2 | 13 |
| 13 | Retinal Anatomy of the New Zealand Kiwi: Structural Traits Consistent With Their Nocturnal Behavior. <i>Anatomical Record</i> , 2015, 298, 771-779. | 1.4 | 12 |
| 14 | Effects of Clear-Fell Harvest on Bat Home Range. <i>PLoS ONE</i> , 2014, 9, e86163. | 2.5 | 10 |
| 15 | Temporal and spatial distribution and habitat associations of an urban population of New Zealand long-tailed bats (<i>Chalinobus tuberculatus</i>). <i>New Zealand Journal of Zoology</i> , 2014, 41, 285-295. | 1.1 | 12 |
| 16 | Anatomical Specializations for Enhanced Olfactory Sensitivity in Kiwi, <i>Apteryx mantelli</i> . <i>Brain, Behavior and Evolution</i> , 2014, 84, 214-226. | 1.7 | 27 |
| 17 | Competition for pollination by the lesser short-tailed bat and its influence on the flowering phenology of some New Zealand endemics. <i>Journal of Zoology</i> , 2014, 293, 281-288. | 1.7 | 9 |
| 18 | Is lek breeding rare in bats?. <i>Journal of Zoology</i> , 2013, 291, 3-11. | 1.7 | 24 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 19 | Impacts of wind energy developments on wildlife: a southern hemisphere perspective. <i>New Zealand Journal of Zoology</i> , 2013, 40, 1-4. | 1.1 | 9 |
| 20 | The Anatomy of the bill Tip of Kiwi and Associated Somatosensory Regions of the Brain: Comparisons with Shorebirds. <i>PLoS ONE</i> , 2013, 8, e80036. | 2.5 | 59 |
| 21 | Vocal repertoire of the New Zealand kea parrot <i>Nestor notabilis</i> . <i>Environmental Epigenetics</i> , 2012, 58, 727-740. | 1.8 | 25 |
| 22 | A continental-scale tool for acoustic identification of European bats. <i>Journal of Applied Ecology</i> , 2012, 49, 1064-1074. | 4.0 | 144 |
| 23 | Inner-Ear Morphology of the New Zealand Kiwi (<i>Apteryx mantelli</i>) Suggests High-Frequency Specialization. <i>JARO - Journal of the Association for Research in Otolaryngology</i> , 2012, 13, 629-639. | 1.8 | 9 |
| 24 | Morphometric Analysis of Telencephalic Structure in a Variety of Neognath and Paleognath Bird Species Reveals Regional Differences Associated with Specific Behavioral Traits. <i>Brain, Behavior and Evolution</i> , 2012, 80, 181-195. | 1.7 | 27 |
| 25 | Discovery of a Lipid Synthesising Organ in the Auditory System of an Insect. <i>PLoS ONE</i> , 2012, 7, e51486. | 2.5 | 9 |
| 26 | Sex-Specific Roost Selection by Bats in Clearfell Harvested Plantation Forest: Improved Knowledge Advises Management. <i>Acta Chiropterologica</i> , 2011, 13, 373-383. | 0.6 | 11 |
| 27 | Mechanical filtering for narrow-band hearing in the weta. <i>Journal of Experimental Biology</i> , 2011, 214, 778-785. | 1.7 | 8 |
| 28 | Home range and habitat selection by a threatened bat in exotic plantation forest. <i>Forest Ecology and Management</i> , 2011, 262, 845-852. | 3.2 | 19 |
| 29 | Evidence for an Auditory Fovea in the New Zealand Kiwi (<i>Apteryx mantelli</i>). <i>PLoS ONE</i> , 2011, 6, e23771. | 2.5 | 42 |
| 30 | Seasonal occurrence and distribution of Bryde's whales in the Hauraki Gulf, New Zealand. <i>Marine Mammal Science</i> , 2011, 27, E253. | 1.8 | 27 |
| 31 | Bat colony size reduction coincides with clear-fell harvest operations and high rates of roost loss in plantation forest. <i>Biodiversity and Conservation</i> , 2011, 20, 3537-3548. | 2.6 | 19 |
| 32 | The conservation status of New Zealand bats, 2009. <i>New Zealand Journal of Zoology</i> , 2010, 37, 297-311. | 1.1 | 24 |
| 33 | Plantation forests are used by the lesser short-tailed bat, <i>Mystacina tuberculata rhyacobia</i> . <i>New Zealand Journal of Zoology</i> , 2010, 37, 13-17. | 1.1 | 5 |
| 34 | The importance of exotic plantation forest for the New Zealand long-tailed bat (<i>Chalinolobus</i>). <i>Overlock 10, 11, 12, 13, 14, 15</i> | 1.1 | 13 |
| 35 | Echolocation call production during aerial and terrestrial locomotion by New Zealand's enigmatic lesser short-tailed bat, <i>Mystacina tuberculata</i> . <i>Journal of Experimental Biology</i> , 2010, 213, 551-557. | 1.7 | 12 |
| 36 | Variation in the abundance of ectoparasitic mites of flat-headed bats. <i>Journal of Mammalogy</i> , 2010, 91, 136-143. | 1.3 | 14 |

| # | ARTICLE | IF | CITATIONS |
|----|--|------|-----------|
| 37 | Classification of Echolocation Calls from 14 Species of Bat by Support Vector Machines and Ensembles of Neural Networks. <i>Algorithms</i> , 2009, 2, 907-924. | 2.1 | 54 |
| 38 | Recent Surveys of Bats (Mammalia: Chiroptera) from China. I. Rhinolophidae and Hipposideridae. <i>Acta Chiropterologica</i> , 2009, 11, 71-88. | 0.6 | 46 |
| 39 | Assessment of the short-term success of a translocation of lesser short-tailed bats <i>Mystacina tuberculata</i> . <i>Endangered Species Research</i> , 2009, 8, 33-39. | 2.4 | 14 |
| 40 | Translocation of bats as a conservation strategy: previous attempts and potential problems. <i>Endangered Species Research</i> , 2009, 8, 25-31. | 2.4 | 13 |
| 41 | MRI of postmortem specimens of endangered species for comparative brain anatomy. <i>Nature Protocols</i> , 2008, 3, 597-605. | 12.0 | 30 |
| 42 | Human vs. machine: identification of bat species from their echolocation calls by humans and by artificial neural networks. <i>Canadian Journal of Zoology</i> , 2008, 86, 371-377. | 1.0 | 58 |
| 43 | Wing morphology, echolocation calls, diet and emergence time of black-bearded tomb bats (<i>Taphozous melanopogon</i> , Emballonuridae) from southwest China. <i>Acta Chiropterologica</i> , 2008, 10, 51-59. | 0.6 | 10 |
| 44 | VOCALIZATIONS OF THE NORTH ISLAND BROWN KIWI (<i>APTERYX MANTELLI</i>). <i>Auk</i> , 2008, 125, 326-335. | 1.4 | 22 |
| 45 | Evolution of Brain Size in the Palaeognath Lineage, with an Emphasis on New Zealand Ratites. <i>Brain, Behavior and Evolution</i> , 2008, 71, 87-99. | 1.7 | 45 |
| 46 | The potential availability of roosting sites for lesser short-tailed bats (<i>Mystacina tuberculata</i>) on Kapiti Island, New Zealand: Implications for a translocation. <i>New Zealand Journal of Zoology</i> , 2007, 34, 219-226. | 1.1 | 2 |
| 47 | Temporal and spatial patterns of seed dispersal of <i>Musa acuminata</i> by <i>Cynopterus sphinx</i> . <i>Acta Chiropterologica</i> , 2007, 9, 229-235. | 0.6 | 14 |
| 48 | Evidence of homing following translocation of long-tailed bats (<i>Chalinolobus tuberculatus</i>) at Grand Canyon Cave, New Zealand. <i>New Zealand Journal of Zoology</i> , 2007, 34, 239-246. | 1.1 | 11 |
| 49 | Bats respond to polarity of a magnetic field. <i>Proceedings of the Royal Society B: Biological Sciences</i> , 2007, 274, 2901-2905. | 2.6 | 75 |
| 50 | Kiwi Forego Vision in the Guidance of Their Nocturnal Activities. <i>PLoS ONE</i> , 2007, 2, e198. | 2.5 | 91 |
| 51 | Fruit-feeding behaviour and use of olfactory cues by the fruit bat <i>Rousettus leschenaulti</i> : an experimental study. <i>Acta Theriologica</i> , 2007, 52, 285-290. | 1.1 | 7 |
| 52 | PHYLOGENETICS OF SMALL HORSESHOE BATS FROM EAST ASIA BASED ON MITOCHONDRIAL DNA SEQUENCE VARIATION. <i>Journal of Mammalogy</i> , 2006, 87, 1234-1240. | 1.3 | 56 |
| 53 | Echolocation calls, wing shape, diet and phylogenetic diagnosis of the endemic Chinese bat <i>Myotis pequinus</i> . <i>Acta Chiropterologica</i> , 2006, 8, 451-463. | 0.6 | 20 |
| 54 | Terrestrial locomotion of the New Zealand short-tailed bat <i>Mystacina tuberculata</i> and the common vampire bat <i>Desmodus rotundus</i> . <i>Journal of Experimental Biology</i> , 2006, 209, 1725-1736. | 1.7 | 67 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 55 | Genetic divergence and echolocation call frequency in cryptic species of <i>Hipposideros larvatus</i> s.l. (Chiroptera: Hipposideridae) from the Indo-Malayan region. <i>Biological Journal of the Linnean Society</i> , 2006, 88, 119-130. | 1.6 | 93 |
| 56 | Echolocation call intensity in the aerial hawking bat <i>Eptesicus bottae</i> (Vespertilionidae) studied using stereo videogrammetry. <i>Journal of Experimental Biology</i> , 2005, 208, 1321-1327. | 1.7 | 103 |
| 57 | Development of vocalizations in the flat-headed bats, <i>Tylonycteris pachypus</i> and <i>T. robustula</i> (Chiroptera: Vespertilionidae). <i>Acta Chiropterologica</i> , 2005, 7, 91-99. | 0.6 | 20 |
| 58 | Echolocation Calls and Wing Morphology of Bats from the West Indies. <i>Acta Chiropterologica</i> , 2004, 6, 75-90. | 0.6 | 48 |
| 59 | The influence of flight speed on the ranging performance of bats using frequency modulated echolocation pulses. <i>Journal of the Acoustical Society of America</i> , 2003, 113, 617-628. | 1.1 | 24 |
| 60 | Female greater wax moths reduce sexual display behavior in relation to the potential risk of predation by echolocating bats. <i>Behavioral Ecology</i> , 2002, 13, 375-380. | 2.2 | 58 |
| 61 | Effects of Different Surfaces on the Perception of Prey-Generated Noise by the Indian False Vampire Bat <i>Megaderma lyra</i> . <i>Acta Chiropterologica</i> , 2002, 4, 25-32. | 0.6 | 6 |
| 62 | Identification of New Zealand bats (<i>Chalinolobus tuberculatus</i> and <i>Mystacina tuberculata</i>) in flight from analysis of echolocation calls by artificial neural networks. <i>Journal of Zoology</i> , 2001, 253, 447-456. | 1.7 | 42 |
| 63 | ADVANTAGES AND DISADVANTAGES OF TECHNIQUES FOR TRANSFORMING AND ANALYZING CHIROPTERAN ECHOLOCATION CALLS. <i>Journal of Mammalogy</i> , 2000, 81, 927-938. | 1.3 | 54 |
| 64 | The Long and Short of It: Branch Lengths and the Problem of Placing the New Zealand Short-Tailed Bat, <i>Mystacina</i> . <i>Molecular Phylogenetics and Evolution</i> , 1999, 13, 405-416. | 2.7 | 42 |
| 65 | The effect of recording situation on the echolocation calls of the New Zealand lesser short-tailed bat (<i>Mystacina tuberculata</i> Gray). <i>New Zealand Journal of Zoology</i> , 1998, 25, 147-156. | 1.1 | 12 |
| 66 | Search-phase echolocation calls of the New Zealand lesser short-tailed bat (<i>Mystacina</i>) <i>Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 307 Td</i> . <i>Zoology</i> , 1997, 75, 1487-1494. | 1.0 | 35 |
| 67 | A COMPARISON OF THE PERFORMANCE OF A BRAND OF BROAD-BAND AND SEVERAL BRANDS OF NARROW-BAND BAT DETECTORS IN TWO DIFFERENT HABITAT TYPES. <i>Bioacoustics</i> , 1996, 7, 33-43. | 1.7 | 23 |
| 68 | Stressful summers? Torpor expression differs between high- and low-latitude populations of bats. <i>Journal of Mammalogy</i> , 0, , . | 1.3 | 8 |