
## Sang-We Kim

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/7691411/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Afatinib versus gefitinib as first-line treatment of patients with EGFR mutation-positive non-small-cell lung cancer (LUX-Lung 7): a phase 2B, open-label, randomised controlled trial. Lancet Oncology, The, 2016, 17, 577-589.	10.7	950
2	Brigatinib in Patients With Crizotinib-Refractory Anaplastic Lymphoma Kinase–Positive Non–Small-Cell Lung Cancer: A Randomized, Multicenter Phase II Trial. Journal of Clinical Oncology, 2017, 35, 2490-2498.	1.6	506
3	Osimertinib in Pretreated T790M-Positive Advanced Non–Small-Cell Lung Cancer: AURA Study Phase II Extension Component. Journal of Clinical Oncology, 2017, 35, 1288-1296.	1.6	470
4	Gefitinib plus chemotherapy versus placebo plus chemotherapy in EGFR-mutation-positive non-small-cell lung cancer after progression on first-line gefitinib (IMPRESS): a phase 3 randomised trial. Lancet Oncology, The, 2015, 16, 990-998.	10.7	353
5	Amivantamab in EGFR Exon 20 Insertion–Mutated Non–Small-Cell Lung Cancer Progressing on Platinum Chemotherapy: Initial Results From the CHRYSALIS Phase I Study. Journal of Clinical Oncology, 2021, 39, 3391-3402.	1.6	320
6	Entrectinib in ROS1 fusion-positive non-small-cell lung cancer: integrated analysis of three phase 1–2 trials. Lancet Oncology, The, 2020, 21, 261-270.	10.7	303
7	Osimertinib plus savolitinib in patients with EGFR mutation-positive, MET-amplified, non-small-cell lung cancer after progression on EGFR tyrosine kinase inhibitors: interim results from a multicentre, open-label, phase 1b study. Lancet Oncology, The, 2020, 21, 373-386.	10.7	300
8	Osimertinib in Patients With Epidermal Growth Factor Receptor Mutation–Positive Non–Small-Cell Lung Cancer and Leptomeningeal Metastases: The BLOOM Study. Journal of Clinical Oncology, 2020, 38, 538-547.	1.6	221
9	Multinational Randomized Phase III Trial With or Without Consolidation Chemotherapy Using Docetaxel and Cisplatin After Concurrent Chemoradiation in Inoperable Stage III Non–Small-Cell Lung Cancer: KCSG-LU05-04. Journal of Clinical Oncology, 2015, 33, 2660-2666.	1.6	215
10	First-Line Erlotinib Therapy Until and Beyond Response Evaluation Criteria in Solid Tumors Progression in Asian Patients With Epidermal Growth Factor Receptor Mutation–Positive Non–Small-Cell Lung Cancer. JAMA Oncology, 2016, 2, 305.	7.1	201
11	Alectinib versus crizotinib in untreated Asian patients with anaplastic lymphoma kinase-positive non-small-cell lung cancer (ALESIA): a randomised phase 3 study. Lancet Respiratory Medicine,the, 2019, 7, 437-446.	10.7	192
12	First-Line Nivolumab Plus Ipilimumab in Advanced NSCLC: 4-Year Outcomes From the Randomized, Open-Label, Phase 3 CheckMate 227 Part 1 Trial. Journal of Thoracic Oncology, 2022, 17, 289-308.	1.1	173
13	Tepotinib plus gefitinib in patients with EGFR-mutant non-small-cell lung cancer with MET overexpression or MET amplification and acquired resistance to previous EGFR inhibitor (INSIGHT) Tj ETQq1 1 0.7 8. 1132-1143.	′84314 rg' 10.7	BT/Overlock
14	o: 11921149. Treatment Outcomes and Safety of Mobocertinib in Platinum-Pretreated Patients With <i>EGFR</i> Exon 20 Insertion–Positive Metastatic Non–Small Cell Lung Cancer. JAMA Oncology, 2021, 7, e214761.	7.1	160
15	ASCEND-8: A Randomized Phase 1 Study of Ceritinib, 450 mg or 600 mg, Taken with a Low-Fat Meal versus 750 mg in Fasted State in Patients with Anaplastic Lymphoma Kinase (ALK)-Rearranged Metastatic Non–Small Cell Lung Cancer (NSCLC). Journal of Thoracic Oncology, 2017, 12, 1357-1367.	1.1	144
16	Efficacy and Safety of Patritumab Deruxtecan (HER3-DXd) in EGFR Inhibitor–Resistant, <i>EGFR</i> -Mutated Non–Small Cell Lung Cancer. Cancer Discovery, 2022, 12, 74-89.	9.4	133
17	Osimertinib Western and Asian clinical pharmacokinetics in patients and healthy volunteers: implications for formulation, dose, and dosing frequency in pivotal clinical studies. Cancer Chemotherapy and Pharmacology, 2016, 77, 767-776.	2.3	118
18	Brigatinib in Crizotinib-Refractory ALK+ NSCLC: 2-Year Follow-up on Systemic and Intracranial Outcomes in the Phase 2 ALTA Trial. Journal of Thoracic Oncology, 2020, 15, 404-415.	1.1	102

#	Article	IF	CITATIONS
19	Epidermal growth factor receptor mutation analysis in tissue and plasma from the AURA3 trial: Osimertinib versus platinumâ€pemetrexed for T790M mutationâ€positive advanced non–small cell lung cancer. Cancer, 2020, 126, 373-380.	4.1	95
20	Activity and safety of AZD3759 in EGFR-mutant non-small-cell lung cancer with CNS metastases (BLOOM): a phase 1, open-label, dose-escalation and dose-expansion study. Lancet Respiratory Medicine,the, 2017, 5, 891-902.	10.7	92
21	Lazertinib in patients with EGFR mutation-positive advanced non-small-cell lung cancer: results from the dose escalation and dose expansion parts of a first-in-human, open-label, multicentre, phase 1–2 study. Lancet Oncology, The, 2019, 20, 1681-1690.	10.7	92
22	Postoperative Chemotherapy Use and Outcomes From ADAURA: Osimertinib as Adjuvant Therapy for Resected EGFR-Mutated NSCLC. Journal of Thoracic Oncology, 2022, 17, 423-433.	1.1	89
23	Development of thyroid dysfunction is associated with clinical response to PD-1 blockade treatment in patients with advanced non-small cell lung cancer. Oncolmmunology, 2018, 7, e1375642.	4.6	83
24	Clinical activity of the mutant-selective EGFR inhibitor AZD9291 in patients (pts) with EGFR inhibitor–resistant non-small cell lung cancer (NSCLC) Journal of Clinical Oncology, 2014, 32, 8009-8009.	1.6	81
25	JNJ-61186372 (JNJ-372), an EGFR-cMet bispecific antibody, in EGFR-driven advanced non-small cell lung cancer (NSCLC) Journal of Clinical Oncology, 2019, 37, 9009-9009.	1.6	74
26	Clinical Activity, Tolerability, and Long-Term Follow-Up of Durvalumab in Patients With Advanced NSCLC. Journal of Thoracic Oncology, 2019, 14, 1794-1806.	1.1	69
27	Osimertinib for patients (pts) with leptomeningeal metastases (LM) from EGFR-mutant non-small cell lung cancer (NSCLC): Updated results from the BLOOM study Journal of Clinical Oncology, 2017, 35, 2020-2020.	1.6	63
28	Phase 2 Study of the HSP-90 Inhibitor AUY922 in Previously Treated and Molecularly Defined Patients with Advanced Non–Small Cell Lung Cancer. Journal of Thoracic Oncology, 2018, 13, 576-584.	1.1	62
29	Efficacy and Safety of Ceritinib (450 mg/d or 600 mg/d) With Food Versus 750-mg/d Fasted in Patients With ALK Receptor Tyrosine Kinase (ALK)–Positive NSCLC: Primary Efficacy Results From the ASCEND-8 Study. Journal of Thoracic Oncology, 2019, 14, 1255-1265.	1.1	59
30	Osimertinib activity in patients (pts) with leptomeningeal (LM) disease from non-small cell lung cancer (NSCLC): Updated results from BLOOM, a phase I study Journal of Clinical Oncology, 2016, 34, 9002-9002.	1.6	59
31	Osimertinib as adjuvant therapy in patients (pts) with stage IB–IIIA EGFR mutation positive (EGFRm) NSCLC after complete tumor resection: ADAURA Journal of Clinical Oncology, 2020, 38, LBA5-LBA5.	1.6	56
32	Amivantamab in combination with lazertinib for the treatment of osimertinib-relapsed, chemotherapy-naìve EGFR mutant (EGFRm) non-small cell lung cancer (NSCLC) and potential biomarkers for response Journal of Clinical Oncology, 2021, 39, 9006-9006.	1.6	55
33	A Phase 1 study of gefitinib combined with durvalumab in EGFR TKI-naive patients with EGFR mutation-positive locally advanced/metastatic non-small-cell lung cancer. British Journal of Cancer, 2021, 124, 383-390.	6.4	54
34	Amivantamab (JNJ-61186372), an anti-EGFR-MET bispecific antibody, in patients with EGFR exon 20 insertion (exon20ins)-mutated non-small cell lung cancer (NSCLC) Journal of Clinical Oncology, 2020, 38, 9512-9512.	1.6	54
35	Phase II evaluation of LY2603618, a first-generation CHK1 inhibitor, in combination with pemetrexed in patients with advanced or metastatic non-small cell lung cancer. Investigational New Drugs, 2016, 34, 625-635.	2.6	52
36	Avelumab (anti–PD-L1) in combination with crizotinib or lorlatinib in patients with previously treated advanced NSCLC: Phase 1b results from JAVELIN Lung 101 Journal of Clinical Oncology, 2018, 36, 9008-9008.	1.6	47

#	Article	IF	CITATIONS
37	A Phase 1/2 Study of Lazertinib 240 mg in Patients With Advanced EGFR T790M-Positive NSCLC After Previous EGFR Tyrosine Kinase Inhibitors. Journal of Thoracic Oncology, 2022, 17, 558-567.	1.1	43
38	ASTRIS: a global real-world study of osimertinib in >3000 patients with <i>EGFR</i> T790M positive non-small-cell lung cancer. Future Oncology, 2019, 15, 3003-3014.	2.4	42
39	Nivolumab + ipilimumab versus platinum-doublet chemotherapy as first-line treatment for advanced non-small cell lung cancer: Three-year update from CheckMate 227 Part 1 Journal of Clinical Oncology, 2020, 38, 9500-9500.	1.6	42
40	Survival Benefit of Pemetrexed in Lung Adenocarcinoma Patients With Anaplastic Lymphoma Kinase Gene Rearrangements. Clinical Lung Cancer, 2015, 16, e83-e89.	2.6	40
41	The HSP90 inhibitor, NVP-AUY922, sensitizes KRAS-mutant non-small cell lung cancer with intrinsic resistance to MEK inhibitor, trametinib. Cancer Letters, 2016, 372, 75-81.	7.2	35
42	A Phase II Study of Poziotinib in Patients with Epidermal Growth Factor Receptor ( <i>EGFR</i> )-Mutant Lung Adenocarcinoma Who Have Acquired Resistance to EGFR–Tyrosine Kinase Inhibitors. Cancer Research and Treatment, 2017, 49, 10-19.	3.0	35
43	Safety and clinical activity of durvalumab (MEDI4736), an anti-PD-L1 antibody, in treatment-naÃ⁻ve patients with advanced non‒small-cell lung cancer Journal of Clinical Oncology, 2016, 34, 9029-9029.	1.6	32
44	Multiple resistant factors in lung cancer with primary resistance to EGFR-TK inhibitors confer poor survival. Lung Cancer, 2015, 88, 139-146.	2.0	31
45	First-line afatinib vs gefitinib for patients with EGFR mutation-positive NSCLC (LUX-Lung 7): impact of afatinib dose adjustment and analysis of mode of initial progression for patients who continued treatment beyond progression. Journal of Cancer Research and Clinical Oncology, 2019, 145, 1569-1579.	2.5	31
46	Phase Ib Study of Lumretuzumab Plus Cetuximab or Erlotinib in Solid Tumor Patients and Evaluation of HER3 and Heregulin as Potential Biomarkers of Clinical Activity. Clinical Cancer Research, 2017, 23, 5406-5415.	7.0	29
47	Safety and efficacy of the anti-CD73 monoclonal antibody (mAb) oleclumab ± durvalumab in patients (pts) with advanced colorectal cancer (CRC), pancreatic ductal adenocarcinoma (PDAC), or EGFR-mutant non-small cell lung cancer (EGFRm NSCLC) Journal of Clinical Oncology, 2021, 39, 9047-9047.	1.6	28
48	Phase II study of the HSP90 inhibitor AUY922 in patients with previously treated, advanced non-small cell lung cancer (NSCLC) Journal of Clinical Oncology, 2012, 30, 7543-7543.	1.6	28
49	Prognostic Significance of the Number of Metastatic pN2 Lymph Nodes in Stage IIIA-N2 Non–Small-Cell Lung Cancer After Curative Resection. Clinical Lung Cancer, 2015, 16, e203-e212.	2.6	27
50	The HSP90 inhibitor, NVP-AUY922, attenuates intrinsic PI3K inhibitor resistance in KRAS-mutant non-small cell lung cancer. Cancer Letters, 2017, 406, 47-53.	7.2	27
51	Brigatinib (BRG) in patients (pts) with crizotinib (CRZ)-refractory ALK+ non-small cell lung cancer (NSCLC): First report of efficacy and safety from a pivotal randomized phase (ph) 2 trial (ALTA) Journal of Clinical Oncology, 2016, 34, 9007-9007.	1.6	25
52	Efficacy and safety of patritumab deruxtecan (HER3-DXd) in EGFR inhibitor-resistant, EGFR-mutated (EGFRm) non-small cell lung cancer (NSCLC) Journal of Clinical Oncology, 2021, 39, 9007-9007.	1.6	24
53	Mobocertinib (TAK-788) in EGFR exon 20 insertion (ex20ins)+ metastatic NSCLC (mNSCLC): Additional results from platinum-pretreated patients (pts) and EXCLAIM cohort of phase 1/2 study Journal of Clinical Oncology, 2021, 39, 9014-9014.	1.6	23
54	Bone metastasis in pulmonary sclerosing hemangioma. Korean Journal of Internal Medicine, 2015, 30, 928-930.	1.7	23

#	Article	IF	CITATIONS
55	Randomized Phase III Trial of Irinotecan Plus Cisplatin versus Etoposide Plus Cisplatin in Chemotherapy-NaÃ <sup>-</sup> ve Korean Patients with Extensive-Disease Small Cell Lung Cancer. Cancer Research and Treatment, 2019, 51, 119-127.	3.0	23
56	Nivolumab in advanced non-small-cell lung cancer patients who failed prior platinum-based chemotherapy. Lung Cancer, 2018, 122, 234-242.	2.0	22
57	Clinical activity of fianlimab (REGN3767), a human anti-LAG-3 monoclonal antibody, combined with cemiplimab (anti-PD-1) in patients (pts) with advanced melanoma Journal of Clinical Oncology, 2021, 39, 9515-9515.	1.6	22
58	First-Line Pemetrexed plus Cisplatin followed by Gefitinib Maintenance Therapy versus Gefitinib Monotherapy in East Asian Never-Smoker Patients with Locally Advanced or Metastatic Nonsquamous Non–Small Cell Lung Cancer: Final Overall Survival Results from a Randomized Phase 3 Study. Journal of Thoracic Oncology, 2016, 11, 370-379.	1.1	21
59	HSP90 inhibitor (NVP-AUY922) enhances the anti-cancer effect of BCL-2 inhibitor (ABT-737) in small cell lung cancer expressing BCL-2. Cancer Letters, 2017, 411, 19-26.	7.2	21
60	Non–Small Cell Lung Cancer with Resistance to EGFR-TKI Therapy: CT Characteristics of T790M Mutation–positive Cancer. Radiology, 2018, 289, 227-237.	7.3	19
61	Impact of pseudoprogression and treatment beyond progression on outcome in patients with non-small cell lung cancer treated with immune checkpoint inhibitors. OncoImmunology, 2020, 9, 1776058.	4.6	19
62	Randomized phase II study of paclitaxel/carboplatin intercalated with gefitinib compared to paclitaxel/carboplatin alone for chemotherapy-naÃ <sup>-</sup> ve non-small cell lung cancer in a clinically selected population excluding patients with non-smoking adenocarcinoma or mutated EGFR. BMC Cancer, 2015, 15, 763.	2.6	18
63	Brigatinib (BRG) in crizotinib (CRZ)-refractory ALK+ non–small cell lung cancer (NSCLC): Efficacy updates and exploratory analysis of CNS ORR and overall ORR by baseline (BL) brain lesion status Journal of Clinical Oncology, 2018, 36, 9061-9061.	1.6	18
64	Predictive factors for a long-term response duration in non-squamous cell lung cancer patients treated with pemetrexed. BMC Cancer, 2016, 16, 417.	2.6	17
65	Phase I study (BLOOM) of AZD3759, a BBB penetrable EGFR inhibitor, in patients with TKI-naÃ <sup>-</sup> ve, EGFRm NSCLC with CNS metastases Journal of Clinical Oncology, 2017, 35, 2006-2006.	1.6	17
66	Genomic Alterations in the RB Pathway Indicate Prognostic Outcomes of Early-Stage Lung Adenocarcinoma. Clinical Cancer Research, 2015, 21, 2613-2623.	7.0	16
67	Mutational Profiling of Malignant Mesothelioma Revealed Potential Therapeutic Targets in EGFR and NRAS. Translational Oncology, 2018, 11, 268-274.	3.7	16
68	Clinical activity of <scp>ASP</scp> 8273 in Asian patients with nonâ€smallâ€cell lung cancer with <scp>EGFR</scp> activating and T790M mutations. Cancer Science, 2018, 109, 2852-2862.	3.9	15
69	MEK114653: A randomized, multicenter, phase II study to assess efficacy and safety of trametinib (T) compared with docetaxel (D) in <i>KRAS</i> -mutant advanced non–small cell lung cancer (NSCLC) Journal of Clinical Oncology, 2013, 31, 8029-8029.	1.6	15
70	Comparison of T790M Acquisition Between Patients Treated with Afatinib and Gefitinib as First-Line Therapy: Retrospective Propensity Score Matching Analysis. Translational Oncology, 2019, 12, 852-858.	3.7	14
71	Clinical outcomes of nivolumab in patients with advanced non-small cell lung cancer in real-world practice, with an emphasis on hyper-progressive disease. Journal of Cancer Research and Clinical Oncology, 2020, 146, 3025-3036.	2.5	14
72	HSP90 inhibitor, AUY922, debilitates intrinsic and acquired lapatinib-resistant HER2-positive gastric cancer cells. BMB Reports, 2018, 51, 660-665.	2.4	14

#	Article	IF	CITATIONS
73	The GTP binding activity of transglutaminase 2 promotes bone metastasis of breast cancer cells by downregulating microRNA-205. American Journal of Cancer Research, 2019, 9, 597-607.	1.4	14
74	Health-Related Quality of Life Outcomes in Patients with Resected Epidermal Growth Factor Receptor–Mutated Non–Small Cell Lung Cancer Who Received Adjuvant Osimertinib in the Phase III ADAURA Trial. Clinical Cancer Research, 2022, 28, 2286-2296.	7.0	14
75	Weekly low dose paclitaxel and cisplatin as first-line chemotherapy for advanced non-small cell lung cancer. Lung Cancer, 2003, 41, 221-226.	2.0	13
76	A multicenter phase II study of sorafenib in combination with erlotinib in patients with advanced non-small cell lung cancer (KCSG-0806). Lung Cancer, 2016, 93, 1-8.	2.0	13
77	Symptom experiences and healthâ€related quality of life among nonâ€small cell lung cancer patients participating in clinical trials. Journal of Clinical Nursing, 2019, 28, 2111-2123.	3.0	13
78	Safety, Pharmacokinetics, and Clinical Activity of Adavosertib in Combination with Chemotherapy in Asian Patients with Advanced Solid Tumors: Phase Ib Study. Targeted Oncology, 2020, 15, 75-84.	3.6	13
79	A randomised phase 2b study comparing the efficacy and safety of belotecan vs. topotecan as monotherapy for sensitive-relapsed small-cell lung cancer. British Journal of Cancer, 2021, 124, 713-720.	6.4	13
80	ASPIRATION: Phase II study of continued erlotinib beyond RECIST progression in Asian patients (pts) with epidermal growth factor receptor ( <i>EGFR</i> ) mutation-positive non-small cell lung cancer (NSCLC) Journal of Clinical Oncology, 2012, 30, TPS7614-TPS7614.	1.6	13
81	Molecular characterization of lung adenocarcinoma from Korean patients using next generation sequencing. PLoS ONE, 2019, 14, e0224379.	2.5	12
82	Afatinib With Pembrolizumab for Treatment of Patients With Locally Advanced/Metastatic Squamous Cell Carcinoma of the Lung: The LUX-Lung IO/KEYNOTE 497 Study Protocol. Clinical Lung Cancer, 2019, 20, e407-e412.	2.6	12
83	Different prognostic implications of hepatic metastasis according to front-line treatment in non-small cell lung cancer: a real-world retrospective study. Translational Lung Cancer Research, 2021, 10, 2551-2561.	2.8	12
84	Phase I study (BLOOM) of AZD3759, a BBB penetrable EGFR inhibitor, in EGFRm NSCLC patients with leptomeningeal metastasis (LM) who progressed after other anti-cancer therapy Journal of Clinical Oncology, 2017, 35, 2069-2069.	1.6	12
85	Phase II Study of Afatinib as Thirdâ€Line Treatment for Patients in Korea With Stage IIIB/IV Nonâ€Small Cell Lung Cancer Harboring Wildâ€Type EGFR. Oncologist, 2014, 19, 702-703.	3.7	11
86	Feasibility, safety, and adequacy of research biopsies for cancer clinical trials at an academic medical center. PLoS ONE, 2019, 14, e0221065.	2.5	11
87	First-line afatinib (A) vs gefitinib (G) for patients (pts) with EGFR mutation positive (EGFRm+) NSCLC (LUX-Lung 7): Patient-reported outcomes (PROs) and impact of dose modifications on efficacy and adverse events (AEs) Journal of Clinical Oncology, 2016, 34, 9046-9046.	1.6	11
88	Does Pemetrexed Work in Targetable, Nonsquamous Non-Small-Cell Lung Cancer? A Narrative Review. Cancers, 2020, 12, 2658.	3.7	10
89	Realâ€world utility of nextâ€generation sequencing for targeted gene analysis and its application to treatment in lung adenocarcinoma. Cancer Medicine, 2021, 10, 3197-3204.	2.8	10
90	ASTRIS: A real world treatment study of osimertinib in patients (pts) with EGFR T790M positive non-small cell lung cancer (NSCLC) Journal of Clinical Oncology, 2017, 35, 9036-9036.	1.6	10

#	Article	IF	CITATIONS
91	Preliminary Phase II results of a multicenter, open-label study of nazartinib (EGF816) in adult patients with treatment-naÃ <sup>-</sup> ve <i>EGFR</i> -mutant non-small cell lung cancer (NSCLC) Journal of Clinical Oncology, 2018, 36, 9094-9094.	1.6	10
92	Postoperative radiation therapy following the incomplete resection of a non-small cell lung cancer. Radiation Oncology Journal, 2014, 32, 70.	1.5	10
93	Real World Experience of Nivolumab in Non-Small Cell Lung Cancer in Korea. Cancer Research and Treatment, 2020, 52, 1112-1119.	3.0	10
94	Outcomes and prognostic factors of patients with lung cancer and pneumonia-induced respiratory failure in a medical intensive care unit: A single-center study. Journal of Critical Care, 2014, 29, 414-419.	2.2	9
95	Comprehensive outcomes of on―and offâ€antiviral prophylaxis in hepatitis B patients undergoing cancer chemotherapy: A competing risks analysis. Journal of Medical Virology, 2016, 88, 1576-1586.	5.0	9
96	Real-world use of osimertinib in non–small cell lung cancer: ASTRIS study Korean subgroup analysis. Current Medical Research and Opinion, 2020, 36, 477-482.	1.9	9
97	Outcomes according to initial and subsequent therapies following intracranial progression in patients with EGFR-mutant lung cancer and brain metastasis. PLoS ONE, 2020, 15, e0231546.	2.5	9
98	Exploring the resistance mechanisms of second-line osimertinib and their prognostic implications using next-generation sequencing in patients with non-small-cell lung cancer. European Journal of Cancer, 2021, 148, 202-210.	2.8	9
99	Tivantinib plus erlotinib versus placebo plus erlotinib in Asian patients with previously treated nonsquamous NSCLC with wild-type <i>EGFR:</i> First report of a phase III ATTENTION trial Journal of Clinical Oncology, 2014, 32, 8044-8044.	1.6	9
100	Safety and clinical activity of first-line durvalumab in advanced NSCLC: Updated results from a Phase 1/2 study Journal of Clinical Oncology, 2017, 35, e20504-e20504.	1.6	9
101	A Randomized Double-Blind, Double-Dummy, Multicenter Trial of Azasetron versus Ondansetron to Evaluate Efficacy and Safety in the Prevention of Delayed Nausea and Vomiting Induced by Chemotherapy. Cancer Research and Treatment, 2014, 46, 19-26.	3.0	9
102	Efficacy and safety of patritumab deruxtecan (HER3-DXd) in advanced/metastatic non-small cell lung cancer (NSCLC) without <i>EGFR</i> -activating mutations Journal of Clinical Oncology, 2022, 40, 9017-9017.	1.6	9
103	Impact of clinicopathologic features on leptomeningeal metastasis from lung adenocarcinoma and treatment efficacy with epidermal growth factor receptor tyrosine kinase inhibitor. Thoracic Cancer, 2020, 11, 436-442.	1.9	8
104	CONTACT-01: A phase III, randomized study of atezolizumab plus cabozantinib versus docetaxel in patients with metastatic non-small cell lung cancer (mNSCLC) previously treated with PD-L1/PD-1 inhibitors and platinum-containing chemotherapy Journal of Clinical Oncology, 2021, 39, TPS9134-TPS9134.	1.6	8
105	Abstract CT163: CD73 inhibitor oleclumab plus osimertinib for advanced EGFRm NSCLC: First report of a Phase 1b/2 study. Cancer Research, 2021, 81, CT163-CT163.	0.9	8
106	Clinical significance of NQO1 polymorphism and expression of p53, SOD2, PARP1 in limited-stage small cell lung cancer. International Journal of Clinical and Experimental Pathology, 2014, 7, 6743-51.	0.5	8
107	Afatinib in heavily pretreated advanced NSCLC patients who progressed following prior gefitinib or erlotinib: Compassionate use program in Korea. Lung Cancer, 2018, 119, 36-41.	2.0	7
108	Ramucirumab Safety in East Asian Patients: A Meta-Analysis of Six Global, Randomized, Double-Blind, Placebo-Controlled, Phase III Clinical Trials. Journal of Global Oncology, 2018, 4, 1-12.	0.5	7

#	Article	IF	CITATIONS
109	Updated safety and clinical activity of durvalumab monotherapy in previously treated patients with stage IIIB/IV NSCLC Journal of Clinical Oncology, 2017, 35, 9085-9085.	1.6	7
110	Modulation of Fexofenadine Pharmacokinetics by Osimertinib in Patients With Advanced EGFRâ€Mutated Non–Small Cell Lung Cancer. Journal of Clinical Pharmacology, 2019, 59, 1099-1109.	2.0	6
111	Cardiac Safety Assessment of Lazertinib: Findings From Patients With EGFR Mutation-Positive Advanced NSCLC and Preclinical Studies. JTO Clinical and Research Reports, 2021, 2, 100224.	1.1	6
112	Blockade of CCL2 expression overcomes intrinsic PD-1/PD-L1 inhibitor-resistance in transglutaminase 2-induced PD-L1 positive triple negative breast cancer. American Journal of Cancer Research, 2020, 10, 2878-2894.	1.4	6
113	Safety and clinical activity of MEDI5752, a PD-1/CTLA-4 bispecific checkpoint inhibitor, as monotherapy in patients (pts) with advanced renal cell carcinoma (RCC): Preliminary results from an FTIH trial Journal of Clinical Oncology, 2022, 40, 107-107.	1.6	6
114	Murine Bone Marrow Stromal Cells: Implications for their Use in Gene Modified Cell Therapy. Leukemia and Lymphoma, 2003, 44, 1973-1978.	1.3	5
115	Prognosis of multiâ€level N2â€positive nonâ€small cell lung cancer according to lymph node staging using endobronchial ultrasoundâ€transbronchial biopsy. Thoracic Cancer, 2018, 9, 684-692.	1.9	5
116	Recurrence-associated gene signature in patients with stage I non-small-cell lung cancer. Scientific Reports, 2021, 11, 19596.	3.3	5
117	A multinational phase III randomized trial with or without consolidation chemotherapy using docetaxel and cisplatin after concurrent chemoradiation in inoperable stage III non-small cell lung cancer (CCheIN) Journal of Clinical Oncology, 2014, 32, 7500-7500.	1.6	5
118	Updated overall survival and safety profile of durvalumab monotherapy in advanced NSCLC Journal of Clinical Oncology, 2018, 36, 169-169.	1.6	5
119	Phase 3 trial of lorlatinib in treatment-naive patients (Pts) with <i>ALK</i> -positive advanced non–small cell lung cancer (NSCLC): Comprehensive plasma and tumor genomic analyses Journal of Clinical Oncology, 2022, 40, 9070-9070.	1.6	5
120	Optimizing palliative chemotherapy for advanced invasive mucinous adenocarcinoma of the lung. BMC Cancer, 2021, 21, 731.	2.6	4
121	Nazartinib (EGF816) in patients with treatment-naÃ <sup>-</sup> ve <i>EGFR</i> -mutant non-small cell lung cancer (NSCLC): Updated phase II results Journal of Clinical Oncology, 2020, 38, 9574-9574.	1.6	4
122	Osimertinib in Patients with T790M-Positive Advanced Non-small Cell Lung Cancer: Korean Subgroup Analysis from Phase II Studies. Cancer Research and Treatment, 2020, 52, 284-291.	3.0	4
123	Real-world outcomes of anti-PD1 antibodies in platinum-refractory, PD-L1-positive recurrent and/or metastatic non-small cell lung cancer, and its potential practical predictors: first report from Korean Cancer Study Group LU19-05. Journal of Cancer Research and Clinical Oncology, 2021, 147, 2459-2469.	2.5	3
124	Brigatinib (BRG) in ALK+ crizotinib (CRZ)-refractory non-small cell lung cancer (NSCLC): Final results of the phase 1/2 and phase 2 (ALTA) trials Journal of Clinical Oncology, 2021, 39, 9071-9071.	1.6	3
125	Phase 1b Open-Label Trial of Afatinib Plus Xentuzumab (BI 836845) in Patients With EGFR Mutation-Positive NSCLC After Progression on EGFR Tyrosine Kinase Inhibitors. JTO Clinical and Research Reports, 2021, 2, 100206.	1.1	3
126	Brigatinib (BRG) in patients (pts) with crizotinib (CRZ)-refractory ALK+ non-small cell lung cancer (NSCLC) and brain metastases in the pivotal randomized phase 2 ALTA trial Journal of Clinical Oncology, 2017, 35, e20502-e20502.	1.6	3

#	Article	IF	CITATIONS
127	Brigatinib (BRG) in crizotinib (CRZ)-refractory ALK+ non-small cell lung cancer (NSCLC): Updates from ALTA, a pivotal randomized phase 2 trial Journal of Clinical Oncology, 2017, 35, e20503-e20503.	1.6	3
128	Phase II study of afatinib plus pembrolizumab in patients with squamous cell carcinoma of the lung following progression during or after first-line chemotherapy (LUX-Lung-IO). Lung Cancer, 2022, 166, 107-113.	2.0	3
129	Experience from Asian centers in a named-patient-use program for afatinib in patients with advanced non-small-cell lung cancer who had progressed following prior therapies, including patients with uncommon EGFR mutations. International Journal of Clinical Oncology, 2021, 26, 841-850.	2.2	2
130	The NHance® Mutation-Equipped Anti-MET Antibody ARGX-111 Displays Increased Tissue Penetration and Anti-Tumor Activity in Advanced Cancer Patients. Biomedicines, 2021, 9, 665.	3.2	2
131	Phase III trial of concurrent thoracic radiotherapy (TRT) with either the first cycle or the third cycle of cisplatin and etoposide chemotherapy to determine the optimal timing of TRT for limited-disease small cell lung cancer Journal of Clinical Oncology, 2012, 30, 7004-7004.	1.6	2
132	YH25448, a 3rd generation EGFR-TKI, in patients with EGFR-TKI-resistant NSCLC: Phase I/II study results Journal of Clinical Oncology, 2018, 36, 9033-9033.	1.6	2
133	Renal Parenchymal Malakoplakia Presenting with Abscesses and Hepatic Extension Misdiagnosed as a Malignant Tumor: A Case Report. Korean Journal of Medicine, 2012, 82, 764.	0.3	2
134	Lazertinib, a 3 <sup>rd</sup> generation EGFR-TKI, in patients with EGFR-TKI resistant NSCLC: Updated results of phase I/II Study Journal of Clinical Oncology, 2019, 37, 9037-9037.	1.6	2
135	A phase III, open-label, randomized study of atezolizumab in combination with carboplatin + paclitaxel + bevacizumab compared with pemetrexed + cisplatin or carboplatin with stage IV non-squamous non-small cell lung cancer (NSCLC) with activating EGFR mutation or ALK translocation (ATLAS Trial) lournal of Clinical Oncology, 2020, 38. TPS9636-TPS9636.	1.6	2
136	Transglutaminase 2 induces intrinsic EGFR-TKI resistance in NSCLC harboring EGFR sensitive mutations. American Journal of Cancer Research, 2019, 9, 1708-1721.	1.4	2
137	Efficacy and Safety of Ceritinib 450 mg/day with Food and 750 mg/day in Fasted State in Treatment-NaÃ⁻ve Patients with ALK+ Non–Small Cell Lung Cancer: Results from the ASCEND-8 Asian Subgroup Analysis. Cancer Research and Treatment, 2023, 55, 83-93.	3.0	2
138	Abstract CT016: MEDI5752, a novel PD-1/CTLA-4 bispecific checkpoint inhibitor for advanced solid tumors: First-in-human study. Cancer Research, 2022, 82, CT016-CT016.	0.9	2
139	Integrin αvβ3 Induces HSP90 Inhibitor Resistance via FAK Activation in KRAS-Mutant Non-Small Cell Lung Cancer. Cancer Research and Treatment, 2021, , .	3.0	1
140	Phase Ib study of oral PPARÎ <sup>3</sup> agonist efatutazone (CS-7017) in combination with erlotinib in Korean patients with metastatic or unresectable locally advanced non-small cell lung cancer (NSCLC) who progressed after first-line therapy Journal of Clinical Oncology, 2012, 30, e18052-e18052.	1.6	1
141	Afatinib in combination with pembrolizumab in patients (pts) with stage IIIB/IV squamous cell carcinoma (SCC) of the lung Journal of Clinical Oncology, 2018, 36, TPS9117-TPS9117.	1.6	1
142	ctDNA resistance landscape of lazertinib, a third-generation EGFR tyrosine kinase inhibitor (TKI) Journal of Clinical Oncology, 2020, 38, 9601-9601.	1.6	1
143	PROGNOSTIC SIGNIFICANCE of CD68 EXPRESSION for Korean PATIENTS with HODGKIN'S LYMPHOMA. Blood, 2010, 116, 3888-3888.	1.4	1
144	Gefitinib with pemetrexed as first-line therapy in patients with advanced nonsquamous non-small cell lung cancer with activating epidermal growth factor receptor mutations. Annals of Translational Medicine, 2017, 5, 11-11.	1.7	1

#	Article	IF	CITATIONS
145	A plain language summary of results from the ADAURA study: osimertinib after surgery for patients who have early-stage EGFR-mutated non-small cell lung cancer. Future Oncology, 2021, 17, 4827-4835.	2.4	1
146	The impact of systematic assessment for adverse events on unscheduled hospital utilization in patients receiving neoadjuvant or adjuvant chemotherapy: A retrospective multicenter study. Cancer Medicine, 2022, 11, 705-714.	2.8	1
147	A study of senaparib in combination with temozolomide for the treatment of patients with advanced solid tumors and extensive-stage small cell lung cancer Journal of Clinical Oncology, 2022, 40, 3102-3102.	1.6	1
148	Massive Hemoptysis Cases Intubated with the Univent® Bronchial Blocker for Lung Protection. Tuberculosis and Respiratory Diseases, 2012, 72, 212.	1.8	0
149	Comments on the trial of cisplatin and etoposide plus thoracic radiotherapy followed by nivolumab or placebo for locally advanced non-small cell lung cancer (RTOG 3505). Journal of Thoracic Disease, 2017, 9, 3525-3528.	1.4	0
150	Abstract 663: Longitudinal AXL assessment of circulating tumor cells (CTCs) and its clinical implication in the patients with EGFR mutated non-small cell lung cancer (NSCLC). , 2021, , .		0
151	Initial Cytoreductive Treatment with Thalidomide Plus Bolus Vincristine/Doxorubicin and Reduced Dexamethasone (T-bVAd) Followed by Autologous Stem Cell Transplantation (ASCT) for Multiple Myeloma Blood, 2007, 110, 5094-5094.	1.4	0
152	Short Course Adjuvant Chemotherapy for Localized Diffuse Large B-Cell Lymphoma After Complete Resection Blood, 2010, 116, 2824-2824.	1.4	0
153	Paclitaxel/carboplatin (PC) intercalated with gefitinib or paclitaxel/carboplatin (PC) for advanced non-small cell lung cancer (NSCLC) in selected population who were smoker or wild-type EGFR: Randomized phase II study Journal of Clinical Oncology, 2013, 31, e19079-e19079.	1.6	0
154	Phase II study of nivolumab in patients with advanced non-small cell lung cancer (NSCLC) in Korea Journal of Clinical Oncology, 2017, 35, 92-92.	1.6	0
155	The role of FDC-PET during osimertinib treatment to predict the responsiveness of tumor early in patients with stage IV non-small cell lung cancer: A pilot study Journal of Clinical Oncology, 2018, 36, e21150-e21150.	1.6	0
156	Durvalumab (D) +/- tremelimumab (T) + chemotherapy (CT) in first-line (1L) metastatic (m) NSCLC: AE management in POSEIDON Journal of Clinical Oncology, 2022, 40, 9035-9035.	1.6	0
157	Abstract CT558: Capmatinib vs docetaxel as second- or third-line therapy in patients with locally advanced or metastatic <i>MET</i> ex14-mutated NSCLC: The GeoMETry-III trial. Cancer Research, 2022, 82, CT558-CT558.	0.9	0