
List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/769574/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Polygenic risk scores for prediction of breast cancer risk in Asian populations. Genetics in Medicine, 2022, 24, 586-600.	2.4	27
2	Genetic Risk Score for Plasma Uric Acid Levels Is Associated With Early Rapid Kidney Function Decline in Type 2 Diabetes. Journal of Clinical Endocrinology and Metabolism, 2022, 107, e2792-e2800.	3.6	3
3	Interaction between cigarette smoking and genetic polymorphisms on the associations with age of natural menopause and reproductive lifespan: the Singapore Chinese Health Study. Human Reproduction, 2022, 37, 1351-1359.	0.9	3
4	The association of genetic susceptibility to smoking with cardiovascular disease mortality and the benefits of adhering to a DASH diet: The Singapore Chinese Health Study. American Journal of Clinical Nutrition, 2022, 116, 386-393.	4.7	3
5	NAFLD polygenic risk score and risk of hepatocellular carcinoma in an East Asian population. Hepatology Communications, 2022, 6, 2310-2321.	4.3	11
6	Genetic associations with healthy ageing among Chinese adults. , 2022, 8, .		1
7	Shortened Telomere Length in Sputum Cells of Bronchiectasis Patients is Associated with Dysfunctional Inflammatory Pathways. Lung, 2022, 200, 401-407.	3.3	3
8	Differential and shared genetic effects on kidney function between diabetic and non-diabetic individuals. Communications Biology, 2022, 5, .	4.4	17
9	Midlife Leukocyte Telomere Length as an Indicator for Handgrip Strength in Late Life. Journals of Gerontology - Series A Biological Sciences and Medical Sciences, 2021, 76, 172-175.	3.6	4
10	Association of leukocyte telomere length with obesityâ€related traits in Asian children with earlyâ€onset obesity. Pediatric Obesity, 2021, 16, e12771.	2.8	9
11	Association of leukocyte telomere length with chronic kidney disease in East Asians with type 2 diabetes: a Mendelian randomization study. CKJ: Clinical Kidney Journal, 2021, 14, 2371-2376.	2.9	12
12	DNA methylation and breast cancer-associated variants. Breast Cancer Research and Treatment, 2021, 188, 713-727.	2.5	7
13	Association of Genetic Variants for Plasma LRG1 With Rapid Decline in Kidney Function in Patients With Type 2 Diabetes. Journal of Clinical Endocrinology and Metabolism, 2021, 106, 2384-2394.	3.6	9
14	Low frequency variants associated with leukocyte telomere length in the Singapore Chinese population. Communications Biology, 2021, 4, 519.	4.4	15
15	The association of genetically determined serum glycine with cardiovascular risk in East Asians. Nutrition, Metabolism and Cardiovascular Diseases, 2021, 31, 1840-1844.	2.6	4
16	Discovery of Novel Genetic Risk Loci for Acute Central Serous Chorioretinopathy and Genetic Pleiotropic Effect With Age-Related Macular Degeneration. Frontiers in Cell and Developmental Biology, 2021, 9, 696885.	3.7	2
17	The power of genetic diversity in genome-wide association studies of lipids. Nature, 2021, 600, 675-679.	27.8	353
18	The Interaction between 30b-5p miRNA and MBNL1 mRNA is Involved in Vascular Smooth Muscle Cell Differentiation in Patients with Coronary Atherosclerosis. International Journal of Molecular Sciences, 2020, 21, 11.	4.1	31

#	Article	IF	CITATIONS
19	Correlation of Telomere Length in Adipose Tissue and Leukocytes and its Association with Postsurgical Weight Loss. Obesity, 2020, 28, 2424-2430.	3.0	4
20	Interaction between a haptoglobin genetic variant and coronary artery disease (CAD) risk factors on CAD severity in Singaporean Chinese population. Molecular Genetics & Genomic Medicine, 2020, 8, e1450.	1.2	3
21	Effect of plasma polyunsaturated fatty acid levels on leukocyte telomere lengths in the Singaporean Chinese population. Nutrition Journal, 2020, 19, 119.	3.4	16
22	Gene-educational attainment interactions in a multi-ancestry genome-wide meta-analysis identify novel blood pressure loci. Molecular Psychiatry, 2020, 26, 2111-2125.	7.9	17
23	Mendelian randomization analysis does not support causal associations of birth weight with hypertension risk and blood pressure in adulthood. European Journal of Epidemiology, 2020, 35, 685-697.	5.7	9
24	Association of <i>G6PD</i> variants with hemoglobin A1c and impact on diabetes diagnosis in East Asian individuals. BMJ Open Diabetes Research and Care, 2020, 8, e001091.	2.8	12
25	Elevated β-cell stress levels promote severe diabetes development in mice with MODY4. Journal of Endocrinology, 2020, 244, 323-337.	2.6	4
26	Large-Scale Whole-Genome Sequencing of Three Diverse Asian Populations in Singapore. Cell, 2019, 179, 736-749.e15.	28.9	126
27	Associations of autozygosity with a broad range of human phenotypes. Nature Communications, 2019, 10, 4957.	12.8	84
28	Association of Birth Weight With Type 2 Diabetes and Glycemic Traits. JAMA Network Open, 2019, 2, e1910915.	5.9	41
29	Multiancestry Genome-Wide Association Study of Lipid Levels Incorporating Gene-Alcohol Interactions. American Journal of Epidemiology, 2019, 188, 1033-1054.	3.4	85
30	Multi-ancestry study of blood lipid levels identifies four loci interacting with physical activity. Nature Communications, 2019, 10, 376.	12.8	64
31	Loci for human leukocyte telomere length in the Singaporean Chinese population and trans-ethnic genetic studies. Nature Communications, 2019, 10, 2491.	12.8	64
32	A catalog of genetic loci associated with kidney function from analyses of a million individuals. Nature Genetics, 2019, 51, 957-972.	21.4	549
33	A multi-ancestry genome-wide study incorporating gene–smoking interactions identifies multiple new loci for pulse pressure and mean arterial pressure. Human Molecular Genetics, 2019, 28, 2615-2633.	2.9	31
34	Multi-ancestry genome-wide gene–smoking interaction study of 387,272 individuals identifies new loci associated with serum lipids. Nature Genetics, 2019, 51, 636-648.	21.4	112
35	PAX4 R192H is associated with younger onset of Type 2 diabetes in East Asians in Singapore. Journal of Diabetes and Its Complications, 2019, 33, 53-58.	2.3	7
36	A Large-Scale Multi-ancestry Genome-wide Study Accounting for Smoking Behavior Identifies Multiple Significant Loci for Blood Pressure. American Journal of Human Genetics, 2018, 102, 375-400.	6.2	123

#	Article	IF	CITATIONS
37	A Genome-Wide Association Study of Diabetic Kidney Disease in Subjects With Type 2 Diabetes. Diabetes, 2018, 67, 1414-1427.	0.6	136
38	Genetic markers for urine haptoglobin is associated with decline in renal function in type 2 diabetes in East Asians. Scientific Reports, 2018, 8, 5109.	3.3	15
39	Interethnic analyses of blood pressure loci in populations of East Asian and European descent. Nature Communications, 2018, 9, 5052.	12.8	75
40	Gene-diet interaction effects on BMI levels in the Singapore Chinese population. Nutrition Journal, 2018, 17, 31.	3.4	11
41	A study of Kibbutzim in Israel reveals risk factors for cardiometabolic traits and subtle population structure. European Journal of Human Genetics, 2018, 26, 1848-1858.	2.8	12
42	Novel genetic associations for blood pressure identified via gene-alcohol interaction in up to 570K individuals across multiple ancestries. PLoS ONE, 2018, 13, e0198166.	2.5	94
43	Genetic variants in the receptor for advanced glycation end products (<i>RAGE</i>) gene were associated with circulating soluble RAGE level but not with renal function among Asians with type 2 diabetes: a genome-wide association study. Nephrology Dialysis Transplantation, 2017, 32, gfw263.	0.7	18
44	Association analyses of East Asian individuals and trans-ancestry analyses with European individuals reveal new loci associated with cholesterol and triglyceride levels. Human Molecular Genetics, 2017, 26, 1770-1784.	2.9	135
45	Genome-Wide Association Study Meta-Analysis of Long-Term Average Blood Pressure in East Asians. Circulation: Cardiovascular Genetics, 2017, 10, e001527.	5.1	26
46	Exome chip meta-analysis identifies novel loci and East Asian–specific coding variants that contribute to lipid levels and coronary artery disease. Nature Genetics, 2017, 49, 1722-1730.	21.4	129
47	Single-cell transcriptomics of East-Asian pancreatic islets cells. Scientific Reports, 2017, 7, 5024.	3.3	16
48	The genetic variation rs6903956 in the novel androgen-dependent tissue factor pathway inhibitor regulating protein (ADTRP) gene is not associated with levels of plasma coagulation factors in the Singaporean Chinese. Thrombosis Journal, 2017, 15, 1.	2.1	4
49	Utility of genetic and non-genetic risk factors in predicting coronary heart disease in Singaporean Chinese. European Journal of Preventive Cardiology, 2017, 24, 153-160.	1.8	11
50	HDL-cholesterol levels and risk of age-related macular degeneration: a multiethnic genetic study using Mendelian randomization. International Journal of Epidemiology, 2017, 46, 1891-1902.	1.9	73
51	Genome-wide association study identifies a missense variant at APOA5 for coronary artery disease in Multi-Ethnic Cohorts from Southeast Asia. Scientific Reports, 2017, 7, 17921.	3.3	28
52	Interaction Between Peroxisome Proliferator Activated Receptor δ and Epithelial Membrane Protein 2 Polymorphisms Influences HDLâ€C Levels in the Chinese Population. Annals of Human Genetics, 2016, 80, 282-293.	0.8	1
53	Genome-wide association studies in East Asians identify new loci for waist-hip ratio and waist circumference. Scientific Reports, 2016, 6, 17958.	3.3	58
54	A genome-wide association study of n-3 and n-6 plasma fatty acids in a Singaporean Chinese population. Genes and Nutrition, 2015, 10, 53.	2.5	53

#	Article	IF	CITATIONS
55	Genetics of Type 2 Diabetes and Clinical Utility. Genes, 2015, 6, 372-384.	2.4	34
56	Genetic studies of body mass index yield new insights for obesity biology. Nature, 2015, 518, 197-206.	27.8	3,823
57	New loci and coding variants confer risk for age-related macular degeneration in East Asians. Nature Communications, 2015, 6, 6063.	12.8	147
58	Interaction effects between Paraoxonase 1 variants and cigarette smoking on risk of coronary heart disease in a Singaporean Chinese population. Atherosclerosis, 2015, 240, 40-45.	0.8	17
59	Trans-ancestry genome-wide association study identifies 12 genetic loci influencing blood pressure and implicates a role for DNA methylation. Nature Genetics, 2015, 47, 1282-1293.	21.4	294
60	Meta-analysis of genome-wide association studies in East Asian-ancestry populations identifies four new loci for body mass index. Human Molecular Genetics, 2014, 23, 5492-5504.	2.9	192
61	Low copy number of the salivary amylase gene predisposes to obesity. Nature Genetics, 2014, 46, 492-497.	21.4	214
62	Are C-Reactive Protein Associated Genetic Variants Associated with Serum Levels and Retinal Markers of Microvascular Pathology in Asian Populations from Singapore?. PLoS ONE, 2013, 8, e67650.	2.5	23
63	Common variants at CDKAL1 and KLF9 are associated with body mass index in east Asian populations. Nature Genetics, 2012, 44, 302-306.	21.4	240
64	Meta-analysis identifies common variants associated with body mass index in east Asians. Nature Genetics, 2012, 44, 307-311.	21.4	372
65	<i>FTO</i> Variants Are Associated With Obesity in the Chinese and Malay Populations in Singapore. Diabetes, 2008, 57, 2851-2857.	0.6	152