

Hyemyung Seo

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7712218/publications.pdf>

Version: 2024-02-01

47
papers

2,775
citations

218381

26
h-index

214527

47
g-index

48
all docs

48
docs citations

48
times ranked

4357
citing authors

#	ARTICLE	IF	CITATIONS
1	Spotting-based differentiation of functional dopaminergic progenitors from human pluripotent stem cells. <i>Nature Protocols</i> , 2022, , .	5.5	6
2	Analysis of multi-omics data on the relationship between epigenetic changes and nervous system disorders caused by exposure to environmentally harmful substances. <i>Environmental Toxicology</i> , 2022, 37, 802-813.	2.1	5
3	Brain cells derived from Alzheimer's disease patients have multiple specific innate abnormalities in energy metabolism. <i>Molecular Psychiatry</i> , 2021, 26, 5702-5714.	4.1	54
4	Dysfunction of X-linked inhibitor of apoptosis protein (XIAP) triggers neuropathological processes via altered p53 activity in Huntington's disease. <i>Progress in Neurobiology</i> , 2021, 204, 102110.	2.8	8
5	Matrix Metalloproteinase-8 Inhibitor Ameliorates Inflammatory Responses and Behavioral Deficits in LRRK2 G2019S Parkinson's Disease Model Mice. <i>Biomolecules and Therapeutics</i> , 2021, 29, 483-491.	1.1	4
6	Modulation of SETDB1 activity by APQ ameliorates heterochromatin condensation, motor function, and neuropathology in a Huntington's disease mouse model. <i>Journal of Enzyme Inhibition and Medicinal Chemistry</i> , 2021, 36, 856-868.	2.5	7
7	Iroquois Homeobox Protein 2 Identified as a Potential Biomarker for Parkinson's Disease. <i>International Journal of Molecular Sciences</i> , 2020, 21, 3455.	1.8	7
8	Personalized iPSC-Derived Dopamine Progenitor Cells for Parkinson's Disease. <i>New England Journal of Medicine</i> , 2020, 382, 1926-1932.	13.9	298
9	Human autologous iPSC-derived dopaminergic progenitors restore motor function in Parkinson's disease models. <i>Journal of Clinical Investigation</i> , 2020, 130, 904-920.	3.9	102
10	HDAC Inhibition by Valproic Acid Induces Neuroprotection and Improvement of PD-like Behaviors in LRRK2 R1441G Transgenic Mice. <i>Experimental Neurobiology</i> , 2019, 28, 504-515.	0.7	31
11	Alpha-Synuclein Suppresses Retinoic Acid-Induced Neuronal Differentiation by Targeting the Glycogen Synthase Kinase-3 β / β -Catenin Signaling Pathway. <i>Molecular Neurobiology</i> , 2018, 55, 1607-1619.	1.9	14
12	Increase in anti-apoptotic molecules, nucleolin, and heat shock protein 70, against upregulated LRRK2 kinase activity. <i>Animal Cells and Systems</i> , 2018, 22, 273-280.	0.8	9
13	Cell-Penetrating Peptide-Patchy Deformable Polymeric Nanovehicles with Enhanced Cellular Uptake and Transdermal Delivery. <i>Biomacromolecules</i> , 2018, 19, 2682-2690.	2.6	39
14	Oxidized DJ-1 Levels in Urine Samples as a Putative Biomarker for Parkinson's Disease. <i>Parkinson's Disease</i> , 2018, 2018, 1-9.	0.6	13
15	Age-associated bimodal transcriptional drift reduces intergenic disparities in transcription. <i>Aging</i> , 2018, 10, 789-807.	1.4	15
16	Suppression of neuroinflammation by matrix metalloproteinase-8 inhibitor in aged normal and LRRK2 G2019S Parkinson's disease model mice challenged with lipopolysaccharide. <i>Biochemical and Biophysical Research Communications</i> , 2017, 493, 879-886.	1.0	18
17	Age-associated chromatin relaxation is enhanced in Huntington's disease mice. <i>Aging</i> , 2017, 9, 803-822.	1.4	24
18	MiR-126 Regulates Growth Factor Activities and Vulnerability to Toxic Insult in Neurons. <i>Molecular Neurobiology</i> , 2016, 53, 95-108.	1.9	48

#	ARTICLE	IF	CITATIONS
19	G2385R and I2020T Mutations Increase LRRK2 GTPase Activity. <i>BioMed Research International</i> , 2016, 2016, 1-8.	0.9	28
20	Gene therapy by proteasome activator, PA28 β , improves motor coordination and proteasome function in Huntington α ™s disease YAC128 mice. <i>Neuroscience</i> , 2016, 324, 20-28.	1.1	36
21	Leucine-rich repeat kinase 2 exacerbates neuronal cytotoxicity through phosphorylation of histone deacetylase 3 and histone deacetylation. <i>Human Molecular Genetics</i> , 2016, 26, ddw363.	1.4	17
22	β -Lapachone increases phase II antioxidant enzyme expression via NQO1-AMPK/PI3K-Nrf2/ARE signaling in rat primary astrocytes. <i>Free Radical Biology and Medicine</i> , 2016, 97, 168-178.	1.3	44
23	Neuroanatomical Visualization of the Impaired Striatal Connectivity in Huntington α ™s Disease Mouse Model. <i>Molecular Neurobiology</i> , 2016, 53, 2276-2286.	1.9	8
24	Leucine-Rich Repeat Kinase 2 (LRRK2) phosphorylates p53 and induces p21WAF1/CIP1 expression. <i>Molecular Brain</i> , 2015, 8, 54.	1.3	50
25	Enhancement of BACE1 Activity by p25/Cdk5-Mediated Phosphorylation in Alzheimer α ™s Disease. <i>PLoS ONE</i> , 2015, 10, e0136950.	1.1	42
26	An early endosome regulator, Rab5b, is an LRRK2 kinase substrate. <i>Journal of Biochemistry</i> , 2015, 157, 485-495.	0.9	70
27	Reduction of Nfia gene expression and subsequent target genes by binge alcohol in the fetal brain. <i>Neuroscience Letters</i> , 2015, 598, 73-78.	1.0	13
28	Increased TRPC5 glutathionylation contributes to striatal neuron loss in Huntington α ™s disease. <i>Brain</i> , 2015, 138, 3030-3047.	3.7	83
29	LRRK2 G2019S mutation attenuates microglial motility by inhibiting focal adhesion kinase. <i>Nature Communications</i> , 2015, 6, 8255.	5.8	79
30	Increased DJ-1 in Urine Exosome of Korean Males with Parkinson α ™s Disease. <i>BioMed Research International</i> , 2014, 2014, 1-8.	0.9	72
31	miR-126 contributes to Parkinson's disease by dysregulating the insulin-like growth factor/phosphoinositide 3-kinase signaling. <i>Neurobiology of Aging</i> , 2014, 35, 1712-1721.	1.5	120
32	JMJD2A attenuation affects cell cycle and tumourigenic inflammatory gene regulation in lipopolysaccharide stimulated neuroectodermal stem cells. <i>Experimental Cell Research</i> , 2014, 328, 361-378.	1.2	11
33	Prediction of miRNA-mRNA associations in Alzheimer α ™s disease mice using network topology. <i>BMC Genomics</i> , 2014, 15, 644.	1.2	25
34	Systemic injection of LPS induces region-specific neuroinflammation and mitochondrial dysfunction in normal mouse brain. <i>Neurochemistry International</i> , 2014, 69, 35-40.	1.9	151
35	Baclofen, a GABAB receptor agonist, enhances ubiquitin-proteasome system functioning and neuronal survival in Huntington α ™s disease model mice. <i>Biochemical and Biophysical Research Communications</i> , 2014, 443, 706-711.	1.0	37
36	Age-dependent effects of valproic acid in Alzheimer α ™s disease (AD) mice are associated with nerve growth factor (NGF) regulation. <i>Neuroscience</i> , 2014, 266, 255-265.	1.1	33

#	ARTICLE	IF	CITATIONS
37	Identification of cancer-specific biomarkers by using microarray gene expression profiling. <i>Biochip Journal</i> , 2013, 7, 57-62.	2.5	9
38	Pharmacological Rescue of Mitochondrial Deficits in iPSC-Derived Neural Cells from Patients with Familial Parkinson's Disease. <i>Science Translational Medicine</i> , 2012, 4, 141ra90.	5.8	444
39	The hAPP Ψ YAC transgenic model has elevated UPS activity in the frontal cortex similar to Alzheimer's disease and Down's syndrome. <i>Journal of Neurochemistry</i> , 2010, 114, 1819-1826.	2.1	13
40	Compensatory changes in the ubiquitin-proteasome system, brain-derived neurotrophic factor and mitochondrial complex II/III in YAC72 and R6/2 transgenic mice partially model Huntington's disease patients. <i>Human Molecular Genetics</i> , 2008, 17, 3144-3153.	1.4	35
41	Proteasome Activator Enhances Survival of Huntington's Disease Neuronal Model Cells. <i>PLoS ONE</i> , 2007, 2, e238.	1.1	110
42	Abnormal APP, cholinergic and cognitive function in Ts65Dn Down's model mice. <i>Experimental Neurology</i> , 2005, 193, 469-480.	2.0	106
43	Generalized brain and skin proteasome inhibition in Huntington's disease. <i>Annals of Neurology</i> , 2004, 56, 319-328.	2.8	164
44	Alzheimer's disease and Down's syndrome: roles of APP, trophic factors and ACh. <i>Trends in Neurosciences</i> , 2002, 25, 79-84.	4.2	181
45	Spatial memory testing decreases hippocampal amyloid precursor protein in young, but not aged, female rats. <i>Neuroscience Letters</i> , 2002, 328, 50-54.	1.0	26
46	A direct role of the homeodomain proteins Phox2a/2b in noradrenaline neurotransmitter identity determination. <i>Journal of Neurochemistry</i> , 2002, 80, 905-916.	2.1	41
47	Cortico-hippocampal APP and NGF levels are dynamically altered by cholinergic muscarinic antagonist or M1 agonist treatment in normal mice. <i>European Journal of Neuroscience</i> , 2002, 15, 498-506.	1.2	25