
Peng Zhang

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/7746742/publications.pdf Version: 2024-02-01



PENC 7HANC

#	Article	IF	CITATIONS
1	ldentification of risk loci with shared effects on five major psychiatric disorders: a genome-wide analysis. Lancet, The, 2013, 381, 1371-1379.	13.7	2,643
2	Genetic relationship between five psychiatric disorders estimated from genome-wide SNPs. Nature Genetics, 2013, 45, 984-994.	21.4	2,067
3	Large-scale genome-wide association analysis of bipolar disorder identifies a new susceptibility locus near ODZ4. Nature Genetics, 2011, 43, 977-983.	21.4	1,283
4	Psychiatric genome-wide association study analyses implicate neuronal, immune and histone pathways. Nature Neuroscience, 2015, 18, 199-209.	14.8	701
5	Polygenic dissection of diagnosis and clinical dimensions of bipolar disorder and schizophrenia. Molecular Psychiatry, 2014, 19, 1017-1024.	7.9	333
6	Joint Analysis of Psychiatric Disorders Increases Accuracy of Risk Prediction for Schizophrenia, Bipolar Disorder, and Major Depressive Disorder. American Journal of Human Genetics, 2015, 96, 283-294.	6.2	225
7	Identification of Pathways for Bipolar Disorder. JAMA Psychiatry, 2014, 71, 657.	11.0	204
8	Enrichment of cis-regulatory gene expression SNPs and methylation quantitative trait loci among bipolar disorder susceptibility variants. Molecular Psychiatry, 2013, 18, 340-346.	7.9	153
9	Interaction of the <i>ADRB2</i> Gene Polymorphism With Childhood Trauma in Predicting Adult Symptoms of Posttraumatic Stress Disorder. JAMA Psychiatry, 2014, 71, 1174.	11.0	80
10	Genome-Wide Association of Bipolar Disorder Suggests an Enrichment of Replicable Associations in Regions near Genes. PLoS Genetics, 2011, 7, e1002134.	3.5	59
11	Genome-wide association of mood-incongruent psychotic bipolar disorder. Translational Psychiatry, 2012, 2, e180-e180.	4.8	58
12	CLINICAL FEATURES OF BIPOLAR DISORDER COMORBID WITH ANXIETY DISORDERS DIFFER BETWEEN MEN AND WOMEN. Depression and Anxiety, 2012, 29, 739-746.	4.1	47
13	A genomeâ€wide association study of bipolar disorder and comorbid migraine. Genes, Brain and Behavior, 2010, 9, 673-680.	2.2	40
14	Genome-wide association study of personality traits in bipolar patients. Psychiatric Genetics, 2011, 21, 190-194.	1.1	37
15	Familyâ€based association analysis to finemap bipolar linkage peak on chromosome 8q24 using 2,500 genotyped SNPs and 15,000 imputed SNPs. Bipolar Disorders, 2010, 12, 786-792.	1.9	34
16	Genotype Imputation Reference Panel Selection Using Maximal Phylogenetic Diversity. Genetics, 2013, 195, 319-330.	2.9	24
17	Genome-Wide Association Study of Irritable vs. Elated Mania Suggests Genetic Differences between Clinical Subtypes of Bipolar Disorder. PLoS ONE, 2013, 8, e53804.	2.5	22
18	Association study between late-onset Alzheimer's disease and the transferrin gene polymorphisms in Chinese. Neuroscience Letters, 2003, 349, 209-211.	2.1	17

Peng Zhang

#	Article	IF	CITATIONS
19	Polygenic risk for anxiety influences anxiety comorbidity and suicidal behavior in bipolar disorder. Translational Psychiatry, 2020, 10, 298.	4.8	16
20	Genome wide association study identifies variants in NBEA associated with migraine in bipolar disorder. Journal of Affective Disorders, 2015, 172, 453-461.	4.1	15
21	Exonic DNA Sequencing of ERBB4 in Bipolar Disorder. PLoS ONE, 2011, 6, e20242.	2.5	13
22	Suggestive linkage at 9p22 in bipolar disorder weighted by alcohol abuse. American Journal of Medical Genetics Part B: Neuropsychiatric Genetics, 2009, 150B, 1133-1138.	1.7	11
23	Evidence for association of bipolar disorder to haplotypes in the 22q12.3 region near the genes stargazin, ift27 and parvalbumin. American Journal of Medical Genetics Part B: Neuropsychiatric Genetics, 2012, 159B, 941-950.	1.7	10
24	Characterization of the Genomic Architecture and Mutational Spectrum of a Small Cell Prostate Carcinoma. Genes, 2014, 5, 366-384.	2.4	9
25	Detecting significant genotype–phenotype association rules in bipolar disorder: market research meets complex genetics. International Journal of Bipolar Disorders, 2018, 6, 24.	2.2	8
26	Efficient region-based test strategy uncovers genetic risk factors for functional outcome in bipolar disorder. European Neuropsychopharmacology, 2019, 29, 156-170.	0.7	7
27	BDNF expression in lymphoblastoid cell lines carrying BDNF SNPs associated with bipolar disorder. Psychiatric Genetics, 2012, 22, 253-255.	1.1	6
28	Choosing Subsamples for Sequencing Studies by Minimizing the Average Distance to the Closest Leaf. Genetics, 2015, 201, 499-511.	2.9	3