

# Kunitoshi Iseki

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/777153/publications.pdf>

Version: 2024-02-01

27  
papers

2,071  
citations

623734

14  
h-index

552781

26  
g-index

27  
all docs

27  
docs citations

27  
times ranked

2313  
citing authors

#	ARTICLE	IF	CITATIONS
1	Blood Pressure, Hypertension, and the Risk of Aortic Dissection Incidence and Mortality: Results From the J-SCH Study, the UK Biobank Study, and a Meta-Analysis of Cohort Studies. <i>Circulation</i> , 2022, 145, 633-644.	1.6	45
2	Nutrition and quality of life in chronic kidney disease patients: a practical approach for salt restriction. <i>Kidney Research and Clinical Practice</i> , 2022, 41, 657-669.	2.2	8
3	Higher cardiovascular mortality in men with persistent dipstick hematuria. <i>Clinical and Experimental Nephrology</i> , 2021, 25, 150-156.	1.6	9
4	Incidence of shunt infection among patients on maintenance dialysis with buttonhole technique: Okinawa hemodialysis survey. <i>Therapeutic Apheresis and Dialysis</i> , 2021, 25, 354-356.	0.9	0
5	The Effect of CKD on Associations between Lifestyle Factors and All-cause, Cancer, and Cardiovascular Mortality: A Population-based Cohort Study. <i>Internal Medicine</i> , 2021, 60, 2189-2200.	0.7	6
6	Impact of Metabolic Syndrome on the Mortality Rate among Participants in a Specific Health Check and Guidance Program in Japan. <i>Internal Medicine</i> , 2020, 59, 2671-2678.	0.7	10
7	Cause-specific mortality in the general population with transient dipstick-proteinuria. <i>PLoS ONE</i> , 2019, 14, e0223005.	2.5	11
8	Glucosuria and all-cause mortality among general screening participants. <i>Clinical and Experimental Nephrology</i> , 2018, 22, 850-859.	1.6	6
9	Association of dipstick hematuria with all-cause mortality in the general population: results from the specific health check and guidance program in Japan. <i>Nephrology Dialysis Transplantation</i> , 2018, 33, 825-832.	0.7	16
10	Dipstick proteinuria and all-cause mortality among the general population. <i>Clinical and Experimental Nephrology</i> , 2018, 22, 1331-1340.	1.6	20
11	Mortality risk among screened subjects of the specific health check and guidance program in Japan 2008-2012. <i>Clinical and Experimental Nephrology</i> , 2017, 21, 978-985.	1.6	33
12	A practical approach of salt and protein restriction for CKD patients in Japan. <i>BMC Nephrology</i> , 2016, 17, 87.	1.8	11
13	Association of the triglycerides to high-density lipoprotein cholesterol ratio with the risk of chronic kidney disease: Analysis in a large Japanese population. <i>Atherosclerosis</i> , 2014, 233, 260-267.	0.8	64
14	Budget impact analysis of chronic kidney disease mass screening test in Japan. <i>Clinical and Experimental Nephrology</i> , 2014, 18, 885-891.	1.6	13
15	Cost-effectiveness of chronic kidney disease mass screening test in Japan. <i>Clinical and Experimental Nephrology</i> , 2012, 16, 279-291.	1.6	58
16	Evidence for Asymptomatic Microhematuria as a Risk Factor for the Development of ESRD. <i>American Journal of Kidney Diseases</i> , 2012, 60, 12-14.	1.9	6
17	The acute and long-term prognosis of cerebral infarction on the basis of the clinical classification of the Oxfordshire community stroke project. <i>Journal of Cerebral Blood Flow and Metabolism</i> , 2005, 25, S151-S151.	4.3	1
18	Body mass index and the risk of development of end-stage renal disease in a screened cohort. <i>Kidney International</i> , 2004, 65, 1870-1876.	5.2	490

#	ARTICLE	IF	CITATIONS
19	Outcome study of renal biopsy patients in Okinawa, Japan. <i>Kidney International</i> , 2004, 66, 914-919.	5.2	36
20	Proteinuria and the risk of developing end-stage renal disease. <i>Kidney International</i> , 2003, 63, 1468-1474.	5.2	470
21	Effect of the duration of dialysis on survival in a cohort of chronic haemodialysis patients. <i>Nephrology Dialysis Transplantation</i> , 2003, 18, 782-787.	0.7	28
22	Haematocrit and the risk of developing end-stage renal disease. <i>Nephrology Dialysis Transplantation</i> , 2003, 18, 899-905.	0.7	66
23	Demographic trends in the Okinawa Dialysis Study (OKIDS) registry (1971-2000). <i>Kidney International</i> , 2002, 61, 668-675.	5.2	56
24	Hypocholesterolemia is a significant predictor of death in a cohort of chronic hemodialysis patients. <i>Kidney International</i> , 2002, 61, 1887-1893.	5.2	308
25	Serum cholesterol and risk of end-stage renal disease in a cohort of mass screening. <i>Clinical and Experimental Nephrology</i> , 1998, 2, 18-24.	1.6	12
26	Risk of developing end-stage renal disease in a cohort of mass screening. <i>Kidney International</i> , 1996, 49, 800-805.	5.2	285
27	A Case of Syndrome of Inappropriate Secretion of Antidiuretic Hormone(SIADH) with Low Plasma Concentrations of Antidiuretic Hormone.. <i>Internal Medicine</i> , 1992, 31, 246-250.	0.7	3