Manami Inoue

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/7791817/publications.pdf

Version: 2024-02-01

510 papers 21,430 citations

74 h-index

9264

20961 115 g-index

520 all docs 520 docs citations

520 times ranked

27730 citing authors

#	Article	IF	CITATIONS
1	Association analysis identifies 65 new breast cancer risk loci. Nature, 2017, 551, 92-94.	27.8	1,099
2	Metagenomic and metabolomic analyses reveal distinct stage-specific phenotypes of the gut microbiota in colorectal cancer. Nature Medicine, 2019, 25, 968-976.	30.7	748
3	Association between Body-Mass Index and Risk of Death in More Than 1 Million Asians. New England Journal of Medicine, 2011, 364, 719-729.	27.0	730
4	What has made the population of Japan healthy?. Lancet, The, 2011, 378, 1094-1105.	13.7	381
5	Genome-wide association study identifies 112 new loci for body mass index in the Japanese population. Nature Genetics, 2017, 49, 1458-1467.	21.4	380
6	Large-scale genome-wide association study in a Japanese population identifies novel susceptibility loci across different diseases. Nature Genetics, 2020, 52, 669-679.	21.4	304
7	Epidemiology of gastric cancer in Japan. Postgraduate Medical Journal, 2005, 81, 419-424.	1.8	290
8	Identification of ten variants associated with risk of estrogen-receptor-negative breast cancer. Nature Genetics, 2017, 49, 1767-1778.	21.4	289
9	Association of Dietary Intake of Soy, Beans, and Isoflavones With Risk of Cerebral and Myocardial Infarctions in Japanese Populations. Circulation, 2007, 116, 2553-2562.	1.6	247
10	Population-specific and trans-ancestry genome-wide analyses identify distinct and shared genetic risk loci for coronary artery disease. Nature Genetics, 2020, 52, 1169-1177.	21.4	206
11	Adult Mortality Attributable to Preventable Risk Factors for Non-Communicable Diseases and Injuries in Japan: A Comparative Risk Assessment. PLoS Medicine, 2012, 9, e1001160.	8.4	196
12	Soy Product and Isoflavone Consumption in Relation to Prostate Cancer in Japanese Men. Cancer Epidemiology Biomarkers and Prevention, 2007, 16, 538-545.	2.5	185
13	Influence of Coffee Drinking on Subsequent Risk of Hepatocellular Carcinoma: A Prospective Study in Japan. Journal of the National Cancer Institute, 2005, 97, 293-300.	6.3	181
14	Gene-environment interaction between an aldehyde dehydrogenase-2 (ALDH2) polymorphism and alcohol consumption for the risk of esophageal cancer. Carcinogenesis, 2001, 22, 913-916.	2.8	176
15	Consumption of n-3 Fatty Acids and Fish Reduces Risk of Hepatocellular Carcinoma. Gastroenterology, 2012, 142, 1468-1475.	1.3	164
16	Fruit and Vegetable Intake and Risk of Breast Cancer by Hormone Receptor Status. Journal of the National Cancer Institute, 2013, 105, 219-236.	6.3	164
17	Esophageal cancer in high-risk areas of China: research progress and challenges. Annals of Epidemiology, 2017, 27, 215-221.	1.9	164
18	Identification of 28 new susceptibility loci for type 2 diabetes in the Japanese population. Nature Genetics, 2019, 51, 379-386.	21.4	164

#	Article	IF	CITATIONS
19	Population health and regional variations of disease burden in Japan, 1990–2015: a systematic subnational analysis for the Global Burden of Disease Study 2015. Lancet, The, 2017, 390, 1521-1538.	13.7	158
20	Plasma Isoflavone Level and Subsequent Risk of Breast Cancer Among Japanese Women: A Nested Case-Control Study From the Japan Public Health Center-Based Prospective Study Group. Journal of Clinical Oncology, 2008, 26, 1677-1683.	1.6	155
21	Attributable causes of cancer in Japan in 2005â€"systematic assessment to estimate current burden of cancer attributable to known preventable risk factors in Japan. Annals of Oncology, 2012, 23, 1362-1369.	1.2	152
22	Insulin resistance and cancer: Epidemiological evidence. Cancer Science, 2010, 101, 1073-1079.	3.9	149
23	Daily Total Physical Activity Level and Premature Death in Men and Women: Results From a Large-Scale Population-Based Cohort Study in Japan (JPHC Study). Annals of Epidemiology, 2008, 18, 522-530.	1.9	147
24	Identification of six new genetic loci associated with atrial fibrillation in the Japanese population. Nature Genetics, 2017, 49, 953-958.	21.4	136
25	Genome-wide association analysis in East Asians identifies breast cancer susceptibility loci at $1q32.1$, $5q14.3$ and $15q26.1$. Nature Genetics, 2014 , 46 , $886-890$.	21.4	135
26	Association between type 2 diabetes and risk of cancer mortality: a pooled analysis of over 771,000 individuals in the Asia Cohort Consortium. Diabetologia, 2017, 60, 1022-1032.	6.3	132
27	Impact of metabolic factors on subsequent cancer risk: results from a large-scale population-based cohort study in Japan. European Journal of Cancer Prevention, 2009, 18, 240-247.	1.3	131
28	Effect of alcohol consumption, cigarette smoking and flushing response on esophageal cancer risk: A population-based cohort study (JPHC study). Cancer Letters, 2009, 275, 240-246.	7.2	128
29	Reproductive factors, hormone use and the risk of lung cancer among middle-aged never-smoking Japanese women: A large-scale population-based cohort study. International Journal of Cancer, 2005, 117, 662-666.	5.1	127
30	Dairy Product, Saturated Fatty Acid, and Calcium Intake and Prostate Cancer in a Prospective Cohort of Japanese Men. Cancer Epidemiology Biomarkers and Prevention, 2008, 17, 930-937.	2.5	125
31	The Impact of Green Tea and Coffee Consumption on the Reduced Risk of Stroke Incidence in Japanese Population. Stroke, 2013, 44, 1369-1374.	2.0	123
32	Characterizing rare and low-frequency height-associated variants in the Japanese population. Nature Communications, 2019, 10, 4393.	12.8	123
33	Association of Animal and Plant Protein Intake With All-Cause and Cause-Specific Mortality in a Japanese Cohort. JAMA Internal Medicine, 2019, 179, 1509.	5.1	120
34	Fine-mapping of 150 breast cancer risk regions identifies 191 likely target genes. Nature Genetics, 2020, 52, 56-73.	21.4	120
35	Coexpression of the c-kit receptor and the stem cell factor in gynecological tumors. Cancer Research, 1994, 54, 3049-53.	0.9	120
36	Interaction between Adiponectin and Leptin Influences the Risk of Colorectal Adenoma. Cancer Research, 2010, 70, 5430-5437.	0.9	115

#	Article	IF	Citations
37	Impact of population aging on trends in diabetes prevalence: A metaâ€regression analysis of 160,000 Japanese adults. Journal of Diabetes Investigation, 2015, 6, 533-542.	2.4	111
38	Genome-wide association study identifies seven novel susceptibility loci for primary open-angle glaucoma. Human Molecular Genetics, 2018, 27, 1486-1496.	2.9	111
39	Meat intake and cause-specific mortality: a pooled analysis of Asian prospective cohort studies. American Journal of Clinical Nutrition, 2013, 98, 1032-1041.	4.7	109
40	Changing epidemiology of Helicobacter pylori in Japan. Gastric Cancer, 2017, 20, 3-7.	5. 3	109
41	Alcohol Drinking and Colorectal Cancer in Japanese: A Pooled Analysis of Results from Five Cohort Studies. American Journal of Epidemiology, 2008, 167, 1397-1406.	3.4	107
42	Dietary Calcium Intake and Risks of Stroke, Its Subtypes, and Coronary Heart Disease in Japanese. Stroke, 2008, 39, 2449-2456.	2.0	103
43	Association of Diabetes With All-Cause and Cause-Specific Mortality in Asia. JAMA Network Open, 2019, 2, e192696.	5.9	103
44	Tobacco Smoking and Mortality in Asia. JAMA Network Open, 2019, 2, e191474.	5.9	102
45	Alcohol consumption and breast cancer risk by estrogen receptor status: in a pooled analysis of 20 studies. International Journal of Epidemiology, 2016, 45, 916-928.	1.9	101
46	Plasma Isoflavones and Subsequent Risk of Prostate Cancer in a Nested Case-Control Study: The Japan Public Health Center. Journal of Clinical Oncology, 2008, 26, 5923-5929.	1.6	100
47	Consumption of sodium and salted foods in relation to cancer and cardiovascular disease: the Japan Public Health Center–based Prospective Study. American Journal of Clinical Nutrition, 2010, 91, 456-464.	4.7	100
48	Body Mass Index and Mortality From All Causes and Major Causes in Japanese: Results of a Pooled Analysis of 7 Large-Scale Cohort Studies. Journal of Epidemiology, 2011, 21, 417-430.	2.4	100
49	Body mass index, physical activity and the risk of pancreatic cancer in relation to smoking status and history of diabetes: a large-scale population-based cohort study in Japan–The JPHC study. Cancer Causes and Control, 2007, 18, 603-612.	1.8	99
50	Psychological Factors, Coffee and Risk of Diabetes Mellitus among Middle-Aged Japanese: a Population-Based Prospective Study in the JPHC Study Cohort. Endocrine Journal, 2009, 56, 459-468.	1.6	99
51	Effect of Coffee and Green Tea Consumption on the Risk of Liver Cancer: Cohort Analysis by Hepatitis Virus Infection Status. Cancer Epidemiology Biomarkers and Prevention, 2009, 18, 1746-1753.	2.5	98
52	Burden of Total and Cause-Specific Mortality Related to Tobacco Smoking among Adults Aged ≥45 Years in Asia: A Pooled Analysis of 21 Cohorts. PLoS Medicine, 2014, 11, e1001631.	8.4	98
53	Synergistic Promoting Effects ofHelicobacter pyloriInfection and High-salt Diet on Gastric Carcinogenesis in Mongolian Gerbils. Japanese Journal of Cancer Research, 2002, 93, 1083-1089.	1.7	96
54	Tobacco Smoking and Gastric Cancer Risk: An Evaluation Based on a Systematic Review of Epidemiologic Evidence among the Japanese Population. Japanese Journal of Clinical Oncology, 2006, 36, 800-807.	1.3	95

#	Article	IF	CITATIONS
55	Tobacco Smoking and Lung Cancer Risk: An Evaluation Based on a Systematic Review of Epidemiological Evidence Among the Japanese Population. Japanese Journal of Clinical Oncology, 2006, 36, 309-324.	1.3	94
56	Plasma C-Reactive Protein and Risk of Colorectal Cancer in a Nested Case-Control Study: Japan Public Health Center–Based Prospective Study. Cancer Epidemiology Biomarkers and Prevention, 2006, 15, 690-695.	2.5	94
57	Diabetes mellitus and cancer risk: Pooled analysis of eight cohort studies in Japan. Cancer Science, 2013, 104, 1499-1507.	3.9	94
58	Mitogen-Activated Protein Kinase Translocates into the Germinal Vesicle and Induces Germinal Vesicle Breakdown in Porcine Oocytes1. Biology of Reproduction, 1998, 58, 130-136.	2.7	93
59	Impact of the revision of a nutrient database on the validity of a self-administered food frequency questionnaire (FFQ). Journal of Epidemiology, 2006, 16, 107-116.	2.4	92
60	Physical activity and risk of colorectal cancer in Japanese men and women: the Japan Public Health Center-based prospective Study. Cancer Causes and Control, 2007, 18, 199-209.	1.8	88
61	Reproducibility and Validity of Dietary Patterns Assessed by a Food Frequency Questionnaire Used in the 5-Year Follow-Up Survey of the Japan Public Health Center-Based Prospective Study. Journal of Epidemiology, 2012, 22, 205-215.	2.4	88
62	Soft drink intake in relation to incident ischemic heart disease, stroke, and stroke subtypes in Japanese men and women: the Japan Public Health Centre–based study cohort I. American Journal of Clinical Nutrition, 2012, 96, 1390-1397.	4.7	88
63	European polygenic risk score for prediction of breast cancer shows similar performance in Asian women. Nature Communications, 2020, 11 , 3833.	12.8	88
64	Tobacco Smoking and Colorectal Cancer Risk: An Evaluation Based on a Systematic Review of Epidemiologic Evidence among the Japanese Population. Japanese Journal of Clinical Oncology, 2006, 36, 25-39.	1.3	83
65	Dietary factors and lung cancer risk in Japanese: with special reference to fish consumption and adenocarcinomas. British Journal of Cancer, 2001, 84, 1199-1206.	6.4	81
66	Passive smoking and lung cancer in Japanese nonâ€smoking women: A prospective study. International Journal of Cancer, 2008, 122, 653-657.	5.1	81
67	Association between body mass index and the colorectal cancer risk in Japan: pooled analysis of population-based cohort studies in Japan. Annals of Oncology, 2012, 23, 479-490.	1.2	79
68	Soy Intake and Breast Cancer Risk: An Evaluation Based on a Systematic Review of Epidemiologic Evidence Among the Japanese Population. Japanese Journal of Clinical Oncology, 2014, 44, 282-295.	1.3	79
69	Prediction of the 10â€year probability of gastric cancer occurrence in the <scp>J</scp> apanese population: the <scp>JPHC</scp> study cohort <scp>II</scp> . International Journal of Cancer, 2016, 138, 320-331.	5.1	78
70	Monoclonal nature of transient abnormal myelopoiesis in Down's syndrome. Blood, 1991, 77, 1161-1163.	1.4	77
71	Subsite (cervix/endometrium)-specific Risk and Protective Factors in Uterus Cancer. Japanese Journal of Cancer Research, 1996, 87, 1001-1009.	1.7	77
72	Isoflavone intake and risk of lung cancer: a prospective cohort study in Japan. American Journal of Clinical Nutrition, 2010, 91, 722-728.	4.7	77

#	Article	IF	Citations
73	Insulin resistance and cancer: epidemiological evidence. Endocrine-Related Cancer, 2012, 19, F1-F8.	3.1	77
74	Green tea consumption and gastric cancer in Japanese: a pooled analysis of six cohort studies. Gut, 2009, 58, 1323-1332.	12.1	76
75	Identification of IGFBP2 and IGFBP3 As Compensatory Biomarkers for CA19-9 in Early-Stage Pancreatic Cancer Using a Combination of Antibody-Based and LC-MS/MS-Based Proteomics. PLoS ONE, 2016, 11, e0161009.	2.5	76
76	Colorectal cancer screening using fecal occult blood test and subsequent risk of colorectal cancer: A prospective cohort study in Japan. Cancer Detection and Prevention, 2007, 31, 3-11.	2.1	75
77	Heterocyclic amines content of meat and fish cooked by Brazilian methods. Journal of Food Composition and Analysis, 2010, 23, 61-69.	3.9	74
78	Body Mass Index, Body Height, and Subsequent Risk of Colorectal Cancer in Middle-Aged and Elderly Japanese Men and Women: Japan Public Health Center-Based Prospective Study. Cancer Causes and Control, 2005, 16, 839-850.	1.8	72
79	Cancer association as a risk factor for anti-HMGCR antibody-positive myopathy. Neurology: Neuroimmunology and NeuroInflammation, 2016, 3, e290.	6.0	71
80	Protective factor against progression from atrophic gastritis to gastric cancerâ€"data from a cohort study in Japan. , 1996, 66, 309-314.		69
81	Genetic polymorphisms of ADH1B, ADH1C and ALDH2, alcohol consumption, and the risk of gastric cancer: the Japan Public Health Center-based prospective study. Carcinogenesis, 2015, 36, 223-231.	2.8	69
82	Changing trends in the prevalence of H. pylori infection in Japan (1908–2003): a systematic review and meta-regression analysis of 170,752 individuals. Scientific Reports, 2017, 7, 15491.	3.3	69
83	Impact of alcohol drinking on total cancer risk: data from a large-scale population-based cohort study in Japan. British Journal of Cancer, 2005, 92, 182-187.	6.4	66
84	Dietary intake of saturated fatty acids and incident stroke and coronary heart disease in Japanese communities: the JPHC Study. European Heart Journal, 2013, 34, 1225-1232.	2.2	66
85	Association of green tea consumption with mortality due to all causes and major causes of death in a Japanese population: the Japan Public Health Center-based Prospective Study (JPHC Study). Annals of Epidemiology, 2015, 25, 512-518.e3.	1.9	66
86	Associations of All-Cause Mortality with Census-Based Neighbourhood Deprivation and Population Density in Japan: A Multilevel Survival Analysis. PLoS ONE, 2014, 9, e97802.	2.5	65
87	Genetic Predisposition to Ischemic Stroke. Stroke, 2017, 48, 253-258.	2.0	64
88	Impact of Body Mass Index on the Risk of Total Cancer Incidence and Mortality Among Middle-Aged Japanese: Data from a Large-Scale Population-Based Cohort Study – The JPHC Study. Cancer Causes and Control, 2004, 15, 671-680.	1.8	63
89	Tobacco Smoking and Breast Cancer Risk: An Evaluation Based on a Systematic Review of Epidemiological Evidence among the Japanese Population. Japanese Journal of Clinical Oncology, 2006, 36, 387-394.	1.3	62
90	Cigarette Smoking and Completed Suicide among Middle-aged Men: A Population-based Cohort Study in Japan. Annals of Epidemiology, 2005, 15, 286-292.	1.9	61

#	Article	IF	CITATIONS
91	Plasma 25-hydroxyvitamin D concentration and subsequent risk of total and site specific cancers in Japanese population: large case-cohort study within Japan Public Health Center-based Prospective Study cohort. BMJ: British Medical Journal, 2018, 360, k671.	2.3	61
92	Meat Consumption and Colorectal Cancer Risk: An Evaluation Based on a Systematic Review of Epidemiologic Evidence Among the Japanese Population. Japanese Journal of Clinical Oncology, 2014, 44, 641-650.	1.3	60
93	Risk factors of biliary tract cancer in a large-scale population-based cohort study in Japan (JPHC) Tj ETQq1 1 0.78 Causes and Control, 2008, 19, 33-41.	4314 rgB 1.8	T /Overlock 1 59
94	Coffee consumption and risk of endometrial cancer: A prospective study in Japan. International Journal of Cancer, 2008, 123, 2406-2410.	5.1	59
95	Dietary Soy and Isoflavone Intake and Risk of Colorectal Cancer in the Japan Public Health Center–Based Prospective Study. Cancer Epidemiology Biomarkers and Prevention, 2008, 17, 2128-2135.	2.5	59
96	Impact of alcohol intake on total mortality and mortality from major causes in Japan: a pooled analysis of six large-scale cohort studies. Journal of Epidemiology and Community Health, 2012, 66, 448-456.	3.7	59
97	Tobacco control challenges in East Asia: proposals for change in the world's largest epidemic region. Tobacco Control, 2014, 23, 359-368.	3.2	59
98	Associations between unprocessed red and processed meat, poultry, seafood and egg intake and the risk of prostate cancer: A pooled analysis of 15 prospective cohort studies. International Journal of Cancer, 2016, 138, 2368-2382.	5.1	59
99	Increased Levels of Branched-Chain Amino Acid Associated With Increased Risk of Pancreatic Cancer in a Prospective Case–Control Study of a Large Cohort. Gastroenterology, 2018, 155, 1474-1482.e1.	1.3	59
100	Patterns of Alcohol Drinking and All-Cause Mortality: Results from a Large-Scale Population-based Cohort Study in Japan. American Journal of Epidemiology, 2007, 165, 1039-1046.	3.4	58
101	High serum total cholesterol levels is a risk factor of ischemic stroke for general Japanese population: The JPHC study. Atherosclerosis, 2012, 221, 565-569.	0.8	58
102	Association of coffee intake with total and cause-specific mortality in a Japanese population: the Japan Public Health Center–based Prospective Study. American Journal of Clinical Nutrition, 2015, 101, 1029-1037.	4.7	58
103	Chemoprevention of colorectal cancer: Past, present, and future. Cancer Science, 2019, 110, 3018-3026.	3.9	58
104	Association of Sleep Duration With All- and Major-Cause Mortality Among Adults in Japan, China, Singapore, and Korea. JAMA Network Open, 2021, 4, e2122837.	5.9	58
105	10-Year risk of colorectal cancer: Development and validation of a prediction model in middle-aged Japanese men. Cancer Epidemiology, 2010, 34, 534-541.	1.9	56
106	Body mass index and breast cancer risk in Japan: a pooled analysis of eight population-based cohort studies. Annals of Oncology, 2014, 25, 519-524.	1.2	55
107	Role and impact of menstrual and reproductive factors on breast cancer risk in Japan. European Journal of Cancer Prevention, 2007, 16 , 116 - 123 .	1.3	54
108	Visceral Fat Volume and the Prevalence of Colorectal Adenoma. American Journal of Epidemiology, 2009, 170, 1502-1511.	3.4	54

#	Article	IF	CITATIONS
109	Plasma cytokine levels and the presence of colorectal cancer. PLoS ONE, 2019, 14, e0213602.	2.5	54
110	Risk factors for breast cancer: epidemiological evidence from Japanese studies. Cancer Science, 2011, 102, 1607-1614.	3.9	53
111	Cigarette Smoking and Esophageal Cancer Risk: An Evaluation Based on a Systematic Review of Epidemiologic Evidence Among the Japanese Population. Japanese Journal of Clinical Oncology, 2012, 42, 63-73.	1.3	53
112	Green tea drinking and subsequent risk of breast cancer in a population to based cohort of Japanese women. Breast Cancer Research, 2010, 12, R88.	5 . 0	52
113	Alcohol and smoking and subsequent risk of prostate cancer in Japanese men: The Japan Public Health Centerâ€based prospective study. International Journal of Cancer, 2014, 134, 971-978.	5.1	52
114	GWAS identifies two novel colorectal cancer loci at 16q24.1 and 20q13.12. Carcinogenesis, 2018, 39, 652-660.	2.8	52
115	Diabetes and cancer risk: A Mendelian randomization study. International Journal of Cancer, 2020, 146, 712-719.	5.1	52
116	Protective Effects of Raw Vegetables and Fruit against Lung Cancer among Smokers and Ex-smokers: A Case-Control Study in the Tokai Area of Japan. Japanese Journal of Cancer Research, 1993, 84, 594-600.	1.7	51
117	Body Size and Risk for Breast Cancer in Relation to Estrogen and Progesterone Receptor Status in Japan. Annals of Epidemiology, 2007, 17, 304-312.	1.9	51
118	Dietary fiber intake and risk of cardiovascular disease in the Japanese population: the Japan Public Health Center-based study cohort. European Journal of Clinical Nutrition, 2011, 65, 1233-1241.	2.9	51
119	Fermented and nonâ€fermented soy food consumption and gastric cancer in Japanese and Korean populations: A metaâ€analysis of observational studies. Cancer Science, 2011, 102, 231-244.	3.9	51
120	Fermented Soy Product Intake Is Inversely Associated with the Development of High Blood Pressure: The Japan Public Health Center-Based Prospective Study. Journal of Nutrition, 2017, 147, 1749-1756.	2.9	51
121	Cigarette smoking and cervical cancer risk: an evaluation based on a systematic review and meta-analysis among Japanese women. Japanese Journal of Clinical Oncology, 2019, 49, 77-86.	1.3	51
122	Public awareness of risk factors for cancer among the Japanese general population: A population-based survey. BMC Public Health, 2006, 6, 2.	2.9	50
123	Association between GWAS-identified lung adenocarcinoma susceptibility loci andEGFRmutations in never-smoking Asian women, and comparison with findings from Western populations. Human Molecular Genetics, 2016, 26, ddw414.	2.9	50
124	Meta-analysis of genome-wide association studies identifies multiple lung cancer susceptibility loci in never-smoking Asian women. Human Molecular Genetics, 2016, 25, 620-629.	2.9	50
125	Background Characteristics of Basic Health Examination Participants: the JPHC Study Baseline Survey. Journal of Epidemiology, 2003, 13, 216-225.	2.4	49
126	12 new susceptibility loci for prostate cancer identified by genome-wide association study in Japanese population. Nature Communications, 2019, 10, 4422.	12.8	49

#	Article	IF	CITATIONS
127	Metabolic factors and subsequent risk of hepatocellular carcinoma by hepatitis virus infection status: a large-scale population-based cohort study of Japanese men and women (JPHC Study Cohort II). Cancer Causes and Control, 2009, 20, 741-750.	1.8	48
128	Fish, <i>n</i> a€‰â^' 3 polyunsaturated fatty acids and <i>n</i> â^' 6 polyunsaturated fatty acids in breast cancer risk: The <scp>J</scp> apan <scp>P</scp> ublic <scp>H</scp> ealth <scp>C</scp> enterâ€based prospective study. International Journal of Cancer, 2015, 137, 2915-2926.	take and 5.1	48
129	Plasma tea polyphenol levels and subsequent risk of breast cancer among Japanese women: a nested case–control study. Breast Cancer Research and Treatment, 2010, 124, 827-834.	2.5	47
130	Association between mortality and incidence rates of coronary heart disease and stroke: The Japan Public Health Center-based prospective (JPHC) study. International Journal of Cardiology, 2016, 222, 281-286.	1.7	47
131	Green tea and cancer and cardiometabolic diseases: a review of the current epidemiological evidence. European Journal of Clinical Nutrition, 2021, 75, 865-876.	2.9	47
132	Cigarette Smoking and Liver Cancer Risk: An Evaluation Based on a Systematic Review of Epidemiologic Evidence among Japanese. Japanese Journal of Clinical Oncology, 2006, 36, 445-456.	1.3	46
133	Cigarette Smoking and Pancreas Cancer Risk: An Evaluation Based on a Systematic Review of Epidemiologic Evidence in the Japanese Population. Japanese Journal of Clinical Oncology, 2011, 41, 1292-1302.	1.3	46
134	Dietary glycemic index, glycemic load and incidence of type 2 diabetes in Japanese men and women: the Japan public health center-based prospective study. Nutrition Journal, 2013, 12, 165.	3.4	46
135	Identification of novel breast cancer susceptibility loci in meta-analyses conducted among Asian and European descendants. Nature Communications, 2020, 11, 1217.	12.8	46
136	Serum triglycerides and colorectal adenoma in a case–control study among cancer screening examinees (Japan). Cancer Causes and Control, 2006, 17, 1245-1252.	1.8	45
137	Isoflavone consumption and subsequent risk of hepatocellular carcinoma in a populationâ€based prospective cohort of Japanese men and women. International Journal of Cancer, 2009, 124, 1644-1649.	5.1	45
138	Green Tea Consumption and Gastric Cancer Risk: An Evaluation Based on a Systematic Review of Epidemiologic Evidence Among the Japanese Population. Japanese Journal of Clinical Oncology, 2012, 42, 335-346.	1.3	45
139	Dietary patterns and suicide in Japanese adults: The Japan Public Health Center-based Prospective Study. British Journal of Psychiatry, 2013, 203, 422-427.	2.8	45
140	Coping strategies and risk of cardiovascular disease incidence and mortality: the Japan Public Health Center-based prospective Study. European Heart Journal, 2016, 37, 890-899.	2.2	45
141	Association of soy and fermented soy product intake with total and cause specific mortality: prospective cohort study. BMJ, The, 2020, 368, m34.	6.0	45
142	Evaluation Based on Systematic Review of Epidemiological Evidence Among Japanese Populations: Tobacco Smoking and Total Cancer Risk. Japanese Journal of Clinical Oncology, 2005, 35, 404-411.	1.3	44
143	Association between serum organochlorines and global methylation level of leukocyte DNA among Japanese women: a cross-sectional study. Science of the Total Environment, 2014, 490, 603-609.	8.0	44
144	Excess mortality due to indirect health effects of the 2011 triple disaster in Fukushima, Japan: a retrospective observational study. Journal of Epidemiology and Community Health, 2017, 71, 974-980.	3.7	44

#	Article	IF	CITATIONS
145	Transethnic Meta-Analysis of Genome-Wide Association Studies Identifies Three New Loci and Characterizes Population-Specific Differences for Coronary Artery Disease. Circulation Genomic and Precision Medicine, 2020, 13, e002670.	3.6	44
146	Effect of Intermittent Administration of Teriparatide (Parathyroid Hormone 1-34) on Bone Morphogenetic Protein-Induced Bone Formation in a Rat Model of Spinal Fusion. Journal of Bone and Joint Surgery - Series A, 2014, 96, e107.	3.0	43
147	The bone morphogenetic protein-2/7 heterodimer is a stronger inducer of bone regeneration than the individual homodimers in a rat spinal fusion model. Spine Journal, 2015, 15, 1379-1390.	1.3	43
148	Association of high-density lipoprotein cholesterol concentration with different types of stroke and coronary heart disease: The Japan Public Health Center-based prospective (JPHC) study. Atherosclerosis, 2017, 265, 147-154.	0.8	43
149	Dietary acrylamide intake and risk of breast cancer: The Japan Public Health Centerâ€based Prospective Study. Cancer Science, 2018, 109, 843-853.	3.9	43
150	Physical inactivity, prolonged sedentary behaviors, and use of visual display terminals as potential risk factors for dry eye disease: JPHC-NEXT study. Ocular Surface, 2020, 18, 56-63.	4.4	42
151	Development of a prediction model for 10-year risk of hepatocellular carcinoma in middle-aged Japanese: The Japan Public Health Center-based Prospective Study Cohort II. Preventive Medicine, 2012, 55, 137-143.	3.4	41
152	Type A behaviour and risk of coronary heart disease: The JPHC Study. International Journal of Epidemiology, 2008, 37, 1395-1405.	1.9	40
153	Vegetable and fruit intake and risk of type 2 diabetes: Japan Public Health Center-based Prospective Study. British Journal of Nutrition, 2013, 109, 709-717.	2.3	40
154	Rice consumption is not associated with risk of cardiovascular disease morbidity or mortality in Japanese men and women: a large population-based, prospective cohort study. American Journal of Clinical Nutrition, 2014, 100, 199-207.	4.7	40
155	Genome-wide association study in East Asians identifies two novel breast cancer susceptibility loci. Human Molecular Genetics, 2016, 25, 3361-3371.	2.9	40
156	The influence of birth cohort and calendar period on global trends in ovarian cancer incidence. International Journal of Cancer, 2020, 146, 749-758.	5.1	40
157	Dietary isoflavone intake and breast cancer risk in case–control studies in Japanese, Japanese Brazilians, and non-Japanese Brazilians. Breast Cancer Research and Treatment, 2009, 116, 401-411.	2.5	39
158	Alcohol drinking and primary liver cancer: A pooled analysis of four Japanese cohort studies. International Journal of Cancer, 2012, 130, 2645-2653.	5.1	39
159	Fish, n–3 PUFA consumption, and pancreatic cancer risk in Japanese: a large, population-based, prospective cohort study. American Journal of Clinical Nutrition, 2015, 102, 1490-1497.	4.7	39
160	High hemoglobin A1c levels within the nonâ€diabetic range are associated with the risk of all cancers. International Journal of Cancer, 2016, 138, 1741-1753.	5.1	39
161	Impact of Alcohol Intake and Drinking Patterns on Mortality From All Causes and Major Causes of Death in a Japanese Population. Journal of Epidemiology, 2018, 28, 140-148.	2.4	39
162	Genomeâ€wide association study identifies gastric cancer susceptibility loci at 12q24.11â€12 and 20q11.21. Cancer Science, 2018, 109, 4015-4024.	3.9	39

#	Article	IF	CITATIONS
163	Seaweed intake and risk of cardiovascular disease: the Japan Public Health Center–based Prospective (JPHC) Study. American Journal of Clinical Nutrition, 2019, 110, 1449-1455.	4.7	39
164	Body mass, tobacco smoking, alcohol drinking and risk of cancer of the small intestine—a pooled analysis of over 500Â000 subjects in the Asia Cohort Consortium. Annals of Oncology, 2012, 23, 1894-1898.	1.2	38
165	Combined impact of five lifestyle factors and subsequent risk of cancer: The Japan Public Health Center Study. Preventive Medicine, 2012, 54, 112-116.	3.4	38
166	Dietary fiber intake and total and cause-specific mortality: the Japan Public Health Center-based prospective study. American Journal of Clinical Nutrition, 2020, 111, 1027-1035.	4.7	38
167	Validity and applicability of a simple questionnaire for the estimation of total and domain-specific physical activity. Diabetology International, 2011, 2, 47-54.	1.4	37
168	Association between adherence to the Japanese diet and all-cause and cause-specific mortality: the Japan Public Health Center-based Prospective Study. European Journal of Nutrition, 2021, 60, 1327-1336.	3.9	37
169	Coffee and tea consumption and mortality from all causes, cardiovascular disease and cancer: a pooled analysis of prospective studies from the Asia Cohort Consortium. International Journal of Epidemiology, 2022, 51, 626-640.	1.9	37
170	Generalizability of Relative Risk Estimates from a Well-defined Population to a General Population. European Journal of Epidemiology, 2006, 21, 253-262.	5.7	36
171	Isoflavone intake and risk of gastric cancer: a population-based prospective cohort study in Japan. American Journal of Clinical Nutrition, 2012, 95, 147-154.	4.7	36
172	Intermittent administration of teriparatide enhances graft bone healing and accelerates spinal fusion in rats with glucocorticoid-induced osteoporosis. Spine Journal, 2015, 15, 298-306.	1.3	36
173	Alcohol Consumption, Social Support, and Risk of Stroke and Coronary Heart Disease Among Japanese Men: The JPHC Study. Alcoholism: Clinical and Experimental Research, 2009, 33, 1025-1032.	2.4	35
174	Fragment c gamma receptor gene polymorphisms and breast cancer risk in case–control studies in Japanese, Japanese Brazilians, and non-Japanese Brazilians. Breast Cancer Research and Treatment, 2011, 126, 497-505.	2.5	35
175	Dietary magnesium intake and risk of incident coronary heart disease in men: A prospective cohort study. Clinical Nutrition, 2018, 37, 1602-1608.	5.0	35
176	Validating the dietary inflammatory index using inflammatory biomarkers in a Japanese population: A cross-sectional study of the JPHC-FFQ validation study. Nutrition, 2020, 69, 110569.	2.4	35
177	Isoflavone, polymorphisms in estrogen receptor genes and breast cancer risk in caseâ€control studies in Japanese, Japanese Brazilians and nonâ€apanese Brazilians. Cancer Science, 2009, 100, 927-933.	3.9	34
178	Clinical Significance of IgG Antibody Titer against <i> Helicobacter pylori</i> . Helicobacter, 2009, 14, 231-236.	3.5	34
179	Dietary pattern and breast cancer risk in Japanese women: the Japan Public Health Center-based Prospective Study (JPHC Study). British Journal of Nutrition, 2016, 115, 1769-1779.	2.3	34
180	Cruciferous Vegetable Intake Is Inversely Associated with Lung Cancer Risk among Current Nonsmoking Men in the Japan Public Health Center (JPHC) Study. Journal of Nutrition, 2017, 147, 841-849.	2.9	34

#	Article	IF	CITATIONS
181	Perceived stress level and risk of cancer incidence in a Japanese population: the Japan Public Health Center (JPHC)-based Prospective Study. Scientific Reports, 2017, 7, 12964.	3.3	34
182	Genome-wide association meta-analysis identifies GP2 gene risk variants for pancreatic cancer. Nature Communications, 2020, 11, 3175.	12.8	34
183	Hepatitis B and C virus infection and risk of lymphoid malignancies: A population-based cohort study (JPHC Study). Cancer Epidemiology, 2015, 39, 562-566.	1.9	33
184	Hepatitis B and C Virus Infection and Risk of Pancreatic Cancer: A Population-Based Cohort Study (JPHC) Tj ETQq	0 <u>0 0</u> rgB1	- /gyerlock 1
185	Association of leisure-time physical activity with total and cause-specific mortality: a pooled analysis of nearly a half million adults in the Asia Cohort Consortium. International Journal of Epidemiology, 2018, 47, 771-779.	1.9	32
186	Plasma testosterone and sex hormoneâ€binding globulin concentrations and the risk of prostate cancer among Japanese men: A nested caseâ€control study. Cancer Science, 2010, 101, 2652-2657.	3.9	31
187	Association between green tea/coffee consumption and biliary tract cancer: A populationâ€based cohort study in Japan. Cancer Science, 2016, 107, 76-83.	3.9	31
188	Cigarette smoking and bladder cancer risk: an evaluation based on a systematic review of epidemiologic evidence in the Japanese population. Japanese Journal of Clinical Oncology, 2016, 46, 273-283.	1.3	31
189	Changes in the Employment Status and Risk of Stroke and Stroke Types. Stroke, 2017, 48, 1176-1182.	2.0	31
190	Green tea consumption and mortality in Japanese men and women: a pooled analysis of eight population-based cohort studies in Japan. European Journal of Epidemiology, 2019, 34, 917-926.	5.7	31
191	Green Tea Consumption and Breast Cancer Risk in Japanese Women: A Case-Control Study. Nutrition and Cancer, 2014, 66, 57-67.	2.0	30
192	Validity of a Self-Administered Food-Frequency Questionnaire for Assessing Amino Acid Intake in Japan: Comparison With Intake From 4-Day Weighed Dietary Records and Plasma Levels. Journal of Epidemiology, 2016, 26, 36-44.	2.4	30
193	Impact of Moderate-Intensity and Vigorous-Intensity Physical Activity on Mortality. Medicine and Science in Sports and Exercise, 2018, 50, 715-721.	0.4	30
194	Body-Mass Index and Pancreatic Cancer Incidence: A Pooled Analysis of Nine Population-Based Cohort Studies With More Than 340,000 Japanese Subjects. Journal of Epidemiology, 2018, 28, 245-252.	2.4	30
195	The Japan Public Health Center-based Prospective Study for the Next Generation (JPHC-NEXT): Study Design and Participants. Journal of Epidemiology, 2020, 30, 46-54.	2.4	30
196	Non-High-Density Lipoprotein Cholesterol and Risk of Stroke Subtypes and Coronary Heart Disease: The Japan Public Health Center-Based Prospective (JPHC) Study. Journal of Atherosclerosis and Thrombosis, 2020, 27, 363-374.	2.0	30
197	Plasma organochlorine levels and subsequent risk of breast cancer among Japanese women: A nested case–control study. Science of the Total Environment, 2008, 402, 176-183.	8.0	29
198	Education in relation to incidence of and mortality from cancer and cardiovascular disease in Japan. European Journal of Public Health, 2008, 18, 466-472.	0.3	29

#	Article	IF	Citations
199	Association of dietary diversity with total mortality and major causes of mortality in the Japanese population: JPHC study. European Journal of Clinical Nutrition, 2020, 74, 54-66.	2.9	29
200	Impact of tobacco smoking on subsequent cancer risk among middle-aged Japanese men and women: data from a large-scale population-based cohort study in Japanâ€"the JPHC study. Preventive Medicine, 2004, 38, 516-522.	3.4	28
201	Secular trends in cancer mortality among Japanese immigrants in the state of São Paulo, Brazil, 1979–2001. European Journal of Cancer Prevention, 2008, 17, 1-8.	1.3	28
202	Lifestyle, weight perception and change in body mass index of Japanese workers: MY Health Up Study. Public Health, 2010, 124, 530-537.	2.9	28
203	Diabetes Mellitus and Liver Cancer Risk: An Evaluation Based on a Systematic Review of Epidemiologic Evidence among the Japanese Population. Japanese Journal of Clinical Oncology, 2014, 44, 986-999.	1.3	28
204	Death by Suicide and Other Externally Caused Injuries After Stroke in Japan (1990–2010). Psychosomatic Medicine, 2014, 76, 452-459.	2.0	28
205	Diagnosed diabetes and premature death among middle-aged Japanese: results from a large-scale population-based cohort study in Japan (JPHC study). BMJ Open, 2015, 5, e007736-e007736.	1.9	28
206	Cigarette smoking and the risk of head and neck cancer in the Japanese population: a systematic review and meta-analysis. Japanese Journal of Clinical Oncology, 2016, 46, 580-595.	1.3	28
207	Dietary consumption of antioxidant vitamins and subsequent lung cancer risk: The ⟨scp⟩J⟨/scp⟩apan ⟨scp⟩P⟨/scp⟩ublic ⟨scp⟩H⟨/scp⟩ealth ⟨scp⟩C⟨/scp⟩enterâ€based prospective study. International Journal of Cancer, 2018, 142, 2441-2460.	5.1	28
208	Dietary Acrylamide Intake and Risk of Esophageal, Gastric, and Colorectal Cancer: The Japan Public Health Center–Based Prospective Study. Cancer Epidemiology Biomarkers and Prevention, 2019, 28, 1461-1468.	2.5	28
209	Low carbohydrate diet and all cause and cause-specific mortality. Clinical Nutrition, 2021, 40, 2016-2024.	5.0	28
210	The Impact of C-Reactive Protein on Risk of Stroke, Stroke Subtypes, and Ischemic Heart Disease in Middle-Aged Japanese: the Japan Public Health Center-Based Study. Journal of Atherosclerosis and Thrombosis, 2012, 19, .	2.0	27
211	Dietary intake of antioxidant vitamins and risk of stroke: the Japan Public Health Center–based Prospective Study. European Journal of Clinical Nutrition, 2017, 71, 1179-1185.	2.9	27
212	Polygenic risk scores for prediction of breast cancer risk in Asian populations. Genetics in Medicine, 2022, 24, 586-600.	2.4	27
213	The Japanese LupusPRO: A cross-cultural validation of an outcome measure for lupus. Lupus, 2017, 26, 849-856.	1.6	26
214	Dietary acrylamide intake and the risk of endometrial or ovarian cancers in Japanese women. Cancer Science, 2018, 109, 3316-3325.	3.9	26
215	Subluxation of the patella. Computed tomography analysis of patellofemoral congruence. Journal of Bone and Joint Surgery - Series A, 1988, 70, 1331-7.	3.0	26
216	Association of Alcohol Intake with the Risk of Malignant Lymphoma and Plasma Cell Myeloma in Japanese: A Population-Based Cohort Study (Japan Public Health Center–based Prospective Study). Cancer Epidemiology Biomarkers and Prevention, 2010, 19, 429-434.	2.5	25

#	Article	IF	CITATIONS
217	Dietary Heterocyclic Amine Intake, <i>NAT2</i> Genetic Polymorphism, and Colorectal Adenoma Risk: The Colorectal Adenoma Study in Tokyo. Cancer Epidemiology Biomarkers and Prevention, 2015, 24, 613-620.	2.5	25
218	Evidence-based cancer prevention recommendations for Japanese. Japanese Journal of Clinical Oncology, 2018, 48, 576-586.	1.3	25
219	Body mass index and colorectal cancer risk: A Mendelian randomization study. Cancer Science, 2021, 112, 1579-1588.	3.9	25
220	Association of body mass index and height with risk of prostate cancer among middle-aged Japanese men. British Journal of Cancer, 2006, 94, 740-742.	6.4	24
221	Fiber intake and risk of subsequent prostate cancer in Japanese men. American Journal of Clinical Nutrition, 2015, 101, 118-125.	4.7	24
222	Circulating sex hormone levels and colorectal cancer risk in Japanese postmenopausal women: The JPHC nested case–control study. International Journal of Cancer, 2019, 145, 1238-1244.	5.1	24
223	Cancer Mortality Among Japanese Immigrants and their Descendants in the State of Sao Paulo, Brazil, 1999-2001. Japanese Journal of Clinical Oncology, 2004, 34, 673-680.	1.3	23
224	Association of body mass index and risk of death from pancreas cancer in Asians. European Journal of Cancer Prevention, 2013, 22, 244-250.	1.3	23
225	Association between Chlamydophila pneumoniae infection and risk of coronary heart disease for Japanese: The JPHC study. Atherosclerosis, 2014, 233, 338-342.	0.8	23
226	Vegetable consumption and colorectal cancer risk: an evaluation based on a systematic review and meta-analysis among the Japanese population. Japanese Journal of Clinical Oncology, 2015, 45, 973-979.	1.3	23
227	Predictors of poor sleep quality in patients with systemic lupus erythematosus. Clinical Rheumatology, 2017, 36, 1053-1062.	2.2	23
228	Coffee drinking and colorectal cancer and its subsites: A pooled analysis of 8 cohort studies in <scp>J</scp> apan. International Journal of Cancer, 2018, 143, 307-316.	5.1	23
229	Dietary patterns and prostate cancer risk in Japanese: the Japan Public Health Center-based Prospective Study (JPHC Study). Cancer Causes and Control, 2018, 29, 589-600.	1.8	23
230	Dietary Inflammatory Index Is Associated With Inflammation in Japanese Men. Frontiers in Nutrition, 2021, 8, 604296.	3.7	23
231	Food/nutrient intake and risk of atrophic gastritis among the Helicobacter pylori-infected population of northeastern Japan. Cancer Science, 2003, 94, 372-377.	3.9	22
232	Green tea and coffee consumption and its association with thyroid cancer risk: a population-based cohort study in Japan. Cancer Causes and Control, 2011, 22, 985-993.	1.8	22
233	Socioeconomic Status Inconsistency and Risk of Stroke Among Japanese Middle-Aged Women. Stroke, 2014, 45, 2592-2598.	2.0	22
234	<scp>C</scp> offee intake and the risk of colorectal adenoma: The colorectal adenoma study in Tokyo. International Journal of Cancer, 2015, 137, 463-470.	5.1	22

#	Article	IF	CITATIONS
235	Coffee drinking and colorectal cancer risk: an evaluation based on a systematic review and meta-analysis among the Japanese population. Japanese Journal of Clinical Oncology, 2016, 46, 781-787.	1.3	22
236	Coffee and green tea consumption in relation to brain tumor risk in a Japanese population. International Journal of Cancer, 2016, 139, 2714-2721.	5.1	22
237	Circulating sclerostin and dickkopf-1 levels in ossification of the posterior longitudinal ligament of the spine. Journal of Bone and Mineral Metabolism, 2016, 34, 315-324.	2.7	22
238	Plasma tea catechins and risk of cardiovascular disease in middle-aged Japanese subjects: The JPHC study. Atherosclerosis, 2018, 277, 90-97.	0.8	22
239	DNA Adductome Analysis Identifies <i>N</i> -Nitrosopiperidine Involved in the Etiology of Esophageal Cancer in Cixian, China. Chemical Research in Toxicology, 2019, 32, 1515-1527.	3.3	22
240	Revisit of an unanswered question by pooled analysis of eight cohort studies in Japan: Does cigarette smoking and alcohol drinking have interaction for the risk of esophageal cancer?. Cancer Medicine, 2019, 8, 6414-6425.	2.8	22
241	Occupational sitting time and risk of all-cause mortality among Japanese workers. Scandinavian Journal of Work, Environment and Health, 2015, 41, 519-528.	3.4	22
242	IGKC and $Fc\hat{l}^3R$ genotypes and humoral immunity to HER2 in breast cancer. Immunobiology, 2014, 219, 113-117.	1.9	21
243	Adjustment of Cell-Type Composition Minimizes Systematic Bias in Blood DNA Methylation Profiles Derived by DNA Collection Protocols. PLoS ONE, 2016, 11, e0147519.	2.5	21
244	Vitamin D Receptor Gene Polymorphism and the Risk of Colorectal Cancer: A Nested Case-Control Study. PLoS ONE, 2016, 11, e0164648.	2.5	21
245	Inclusion of a Genetic Risk Score into a Validated Risk Prediction Model for Colorectal Cancer in Japanese Men Improves Performance. Cancer Prevention Research, 2017, 10, 535-541.	1.5	21
246	The relationship between vegetable/fruit consumption and gallbladder/bile duct cancer: A populationâ€based cohort study in ⟨scp⟩J⟨/scp⟩apan. International Journal of Cancer, 2017, 140, 1009-1019.	5.1	21
247	Helicobacter pylori infection, atrophic gastritis, and risk of pancreatic cancer: A population-based cohort study in a large Japanese population: the JPHC Study. Scientific Reports, 2019, 9, 6099.	3.3	21
248	Smoking and colorectal cancer: A pooled analysis of 10 populationâ€based cohort studies in Japan. International Journal of Cancer, 2021, 148, 654-664.	5.1	21
249	Characterization of mRNA Expression of lκBα and NF-κB Subfamilies in Primary Adult T-cell Leukemia Cells. Japanese Journal of Cancer Research, 1998, 89, 53-59.	1.7	20
250	Alcohol and risk of lung cancer among Japanese men: data from a large-scale population-based cohort study, the JPHC study. Cancer Causes and Control, 2008, 19, 1095-1102.	1.8	20
251	Association of Anthropometric Characteristics with the Risk of Malignant Lymphoma and Plasma Cell Myeloma in a Japanese Population: A Population-Based Cohort Study. Cancer Epidemiology Biomarkers and Prevention, 2010, 19, 1623-1631.	2.5	20
252	Breastfeeding and Breast Cancer Risk: An Evaluation Based on a Systematic Review of Epidemiologic Evidence Among the Japanese Population. Japanese Journal of Clinical Oncology, 2012, 42, 124-130.	1.3	20

#	Article	lF	CITATIONS
253	Coping behaviors and suicide in the middle-aged and older Japanese general population: the Japan Public Health Center-based Prospective Study. Annals of Epidemiology, 2014, 24, 199-205.	1.9	20
254	<i>CYP1A1</i> , <i>GSTM1</i> and <i>GSTT1</i> genetic polymorphisms and gastric cancer risk among Japanese: A nested caseâ€"control study within a largeâ€scale populationâ€based prospective study. International Journal of Cancer, 2016, 139, 759-768.	5.1	20
255	Marital Transition and Risk of Stroke. Stroke, 2016, 47, 991-998.	2.0	20
256	Smoking cessation and subsequent risk of cancer: A pooled analysis of eight population-based cohort studies in Japan. Cancer Epidemiology, 2017, 51, 98-108.	1.9	20
257	Association between NAT2, CYP1A1, and CYP1A2 genotypes, heterocyclic aromatic amines, and prostate cancer risk: a case control study in Japan. Environmental Health and Preventive Medicine, 2017, 22, 72.	3.4	20
258	Dietary patterns and colorectal cancer risk in middle-aged adults: AÂlarge population-based prospective cohort study. Clinical Nutrition, 2018, 37, 1019-1026.	5.0	20
259	Validity of a Self-administered Food Frequency Questionnaire for the Estimation of Acrylamide Intake in the Japanese Population: The JPHC FFQ Validation Study. Journal of Epidemiology, 2018, 28, 482-487.	2.4	20
260	Reproductive history and risk of cognitive impairment in Japanese women. Maturitas, 2019, 128, 22-28.	2.4	20
261	Isolation and characterization of a human cDNA clone (GCN5L1) homologous to GCN5, a yeast transcription activator. Cytogenetic and Genome Research, 1996, 73, 134-136.	1.1	19
262	Dietary Isoflavone Intake, Polymorphisms in the CYP17, CYP19, 17β-HSD1, and SHBG Genes, and Risk of Breast Cancer in Case-Control Studies in Japanese, Japanese Brazilians, and Non-Japanese Brazilians. Nutrition and Cancer, 2010, 62, 466-475.	2.0	19
263	Differences in suicide risk according to living arrangements in Japanese men and women – The Japan Public Health Center-based (JPHC) prospective study. Journal of Affective Disorders, 2011, 131, 113-119.	4.1	19
264	Neighborhood Deprivation and Risk of Cancer Incidence, Mortality and Survival: Results from a Population-Based Cohort Study in Japan. PLoS ONE, 2014, 9, e106729.	2.5	19
265	The association of active and secondhand smoking with oral health in adults: Japan public health center-based study. Tobacco Induced Diseases, 2015, 13, 19.	0.6	19
266	High serum total cholesterol is associated with suicide mortality in Japanese women. Acta Psychiatrica Scandinavica, 2017, 136, 259-268.	4.5	19
267	Smoking and Pancreatic Cancer Incidence: A Pooled Analysis of 10 Population-Based Cohort Studies in Japan. Cancer Epidemiology Biomarkers and Prevention, 2019, 28, 1370-1378.	2.5	19
268	Prediagnostic circulating inflammation biomarkers and esophageal squamous cell carcinoma: A case–cohort study in Japan. International Journal of Cancer, 2020, 147, 686-691.	5.1	19
269	High-Negative Anti– <i>Helicobacter pylori</i> IgG Antibody Titers and Long-Term Risk of Gastric Cancer: Results from a Large-Scale Population-Based Cohort Study in Japan. Cancer Epidemiology Biomarkers and Prevention, 2020, 29, 420-426.	2.5	19
270	Fermented soy products intake and risk of cardiovascular disease and total cancer incidence: The Japan Public Health Center-based Prospective study. European Journal of Clinical Nutrition, 2021, 75, 954-968.	2.9	19

#	Article	IF	Citations
271	Cigarette smoking, alcohol drinking, and oral cavity and pharyngeal cancer in the Japanese: a population-based cohort study in Japan. European Journal of Cancer Prevention, 2018, 27, 171-179.	1.3	19
272	Coping strategies and cancer incidence and mortality: The Japan Public Health Center-based prospective study. Cancer Epidemiology, 2016, 40, 126-133.	1.9	18
273	Dietary fiber intake and risk of breast cancer defined by estrogen and progesterone receptor status: the Japan Public Health Center-based Prospective Study. Cancer Causes and Control, 2017, 28, 569-578.	1.8	18
274	Plasma levels of n-3 fatty acids and risk of coronary heart disease among Japanese: The Japan Public Health Center-based (JPHC) study. Atherosclerosis, 2018, 272, 226-232.	0.8	18
275	Smoking, Alcohol Consumption, and Risks for Biliary Tract Cancer and Intrahepatic Bile Duct Cancer. Journal of Epidemiology, 2019, 29, 180-186.	2.4	18
276	Coffee, green tea and liver cancer risk: an evaluation based on a systematic review of epidemiologic evidence among the Japanese population. Japanese Journal of Clinical Oncology, 2019, 49, 972-984.	1.3	18
277	Cruciferous vegetable intake and mortality in middle-aged adults: A prospective cohort study. Clinical Nutrition, 2019, 38, 631-643.	5.0	18
278	Intensity-specific validity and reliability of the Japan Public Health Center-based prospective study-physical activity questionnaire. Preventive Medicine Reports, 2020, 20, 101169.	1.8	18
279	Plasma Isoflavones and Risk of Primary Liver Cancer in Japanese Women and Men with Hepatitis Virus Infection: A Nested Case–Control Study. Cancer Epidemiology Biomarkers and Prevention, 2015, 24, 532-537.	2.5	17
280	The Japanese Breast Cancer Society clinical practice guidelines for epidemiology and prevention of breast cancer, 2015 edition. Breast Cancer, 2016, 23, 343-356.	2.9	17
281	Receiver operating characteristic analysis of prediction for gastric cancer development using serum pepsinogen and Helicobacter pylori antibody tests. BMC Cancer, 2017, 17, 183.	2.6	17
282	Effect of body-mass index on the risk of gastric cancer: A population-based cohort study in A Japanese population. Cancer Epidemiology, 2019, 63, 101622.	1.9	17
283	Lowâ€earbohydrate diet and risk of cancer incidence: The Japan Public Health Centerâ€based prospective study. Cancer Science, 2022, 113, 744-755.	3.9	17
284	Body Mass Index and Thyroid Cancer Risk: A Pooled Analysis of Half a Million Men and Women in the Asia Cohort Consortium. Thyroid, 2022, 32, 306-314.	4.5	17
285	C3–6 Laminoplasty for Cervical Spondylotic Myelopathy Maintains Satisfactory Long-Term Surgical Outcomes. Global Spine Journal, 2014, 4, 169-173.	2.3	16
286	Trends in the proportions of stroke subtypes and coronary heart disease in the Japanese men and women from 1995 to 2009. Atherosclerosis, 2016, 248, 219-223.	0.8	16
287	Burden of cancer associated with type 2 diabetes mellitus in Japan, 2010–2030. Cancer Science, 2016, 107, 521-527.	3.9	16
288	Plasma adiponectin levels, ADIPOQ variants, and incidence of type 2 diabetes: A nested case-control study. Diabetes Research and Clinical Practice, 2017, 127, 254-264.	2.8	16

#	Article	IF	Citations
289	Female reproductive factors and risk of all-cause and cause-specific mortality among women: The Japan Public Health Center–based Prospective Study (JPHC study). Annals of Epidemiology, 2018, 28, 597-604.e6.	1.9	16
290	Genome-wide association meta-analysis and Mendelian randomization analysis confirm the influence of ALDH2 on sleep duration in the Japanese population. Sleep, $2019,42,.$	1.1	16
291	Coffee consumption and mortality in Japanese men and women: A pooled analysis of eight population-based cohort studies in Japan (Japan Cohort Consortium). Preventive Medicine, 2019, 123, 270-277.	3.4	16
292	Coffee Drinking and Reduced Risk of Liver Cancer: Update on Epidemiological Findings and Potential Mechanisms. Current Nutrition Reports, 2019, 8, 182-186.	4.3	16
293	Association of BMI and height with the risk of endometrial cancer, overall and by histological subtype: a population-based prospective cohort study in Japan. European Journal of Cancer Prevention, 2019, 28, 196-202.	1.3	16
294	Monoclonal nature of transient abnormal myelopoiesis in Down's syndrome. Blood, 1991, 77, 1161-1163.	1.4	16
295	Association Between Physical Activity and Risk of Disabling Dementia in Japan. JAMA Network Open, 2022, 5, e224590.	5.9	16
296	Validity of self-reported cancer among a Japanese population: Recent results from a population-based prospective study in Japan (JPHC Study). Cancer Epidemiology, 2011, 35, 250-253.	1.9	15
297	Neighborhood contextual factors for smoking among middle-aged Japanese: A multilevel analysis. Health and Place, 2015, 31, 17-23.	3.3	15
298	Development of a risk prediction model for lung cancer: The Japan Public Health Centerâ€based Prospective Study. Cancer Science, 2018, 109, 854-862.	3.9	15
299	Adult height and all-cause and cause-specific mortality in the Japan Public Health Center-based Prospective Study (JPHC). PLoS ONE, 2018, 13, e0197164.	2.5	15
300	Tuberculosis infection and lung adenocarcinoma: Mendelian randomization and pathway analysis of genome-wide association study data from never-smoking Asian women. Genomics, 2020, 112, 1223-1232.	2.9	15
301	Dietary Acrylamide Intake and the Risk of Pancreatic Cancer: The Japan Public Health Center-Based Prospective Study. Nutrients, 2020, 12, 3584.	4.1	15
302	Serum perfluoroalkyl substances and breast cancer risk in Japanese women: A case-control study. Science of the Total Environment, 2021, 800, 149316.	8.0	15
303	Burden of cancer attributable to modifiable factors in Japan in 2015. Global Health & Medicine, 2022, 4, 26-36.	1.4	15
304	The association between complete and partial non-response to psychosocial questions and suicide: the JPHC Study. European Journal of Public Health, 2015, 25, 424-430.	0.3	14
305	Alcohol consumption, genetic variants in the alcohol- and folate metabolic pathways and colorectal cancer risk: the JPHC Study. Scientific Reports, 2016, 6, 36607.	3.3	14
306	Body mass index change during adulthood and risk of oesophageal squamous-cell carcinoma in a Japanese population: the Japan Public Health (JPHC)-based prospective study. British Journal of Cancer, 2017, 117, 1715-1722.	6.4	14

#	Article	IF	CITATIONS
307	Alcohol consumption and bladder cancer risk with or without the flushing response: The Japan Public Health Centerâ€based Prospective Study. International Journal of Cancer, 2017, 141, 2480-2488.	5.1	14
308	Plasma 25-hydroxy vitamin D and subsequent prostate cancer risk in a nested Case-Control study in Japan: The JPHC study. European Journal of Clinical Nutrition, 2017, 71, 132-136.	2.9	14
309	Predictive performance of a genetic risk score using 11 susceptibility alleles for the incidence of Type 2 diabetes in a general Japanese population: a nested case–control study. Diabetic Medicine, 2018, 35, 602-611.	2.3	14
310	Coffee and green tea consumption and subsequent risk of acute myeloid leukemia and myelodysplastic syndromes in Japan. International Journal of Cancer, 2018, 142, 1130-1138.	5.1	14
311	Regional genetic differences among Japanese populations and performance of genotype imputation using whole-genome reference panel of the Tohoku Medical Megabank Project. BMC Genomics, 2018, 19, 551.	2.8	14
312	Re-evaluating genetic variants identified in candidate gene studies of breast cancer risk using data from nearly 280,000 women of Asian and European ancestry. EBioMedicine, 2019, 48, 203-211.	6.1	14
313	Development of scoliosis in young children with osteogenesis imperfecta undergoing intravenous bisphosphonate therapy. Journal of Bone and Mineral Metabolism, 2019, 37, 545-553.	2.7	14
314	Association between serum liver enzymes and allâ€cause mortality: The Japan Public Health Centerâ€based Prospective Study. Liver International, 2019, 39, 1566-1576.	3.9	14
315	Epidemiology of nonmelanoma skin cancer in Japan: Occupational type, lifestyle, and family history of cancer. Cancer Science, 2020, 111, 4257-4265.	3.9	14
316	Variations in the estimated intake of acrylamide from food in the Japanese population. Nutrition Journal, 2020, 19, 17.	3.4	14
317	Identification of a novel uterine leiomyoma GWAS locus in a Japanese population. Scientific Reports, 2020, 10, 1197.	3.3	14
318	Effects of <i>Helicobacter pylori </i> eradication on gastric cancer incidence in the Japanese population: a systematic evidence review. Japanese Journal of Clinical Oncology, 2021, 51, 1158-1170.	1.3	14
319	Relationship between unhealthy sleep status and dry eye symptoms in a Japanese population: The JPHC-NEXT study. Ocular Surface, 2021, 21, 306-312.	4.4	14
320	Fermented and nonfermented soy foods and the risk of breast cancer in a Japanese populationâ€based cohort study. Cancer Medicine, 2021, 10, 757-771.	2.8	14
321	Alcohol Drinking and Total Cancer Risk: An Evaluation Based on a Systematic Review of Epidemiologic Evidence among the Japanese Population. Japanese Journal of Clinical Oncology, 2007, 37, 692-700.	1.3	13
322	Validity of a self-administered food frequency questionnaire in the estimation of heterocyclic aromatic amines. Cancer Causes and Control, 2014, 25, 1015-1028.	1.8	13
323	Risk of lung cancer and consumption of vegetables and fruit in Japanese: A pooled analysis of cohort studies in Japan. Cancer Science, 2015, 106, 1057-1065.	3.9	13
324	Smoking and alcohol and subsequent risk of myelodysplastic syndromes in Japan: the Japan Public Health Centreâ€based Prospective Study. British Journal of Haematology, 2017, 178, 747-755.	2.5	13

#	Article	IF	CITATIONS
325	Risk of thyroid cancer in relation to height, weight, and body mass index in Japanese individuals: a population-based cohort study. Cancer Medicine, 2018, 7, 2200-2210.	2.8	13
326	Dietary Acrylamide Intake and the Risk of Liver Cancer: The Japan Public Health Center-Based Prospective Study. Nutrients, 2020, 12, 2503.	4.1	13
327	Urinary neonicotinoids level among pregnant women in Japan. International Journal of Hygiene and Environmental Health, 2021, 236, 113797.	4.3	13
328	Association of $\langle i \rangle$ Escherichia coli $\langle i \rangle$ containing polyketide synthase in the gut microbiota with colorectal neoplasia in Japan. Cancer Science, 2022, 113, 277-286.	3.9	13
329	A multicenter study of a new inotropic agent, piperanometozine (opc-8212) in congestive heart failure: Clinical improvement during short-term treatment. Cardiovascular Drugs and Therapy, 1987, 1, 169-175.	2.6	12
330	Glycemic index and glycemic load and risk of colorectal cancer: a population-based cohort study (JPHC Study). Cancer Causes and Control, 2016, 27, 583-593.	1.8	12
331	Smoking and subsequent risk of leukemia in Japan: The Japan Public Health Center-based Prospective Study. Journal of Epidemiology, 2017, 27, 305-310.	2.4	12
332	Online version of the self-administered food frequency questionnaire for the Japan Public Health Center-based Prospective Study for the Next Generation (JPHC-NEXT) protocol: Relative validity, usability, and comparison with a printed questionnaire. Journal of Epidemiology, 2017, 27, 435-446.	2.4	12
333	Validity and Reproducibility of a Self-Administered Food Frequency Questionnaire for the Assessment of Sugar Intake in Middle-Aged Japanese Adults. Nutrients, 2019, 11, 554.	4.1	12
334	Neurodevelopmental outcome and respiratory management of congenital central hypoventilation syndrome: a retrospective study. BMC Pediatrics, 2020, 20, 342.	1.7	12
335	Dietary Acrylamide Intake and Risk of Lung Cancer: The Japan Public Health Center Based Prospective Study. Nutrients, 2020, 12, 2417.	4.1	12
336	Dietary Intake of Branched-Chain Amino Acids and Risk of Colorectal Cancer. Cancer Prevention Research, 2020, 13, 65-72.	1.5	12
337	Alcohol consumption and breast cancer risk in Japan: A pooled analysis of eight populationâ€based cohort studies. International Journal of Cancer, 2021, 148, 2736-2747.	5.1	12
338	Dietary Acrylamide Intake and the Risk of Hematological Malignancies: The Japan Public Health Center-Based Prospective Study. Nutrients, 2021, 13, 590.	4.1	12
339	Soy product intake and risk of incident disabling dementia: the JPHC Disabling Dementia Study. European Journal of Nutrition, 2022, 61, 4045-4057.	3.9	12
340	Impact of Lifestyle on Overall Cancer Risk among Japanese: The Japan Public Health Center-Based Prospective Study (JPHC Study). Journal of Epidemiology, 2010, 20, 90-96.	2.4	11
341	Effect of monitoring salt concentration of home-prepared dishes and using low-sodium seasonings on sodium intake reduction. European Journal of Clinical Nutrition, 2018, 72, 1413-1420.	2.9	11
342	Association between plasma concentrations of branched-chain amino acids and adipokines in Japanese adults without diabetes. Scientific Reports, 2018, 8, 1043.	3.3	11

#	Article	IF	CITATIONS
343	Correlation between global methylation level of peripheral blood leukocytes and serum C reactive protein level modified by MTHFR polymorphism: a cross-sectional study. BMC Cancer, 2018, 18, 184.	2.6	11
344	Menstrual and reproductive factors in the risk of thyroid cancer in Japanese women: the Japan Public Health Center-Based Prospective Study. European Journal of Cancer Prevention, 2018, 27, 361-369.	1.3	11
345	Changes in Smoking Status and Mortality From All Causes and Lung Cancer: A Longitudinal Analysis of a Population-based Study in Japan. Journal of Epidemiology, 2019, 29, 11-17.	2.4	11
346	Association of BMI, Smoking, and Alcohol with Multiple Myeloma Mortality in Asians: A Pooled Analysis of More than 800,000 Participants in the Asia Cohort Consortium. Cancer Epidemiology Biomarkers and Prevention, 2019, 28, 1861-1867.	2.5	11
347	Association between educational level and total and cause-specific mortality: a pooled analysis of over 694 000 individuals in the Asia Cohort Consortium. BMJ Open, 2019, 9, e026225.	1.9	11
348	Fruit and vegetable intake and pancreatic cancer risk in a populationâ€based cohort study in Japan. International Journal of Cancer, 2019, 144, 1858-1866.	5.1	11
349	Association of Vegetable, Fruit, and Okinawan Vegetable Consumption With Incident Stroke and Coronary Heart Disease. Journal of Epidemiology, 2020, 30, 37-45.	2.4	11
350	Cross-Sectional Association Between Employment Status and Self-Rated Health Among Middle-Aged Japanese Women: The Influence of Socioeconomic Conditions and Work-Life Conflict. Journal of Epidemiology, 2020, 30, 396-403.	2.4	11
351	Occupational sitting time and subsequent risk of cancer: The Japan Public Health Centerâ€based Prospective Study. Cancer Science, 2020, 111, 974-984.	3.9	11
352	Fat mass and obesity-associated gene polymorphisms, pre-diagnostic plasma adipokine levels and the risk of colorectal cancer: The Japan Public Health Center-based Prospective Study. PLoS ONE, 2020, 15, e0229005.	2.5	11
353	Associations between changes in fruit and vegetable consumption and weight change in Japanese adults. European Journal of Nutrition, 2021, 60, 217-227.	3.9	11
354	Reduction in total and major cause-specific mortality from tobacco smoking cessation: a pooled analysis of 16 population-based cohort studies in Asia. International Journal of Epidemiology, 2022, 50, 2070-2081.	1.9	11
355	Toward a third term of Health Japan 21 – implications from the rise in non-communicable disease burden and highly preventable risk factors. The Lancet Regional Health - Western Pacific, 2022, 21, 100377.	2.9	11
356	A nationally representative crossâ€sectional survey on health information access for consumers in Japan: A protocol for the INFORM Study. World Medical and Health Policy, 2022, 14, 225-275.	1.6	11
357	Impact of five modifiable lifestyle habits on the probability of cancer occurrence in a Japanese population-based cohort: Results from the JPHC study. Preventive Medicine, 2013, 57, 685-689.	3.4	10
358	Smoking and subsequent risk of acute myeloid leukaemia: A pooled analysis of 9 cohort studies in Japan. Hematological Oncology, 2018, 36, 262-268.	1.7	10
359	Coffee Consumption and Lung Cancer Risk: The Japan Public Health Center-Based Prospective Study. Journal of Epidemiology, 2018, 28, 207-213.	2.4	10
360	Fish intake and risk of mortality due to aortic dissection and aneurysm: A pooled analysis of the Japan cohort consortium. Clinical Nutrition, 2019, 38, 1678-1683.	5.0	10

#	Article	IF	Citations
361	Passive smoking and typeÂ2 diabetes among neverâ€smoking women: The Japan Public Health Centerâ€based Prospective Study. Journal of Diabetes Investigation, 2020, 11, 1352-1358.	2.4	10
362	Comparison between the impact of fermented and unfermented soy intake on the risk of liver cancer: the JPHC Study. European Journal of Nutrition, 2021, 60, 1389-1401.	3.9	10
363	Associations of coffee and tea consumption with lung cancer risk. International Journal of Cancer, 2021, 148, 2457-2470.	5.1	10
364	Dietary Acrylamide Intake and the Risks of Renal Cell, Prostate, and Bladder Cancers: A Japan Public Health Center-Based Prospective Study. Nutrients, 2021, 13, 780.	4.1	10
365	Reproductive Factors and Lung Cancer Risk among Never-Smoking Japanese Women with 21 Years of Follow-Up: A Cohort Study. Cancer Epidemiology Biomarkers and Prevention, 2021, 30, 1185-1192.	2.5	10
366	Association between meat and saturated fatty acid intake and lung cancer risk: The Japan Public Health Centerâ€based prospective study. International Journal of Cancer, 2020, 147, 3019-3028.	5.1	10
367	Association between meat intake and mortality due to all-cause and major causes of death in a Japanese population. PLoS ONE, 2020, 15, e0244007.	2.5	10
368	Long-term exposure to fine particle matter and all-cause mortality and cause-specific mortality in Japan: the JPHC Study. BMC Public Health, 2022, 22, 466.	2.9	10
369	Sleep duration and risk of cancer incidence and mortality: A pooled analysis of six populationâ€based cohorts in Japan. International Journal of Cancer, 2022, 151, 1068-1080.	5.1	10
370	Rice, bread, noodle and cereal intake and colorectal cancer in Japanese men and women: the Japan Public Health Center-based prospective Study (JPHC Study). British Journal of Cancer, 2014, 110, 1316-1321.	6.4	9
371	Lung Transplantation for Lymphangioleiomyomatosis in a Single Japanese Institute, With a Focus on Late-onset Complications. Transplantation Proceedings, 2015, 47, 1977-1982.	0.6	9
372	Metabolome analysis for pancreatic cancer risk in nested case ontrol study: Japan Public Health Centerâ€based prospective Study. Cancer Science, 2018, 109, 1672-1681.	3.9	9
373	Meat subtypes and colorectal cancer risk: A pooled analysis of 6 cohort studies in Japan. Cancer Science, 2019, 110, 3603-3614.	3.9	9
374	The functional ALDH2 polymorphism is associated with breast cancer risk: A pooled analysis from the Breast Cancer Association Consortium. Molecular Genetics & Enough Genomic Medicine, 2019, 7, e707.	1.2	9
375	Identification of two novel breast cancer loci through large-scale genome-wide association study in the Japanese population. Scientific Reports, 2019, 9, 17332.	3.3	9
376	The Association Between Habitual Sleep Duration and Mortality According to Sex and Age: The Japan Public Health Center-based Prospective Study. Journal of Epidemiology, 2021, 31, 109-118.	2.4	9
377	Working cancer survivors' physical and mental characteristics compared to cancer-free workers in Japan: a nationwide general population-based study. Journal of Cancer Survivorship, 2021, 15, 912-921.	2.9	9
378	Longâ€term antihypertensive drug use and risk of cancer: The Japan Public Health Centerâ€based prospective study. Cancer Science, 2021, 112, 1997-2005.	3.9	9

#	Article	IF	CITATIONS
379	Association between C-reactive protein and risk of overall and 18 site-specific cancers in a Japanese case-cohort. British Journal of Cancer, 2022, 126, 1481-1489.	6.4	9
380	Association of Marital Status With Total and Cause-Specific Mortality in Asia. JAMA Network Open, 2022, 5, e2214181.	5.9	9
381	Projecting the probability of survival free from cancer and cardiovascular incidence through lifestyle modification in Japan. Preventive Medicine, 2009, 48, 128-133.	3.4	8
382	Synthesis of clinical prediction models under different sets of covariates with one individual patient data. BMC Medical Research Methodology, 2015, 15, 101.	3.1	8
383	Trends in cancer prognosis in a population-based cohort survey: Can recent advances in cancer therapy affect the prognosis?. Cancer Epidemiology, 2015, 39, 97-103.	1.9	8
384	Humoral Immunity to Cytomegalovirus Glycoprotein B in Patients With Breast Cancer and Matched Controls: Contribution of Immunoglobulin \hat{l}^3 , \hat{l}^2 , and Fc \hat{l}^3 Receptor Genes. Journal of Infectious Diseases, 2016, 213, 611-617.	4.0	8
385	Association of plasma C-reactive protein level with the prevalence of colorectal adenoma: the Colorectal Adenoma Study in Tokyo. Scientific Reports, 2017, 7, 4456.	3.3	8
386	Genome-wide association study (GWAS) of ovarian cancer in Japanese predicted regulatory variants in 22q13.1. PLoS ONE, 2018, 13, e0209096.	2.5	8
387	Higher Dietary Non-enzymatic Antioxidant Capacity Is Associated with Decreased Risk of All-Cause and Cardiovascular Disease Mortality in Japanese Adults. Journal of Nutrition, 2019, 149, 1967-1976.	2.9	8
388	Relationship between dietary non-enzymatic antioxidant capacity and type 2 diabetes risk in the Japan Public Health Center-based Prospective Study. Nutrition, 2019, 66, 62-69.	2.4	8
389	Soy food and isoflavones are not associated with changes in serum lipids and glycohemoglobin concentrations among Japanese adults: a cohort study. European Journal of Nutrition, 2020, 59, 2075-2087.	3.9	8
390	Doneness preferences, meat and meat-derived heterocyclic amines intake, and N-acetyltransferase 2 polymorphisms: association with colorectal adenoma in Japanese Brazilians. European Journal of Cancer Prevention, 2020, 29, 7-14.	1.3	8
391	Inclusion of a geneâ€environment interaction between alcohol consumption and the aldehyde dehydrogenase 2 genotype in a risk prediction model for upper aerodigestive tract cancer in Japanese men. Cancer Science, 2020, 111, 3835-3844.	3.9	8
392	Dietary fiber intake and risk of gastric cancer: The <scp>Japan Public Health Center</scp> â€based prospective study. International Journal of Cancer, 2021, 148, 2664-2673.	5.1	8
393	Relation Between Body Mass Index and Dry Eye Disease: The Japan Public Health Center–Based Prospective Study for the Next Generation. Eye and Contact Lens, 2021, 47, 449-455.	1.6	8
394	Public Health Interventions for Gastric Cancer Control. Gastrointestinal Endoscopy Clinics of North America, 2021, 31, 441-449.	1.4	8
395	Association of dietary intakes of vitamin B12, vitamin B6, folate, and methionine with the risk of esophageal cancer: the Japan Public Health Center-based (JPHC) prospective study. BMC Cancer, 2021, 21, 982.	2.6	8
396	Comparison of the Japanese Orthopaedic Association Score and the Japanese Orthopaedic Association Cervical Myelopathy Evaluation Questionnaire Scores: Time-Dependent Changes in Patients with Cervical Spondylotic Myelopathy and Posterior Longitudinal Ligament. Asian Spine Journal, 2015, 9, 47.	2.0	8

#	Article	IF	Citations
397	Sugary drink consumption and risk of kidney and bladder cancer in Japanese adults. Scientific Reports, 2021, 11, 21701.	3.3	8
398	Association between body mass index and oesophageal cancer mortality: a pooled analysis of prospective cohort studies with >800 000 individuals in the Asia Cohort Consortium. International Journal of Epidemiology, 2022, 51, 1190-1203.	1.9	8
399	Three cases of lupus nephritis patients with serum interleukin-32γ detection. Lupus, 2014, 23, 1187-1191.	1.6	7
400	Smoking is a risk factor for development of adult T-cell leukemia/lymphoma in Japanese human T-cell leukemia virus type-1 carriers. Cancer Causes and Control, 2016, 27, 1059-1066.	1.8	7
401	Comparison of weighed food record procedures for the reference methods in two validation studies of food frequency questionnaires. Journal of Epidemiology, 2017, 27, 331-337.	2.4	7
402	Meta-regression Analysis of Sex- and Birth Year-Specific Prevalence of HBsAg and Anti-HCV Among Un-diagnosed Japanese: Data From the First-time Blood Donors, Periodical Health Checkup, and the Comprehensive Health Checkup With Lifestyle Education (Ningen Dock). Journal of Epidemiology, 2020, 30, 420-425.	2.4	7
403	Association between dietary sugar intake and colorectal adenoma among cancer screening examinees in Japan. Cancer Science, 2020, 111, 3862-3872.	3.9	7
404	Quantifying the association of low-intensity and late initiation of tobacco smoking with total and cause-specific mortality in Asia. Tobacco Control, 2021, 30, 328-335.	3.2	7
405	Sugary Drink Consumption and Subsequent Colorectal Cancer Risk: The Japan Public Health Center–Based Prospective Cohort Study. Cancer Epidemiology Biomarkers and Prevention, 2021, 30, 782-788.	2.5	7
406	Alcohol consumption, tobacco smoking, and subsequent risk of renal cell carcinoma: The JPHC study. Cancer Science, 2021, 112, 5068-5077.	3.9	7
407	Comparison of postmenopausal endogenous sex hormones among Japanese, Japanese Brazilians, and non-Japanese Brazilians. BMC Medicine, 2011, 9, 16.	5.5	6
408	Physical activity and subsequent risk of kidney, bladder and upper urinary tract cancer in the Japanese population: the Japan Public Health Centre-based Prospective Study. British Journal of Cancer, 2019, 120, 571-574.	6.4	6
409	Cruciferous vegetable intake and colorectal cancer risk: Japan public health center-based prospective study. European Journal of Cancer Prevention, 2019, 28, 420-427.	1.3	6
410	Family history of cancer and subsequent risk of cancer: A largeâ€scale populationâ€based prospective study in Japan. International Journal of Cancer, 2020, 147, 331-337.	5.1	6
411	Soy and isoflavone consumption and subsequent risk of prostate cancer mortality: the Japan Public Health Center-based Prospective Study. International Journal of Epidemiology, 2020, 49, 1553-1561.	1.9	6
412	Metabolic Syndrome, Physical Activity, and Inflammation: A Cross-Sectional Analysis of 110 Circulating Biomarkers in Japanese Adults. Cancer Epidemiology Biomarkers and Prevention, 2020, 29, 1639-1646.	2.5	6
413	Intake of Vegetables and Fruits and the Risk of Cataract Incidence in a Japanese Population: The Japan Public Health Center-Based Prospective Study. Journal of Epidemiology, 2021, 31, 21-29.	2.4	6
414	Association Between Birth Weight and Risk of Pregnancy-Induced Hypertension and Gestational Diabetes in Japanese Women: JPHC-NEXT Study. Journal of Epidemiology, 2022, 32, 168-173.	2.4	6

#	Article	IF	CITATIONS
415	OUP accepted manuscript. International Journal of Epidemiology, 2021, , .	1.9	6
416	Functional annotation of the 2q35 breast cancer risk locus implicates a structural variant in influencing activity of a long-range enhancer element. American Journal of Human Genetics, 2021, 108, 1190-1203.	6.2	6
417	Dietary heterocyclic aromatic amine intake and cancer risk: epidemiological evidence from Japanese studies. Genes and Environment, 2021, 43, 33.	2.1	6
418	Prediagnostic circulating inflammation-related biomarkers and gastric cancer: A case-cohort study in Japan. Cytokine, 2021, 144, 155558.	3.2	6
419	Food frequency questionnaire reproducibility for middle-aged and elderly Japanese. Asia Pacific Journal of Clinical Nutrition, 2019, 28, 362-370.	0.4	6
420	Meat consumption and gastric cancer risk: The Japan Public Health Center-based Prospective Study. American Journal of Clinical Nutrition, 2021, , .	4.7	6
421	Burden of cancer attributable to tobacco smoke in Japan in 2015. GHM Open, 2021, 1, 43-50.	0.6	6
422	Inverse Association between Fruit and Vegetable Intake and All-Cause Mortality: Japan Public Health Center-Based Prospective Study. Journal of Nutrition, 2022, 152, 2245-2254.	2.9	6
423	Prediagnostic plasma polyphenol concentrations and colon cancer risk: The JPHC nested case–control study. Clinical Nutrition, 2022, 41, 1950-1960.	5.0	6
424	History of Having a Macrosomic Infant and the Risk of Diabetes: The Japan Public Health Center-Based Prospective Diabetes Study. PLoS ONE, 2013, 8, e84542.	2.5	5
425	Fruit and vegetable intake and the risk of overall cancer in Japanese: AÂpooled analysis of population-based cohort studies. Journal of Epidemiology, 2017, 27, 152-162.	2.4	5
426	Coffee and Green Tea Consumption and Subsequent Risk of Malignant Lymphoma and Multiple Myeloma in Japan: The Japan Public Health Center-based Prospective Study. Cancer Epidemiology Biomarkers and Prevention, 2017, 26, 1352-1356.	2.5	5
427	High serum total cholesterol is associated with suicide mortality in Japanese women independently of menopause. Acta Psychiatrica Scandinavica, 2018, 137, 80-81.	4.5	5
428	Two truncating variants in FANCC and breast cancer risk. Scientific Reports, 2019, 9, 12524.	3.3	5
429	Female reproductive factors and risk of lymphoid neoplasm: The Japan Public Health Centerâ€based Prospective Study. Cancer Science, 2019, 110, 1442-1452.	3.9	5
430	Plasma Câ€peptide and glycated albumin and subsequent risk of cancer: From a large prospective caseâ€cohort study in Japan. International Journal of Cancer, 2019, 144, 718-729.	5.1	5
431	Soy Intake and Colorectal Cancer Risk: Results from a Pooled Analysis of Prospective Cohort Studies Conducted in China and Japan. Journal of Nutrition, 2020, 150, 2442-2450.	2.9	5
432	Estimation of the performance of a risk prediction model for gastric cancer occurrence in Japan: Evidence from a small external population. Cancer Epidemiology, 2020, 67, 101766.	1.9	5

#	Article	IF	CITATIONS
433	Dietary glycemic index, glycemic load, and endometrial cancer risk: The Japan Public Health Centerâ€based Prospective Study. Cancer Science, 2021, 112, 3682-3690.	3.9	5
434	Association of sugary drink consumption with all-cause and cause-specific mortality: the Japan Public Health Center-based Prospective Study. Preventive Medicine, 2021, 148, 106561.	3.4	5
435	Reliability of self-reported questionnaire for epidemiological investigation of Helicobacter pylori eradication in a population-based cohort study. Scientific Reports, 2021, 11, 15605.	3. 3	5
436	Association between sugar and starch intakes and type 2 diabetes risk in middle-aged adults in a prospective cohort study. European Journal of Clinical Nutrition, 2022, 76, 746-755.	2.9	5
437	Peanut Consumption and Risk of Stroke and Ischemic Heart Disease in Japanese Men and Women: The JPHC Study. Stroke, 2021, 52, 3543-3550.	2.0	5
438	Burden of cancer attributable to consumption of alcohol in Japan in 2015. GHM Open, 2021, 1, 51-55.	0.6	5
439	Hobby Engagement and Risk of Disabling Dementia. Journal of Epidemiology, 2023, 33, 456-463.	2.4	5
440	Association between Alcohol Consumption and Colorectal Cancer Risk. Current Nutrition Reports, 2013, 2, 71-73.	4.3	4
441	Comparison of plasma levels of nutrient-related biomarkers among Japanese populations in Tokyo, Japan, São Paulo, Brazil, and Hawaii, USA. European Journal of Cancer Prevention, 2015, 24, 155-161.	1.3	4
442	The Validity and Reproducibility of Dietary Non-enzymatic Antioxidant Capacity Estimated by Self-administered Food Frequency Questionnaires. Journal of Epidemiology, 2018, 28, 428-436.	2.4	4
443	IGHG, IGKC, and FCGR genes and endogenous antibody responses to GARP in patients with breast cancer and matched controls. Human Immunology, 2018, 79, 632-637.	2.4	4
444	Exploring predictive biomarkers from clinical genome-wide association studies via multidimensional hierarchical mixture models. European Journal of Human Genetics, 2019, 27, 140-149.	2.8	4
445	Relationship between Meat/Fish Consumption and Biliary Tract Cancer: The Japan Public Health Center–Based Prospective Study. Cancer Epidemiology Biomarkers and Prevention, 2020, 29, 95-102.	2.5	4
446	Soy Food Intake and Pancreatic Cancer Risk: The Japan Public Health Center–based Prospective Study. Cancer Epidemiology Biomarkers and Prevention, 2020, 29, 1214-1221.	2.5	4
447	Validation Study of Diabetes Definitions Using Japanese Diagnosis Procedure Combination Data Among Hospitalized Patients. Journal of Epidemiology, 2023, 33, 165-169.	2.4	4
448	Risk Stratification Score Improves Sensitivity for Advanced Colorectal Neoplasia in Colorectal Cancer Screening: The Oshima Study Workgroup. Clinical and Translational Gastroenterology, 2021, 12, e00319.	2. 5	4
449	Body Mass Index, Height, Weight Change, and Subsequent Lung Cancer Risk: The Japan Public Health Center–Based Prospective Study. Cancer Epidemiology Biomarkers and Prevention, 2021, 30, 1708-1716.	2.5	4
450	A Personal Breast Cancer Risk Stratification Model Using Common Variants and Environmental Risk Factors in Japanese Females. Cancers, 2021, 13, 3796.	3.7	4

#	Article	IF	CITATIONS
451	Burden of cancer attributable to infection in Japan in 2015. GHM Open, 2021, 1, 63-69.	0.6	4
452	Vegetable and fruit intake and the risk of bladder cancer: Japan Public Health Center-based prospective study. British Journal of Cancer, 2022, 126, 1647-1658.	6.4	4
453	Three dimensional analysis of abnormal filaments of Pick's disease by scanning electron microscopy. Neuropathology and Applied Neurobiology, 1997, 23, 326-330.	3.2	3
454	Diabetes mellitus defined by hemoglobin A1c value: Risk characterization for incidence among Japanese subjects in the JPHC Diabetes Study. Journal of Diabetes Investigation, 2011, 2, 359-365.	2.4	3
455	Privacy-Preserving Hypothesis Testing for Reduced Cancer Risk on Daily Physical Activity. Journal of Medical Systems, 2018, 42, 90.	3.6	3
456	Endogenous antibody responses to mucin 1 in a large multiethnic cohort of patients with breast cancer and healthy controls: Role of immunoglobulin and $Fc\hat{l}^3$ receptor genes. Immunobiology, 2018, 223, 178-182.	1.9	3
457	The association between plasma C-peptide concentration and the risk of prostate cancer: a nested case–control study within a Japanese population-based prospective study. European Journal of Cancer Prevention, 2018, 27, 461-467.	1.3	3
458	Female reproductive factors and risk of external causes of death among women: The Japan Public Health Center-based Prospective Study (JPHC Study). Scientific Reports, 2019, 9, 14329.	3.3	3
459	Lack of social support and social trust as potential risk factors for dry eye disease: JPHC-NEXT study. Ocular Surface, 2019, 17, 278-284.	4.4	3
460	Impact of alcohol drinking on cancer risk with consideration of flushing response: The Japan Public Health Center-based Prospective Study Cohort (JPHC study). Preventive Medicine, 2020, 133, 106026.	3.4	3
461	Effectiveness of Screening Using Fecal Occult Blood Testing and Colonoscopy on the Risk of Colorectal Cancer: The Japan Public Health Center-based Prospective Study. Journal of Epidemiology, 2023, 33, 91-100.	2.4	3
462	Apolipoprotein A2 Isoforms in Relation to the Risk of Myocardial Infarction: A Nested Case-Control Analysis in the JPHC Study. Journal of Atherosclerosis and Thrombosis, 2021, 28, 483-490.	2.0	3
463	Long-term Response of <i>Helicobacter pylori</i> Antibody Titer After Eradication Treatment in Middle-aged Japanese: JPHC-NEXT Study. Journal of Epidemiology, 2023, 33, 1-7.	2.4	3
464	Exploratory Research on Determinants of Place of Death in a Large-scale Cohort Study: The JPHC Study. Journal of Epidemiology, 2023, 33, 120-126.	2.4	3
465	Alcohol intake and stomach cancer risk in Japan: A pooled analysis of six cohort studies. Cancer Science, 2022, 113, 261-276.	3.9	3
466	Association between coffee consumption and risk of prostate cancer in Japanese men: a population-based cohort study in Japan. Cancer Epidemiology Biomarkers and Prevention, 2021, , cebp.0484.2021.	2.5	3
467	Association of B Vitamins and Methionine Intake with the Risk of Gastric Cancer: The Japan Public Health Center–based Prospective Study. Cancer Prevention Research, 2022, 15, 101-110.	1.5	3
468	Burden of cancer attributable to excess bodyweight and physical inactivity in Japan in 2015. CHM Open, 2021, 1, 56-62.	0.6	3

#	Article	IF	CITATIONS
469	Burden of cancer attributable to insufficient vegetable, fruit and dietary fiber consumption in Japan in 2015. GHM Open, 2021, 1, 70-75.	0.6	3
470	Association between Meat, Fish, and Fatty Acid Intake and Non-Hodgkin Lymphoma Incidence: The Japan Public Health Center–Based Prospective Study. Journal of Nutrition, 2022, 152, 1895-1906.	2.9	3
471	Effect of Respiratory Rate on Respiratory Patterns in Patients with Chronic Obstructive Pulmonary Disease Internal Medicine, 1997, 36, 771-775.	0.7	2
472	Comparison of plasma levels of obesity-related biomarkers among Japanese populations in Tokyo, Japan, São Paulo, Brazil, and Hawaii, USA. European Journal of Cancer Prevention, 2016, 25, 41-49.	1.3	2
473	Alcohol Drinking and Bladder Cancer Risk From a Pooled Analysis of Ten Cohort Studies in Japan. Journal of Epidemiology, 2020, 30, 309-313.	2.4	2
474	Germline HOXB13 mutations p.G84E and p.R217C do not confer an increased breast cancer risk. Scientific Reports, 2020, 10, 9688.	3.3	2
475	Risk of stroke in cancer survivors using a propensity score-matched cohort analysis. Scientific Reports, 2021, 11, 5599.	3.3	2
476	Impact of reproductive factors on breast cancer incidence: Pooled analysis of nine cohort studies in Japan. Cancer Medicine, 2021, 10, 2153-2163.	2.8	2
477	Effectiveness of a Cancer Risk Prediction Tool on Lifestyle Habits: A Randomized Controlled Trial. Cancer Epidemiology Biomarkers and Prevention, 2021, 30, 1063-1071.	2.5	2
478	Dietary glycemic index, glycemic load and mortality: Japan Public Health Center-based prospective study. European Journal of Nutrition, 2021, 60, 4607-4620.	3.9	2
479	International strategy in cancer epidemiology: Japan's involvement in global projects and future role. Global Health & Medicine, 2021, 3, 187-195.	1.4	2
480	Circulating Inflammation Markers and Pancreatic Cancer Risk: A Prospective Case-Cohort Study in Japan. Cancer Epidemiology Biomarkers and Prevention, 2022, 31, 236-241.	2.5	2
481	Burden of cancer attributable to exogenous hormone use in Japan in 2015. GHM Open, 2021, 1, 97-101.	0.6	2
482	A Predictive Model of Noncardia Gastric Adenocarcinoma Risk Using Antibody Response to <i>Helicobacter pylori</i> Proteins and Pepsinogen. Cancer Epidemiology Biomarkers and Prevention, 2022, 31, 811-820.	2.5	2
483	Burden of cancer attributable to consumption of highly salted food in Japan in 2015. GHM Open, 2021, 1, 85-90.	0.6	2
484	Burden of cancer attributable to excess red and processed meat consumption in Japan in 2015. GHM Open, 2021, 1, 91-96.	0.6	2
485	Burden of cancer attributable to never breastfeeding in Japan in 2015. GHM Open, 2021, 1, 102-105.	0.6	2
486	Burden of cancer attributable to air pollution in Japan in 2015. GHM Open, 2021, 1, 76-84.	0.6	2

#	Article	IF	CITATIONS
487	Dietary fibre intake is associated with reduced risk of lung cancer: a Japan public health centre-based prospective study (JPHC). International Journal of Epidemiology, 2022, 51, 1142-1152.	1.9	2
488	Adult height in relation to the risk of colorectal cancer among the Japanese population: an evaluation based on systematic review and meta-analysis. Japanese Journal of Clinical Oncology, 2022, 52, 322-330.	1.3	2
489	Subclinical Chronic Atlanto-Occipital Rotatory Fixation. JBJS Case Connector, 2012, 2, e41.	0.3	1
490	Commentary: Factors Associated With Non-participation in Cohort Studies Emphasize the Need to Generalize the Results With Care. Journal of Epidemiology, 2015, 25, 89-90.	2.4	1
491	Body mass index and height in relation to brain tumor risk in a Japanese population. Annals of Epidemiology, 2020, 51, 1-6.	1.9	1
492	Impact of reduced smoking rates on lung cancer screening programs in Japan. Japanese Journal of Clinical Oncology, 2020, 50, 1126-1132.	1.3	1
493	Validity of dietary isothiocyanate intake estimates from a food frequency questionnaire using 24 h urinary isothiocyanate excretion as an objective biomarker: the JPHC-NEXT protocol area. European Journal of Clinical Nutrition, 2021, , .	2.9	1
494	Total, animal, and plant protein intake and pneumonia mortality in the Japan Public Health Center–based Prospective Study. American Journal of Clinical Nutrition, 2022, 115, 781-789.	4.7	1
495	Cross-sectional associations between the types/amounts of beverages consumed and the glycemia status: The Japan public health center-based Prospective Diabetes study. Metabolism Open, 2022, 14, 100185.	2.9	1
496	Relevance of the MHC region for breast cancer susceptibility in Asians. Breast Cancer, 2022, 29, 869-879.	2.9	1
497	Association of Plasma Iron Status with Subsequent Risk of Total and Site-Specific Cancer: A Large Case–Cohort Study within JPHC Study. Cancer Prevention Research, 2022, 15, 669-678.	1.5	1
498	Reply to Brockton's letter. Cancer Causes and Control, 2008, 19, 1003-1003.	1.8	0
499	P1-348 Leisure-time physical activity and breast cancer risk defined by oestrogen and progesterone receptor status: the Japan public health center-based prospective study. Journal of Epidemiology and Community Health, 2011, 65, A163-A163.	3.7	0
500	P2-338 Association of body mass index with risk of age-related cataracts in a middle-aged Japanese population. Journal of Epidemiology and Community Health, 2011, 65, A315-A316.	3.7	0
501	The Establishment of the Household Air Pollution Consortium (HAPCO). Atmosphere, 2019, 10, 422.	2.3	0
502	Moving towards tailored, region-specific cancer-control measures in China. The Lancet Global Health, 2019, 7, e175-e176.	6.3	0
503	Body mass index and mortality among middle-aged Japanese individuals with diagnosed diabetes: The Japan Public Health Center-based prospective study (JPHC study). Diabetes Research and Clinical Practice, 2020, 164, 108198.	2.8	0
504	Low <i>MICA </i> gene expression confers an increased risk of Graves' disease: a Mendelian randomization study. Thyroid, 2021, , .	4.5	0

#	Article	IF	CITATIONS
505	Applicability of a web-based 24-hour dietary recall tool for Japanese populations in large-scale epidemiological studies. Journal of Epidemiology, 2022, , .	2.4	O
506	COT-6 Body mass index and height in relation to brain tumor risk in a Japanese population. Neuro-Oncology Advances, 2021, 3, vi29-vi29.	0.7	0
507	Title is missing!. , 2020, 15, e0244007.		0
508	Title is missing!. , 2020, 15, e0244007.		0
509	Title is missing!. , 2020, 15, e0244007.		0
510	Title is missing!. , 2020, 15, e0244007.		0