

# Kazuhiko Nakagawa

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7796081/publications.pdf>

Version: 2024-02-01

166  
papers

24,228  
citations

44444

50  
h-index

8627

151  
g-index

168  
all docs

168  
docs citations

168  
times ranked

18146  
citing authors

| #  | ARTICLE  | IF   | CITATIONS |
|----|--|------|-----------|
| 1  | Trastuzumab Deruxtecan in <i>HER2</i> -Mutant Non-Small-Cell Lung Cancer. <i>New England Journal of Medicine</i> , 2022, 386, 241-251.   | 13.9 | 393       |
| 2  | Phase III Clinical Trial for the Combination of Erlotinib Plus Ramucirumab Compared With Osimertinib in Previously Untreated Advanced or Recurrent Non-Small Cell Lung Cancer Positive for the L858R Mutation of EGFR: REVOL858R (WJOG14420L). <i>Clinical Lung Cancer</i> , 2022, 23, e257-e263.                | 1.1  | 10        |
| 3  | Real-world data on NGS using the OncoPrint DxTT for detecting genetic alterations in non-small cell lung cancer: WJOG13019L. <i>Cancer Science</i> , 2022, 113, 221-228.   | 1.7  | 31        |
| 4  | Randomized Phase III Study of Gefitinib Versus Cisplatin Plus Vinorelbine for Patients With Resected Stage II-III A Non-Small-Cell Lung Cancer With <i>EGFR</i> Mutation (IMPACT). <i>Journal of Clinical Oncology</i> , 2022, 40, 231-241.  | 0.8  | 61        |
| 5  | Association of tumour burden with the efficacy of programmed cell death-1/programmed cell death ligand-1 inhibitors for treatment-naïve advanced non-small-cell lung cancer. <i>European Journal of Cancer</i> , 2022, 161, 44-54.   | 1.3  | 7         |
| 6  | Phase II study of atezolizumab with bevacizumab for non-squamous non-small cell lung cancer with high PD-L1 expression (@Be Study)., 2022, 10, e004025.  |      | 22        |
| 7  | RELAY, Ramucirumab Plus Erlotinib Versus Placebo Plus Erlotinib in Patients with Untreated, Epidermal Growth Factor Receptor Mutation-Positive, Metastatic Non-Small-Cell Lung Cancer: Safety Profile and Manageability. <i>Drug Safety</i> , 2022, 45, 45-64.   | 1.4  | 6         |
| 8  | A Randomized Phase II Study Comparing Nivolumab with Carboplatin+Pemetrexed for <i>EGFR</i> -Mutated NSCLC with Resistance to EGFR Tyrosine Kinase Inhibitors (WJOG8515L). <i>Clinical Cancer Research</i> , 2022, 28, 893-902.  | 3.2  | 35        |
| 9  | RELAY+: Exploratory Study of Ramucirumab Plus Gefitinib in Untreated Patients With EGFR-Mutated Metastatic NSCLC. <i>JTO Clinical and Research Reports</i> , 2022, 3, 100303.  | 0.6  | 1         |
| 10 | HER3 Augmentation via Blockade of EGFR/AKT Signaling Enhances Anticancer Activity of HER3-Targeting Patritumab Deruxtecan in EGFR-Mutated Non-Small Cell Lung Cancer. <i>Clinical Cancer Research</i> , 2022, 28, 390-403.   | 3.2  | 34        |
| 11 | Alternating Therapy with Osimertinib and Afatinib for Treatment-Naïve Patients with EGFR-Mutated Advanced Non-Small Cell Lung Cancer: A Single-Group, Open-Label Phase 2 Trial (WJOG10818L). <i>Lung Cancer</i> , 2022, 168, 38-45.  | 0.9  | 5         |
| 12 | The significance of micro-EGFR T790M mutation on EGFR-TKI efficacy in patients with NSCLC: The WJOG13119L study.. <i>Journal of Clinical Oncology</i> , 2022, 40, e21177-e21177.   | 0.8  | 1         |
| 13 | Dynamics of HER3 and its correlated gene expression profile in EGFR-mutated NSCLC tumor treated with EGFR-TKI toward enhancing effectiveness of patritumab deruxtecan (HER3-DXd; U3-1402).. <i>Journal of Clinical Oncology</i> , 2022, 40, e21175-e21175.   | 0.8  | 0         |
| 14 | EORTC-1416-LCG/ETOP 8-15 + PEARLS/KEYNOTE-091 study of pembrolizumab versus placebo for completely resected early-stage non-small cell lung cancer (NSCLC): Outcomes in subgroups related to surgery, disease burden, and adjuvant chemotherapy use.. <i>Journal of Clinical Oncology</i> , 2022, 40, 8512-8512. | 0.8  | 14        |
| 15 | Nivolumab Retreatment in Non-Small Cell Lung Cancer Patients Who Responded to Prior Immune Checkpoint Inhibitors and Had ICI-Free Intervals (WJOG9616L). <i>Clinical Cancer Research</i> , 2022, 28, 3207-3213.  | 3.2  | 7         |
| 16 | RELAY, ramucirumab plus erlotinib versus placebo plus erlotinib in untreated EGFR-mutated metastatic non-small cell lung cancer: exposure-response relationship. <i>Cancer Chemotherapy and Pharmacology</i> , 2022, 90, 137-148.  | 1.1  | 4         |
| 17 | Evaluation of pembrolizumab monotherapy in patients with previously treated advanced salivary gland carcinoma in the phase 2 KEYNOTE-158 study. <i>European Journal of Cancer</i> , 2022, 171, 259-268.  | 1.3  | 19        |
| 18 | Optimizing antiemetic treatment for chemotherapy-induced nausea and vomiting in Japan: Update summary of the 2015 Japan Society of Clinical Oncology Clinical Practice Guidelines for Antiemesis. <i>International Journal of Clinical Oncology</i> , 2021, 26, 1-17.  | 1.0  | 56        |

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 19 | Five-year follow-up results from phase II studies of nivolumab in Japanese patients with previously treated advanced non-small cell lung cancer: pooled analysis of the ONO-4538-05 and ONO-4538-06 studies. <i>Japanese Journal of Clinical Oncology</i> , 2021, 51, 106-113.  | 0.6 | 8         |
| 20 | Dual EGFR-VEGF Pathway Inhibition: A Promising Strategy for Patients With EGFR-Mutant NSCLC. <i>Journal of Thoracic Oncology</i> , 2021, 16, 205-215.   | 0.5 | 149       |
| 21 | Phase I/ II Study of Cisplatin plus NabêPaclitaxel with Concurrent Thoracic Radiotherapy for Patients with Locally Advanced NonêSmall Cell Lung Cancer. <i>Oncologist</i> , 2021, 26, 19.   | 1.9 | 1         |
| 22 | The patientâ€™s perspective on treatment with dacomitinib: patient-reported outcomes from the Phase III trial ARCHER 1050. <i>Future Oncology</i> , 2021, 17, 783-794.  | 1.1 | 0         |
| 23 | Brigatinib in Japanese Patients With ALK-Positive NSCLC Previously Treated With Alectinib and Other Tyrosine Kinase Inhibitors: Outcomes of the Phase 2 J-ALTA Trial. <i>Journal of Thoracic Oncology</i> , 2021, 16, 452-463.  | 0.5 | 51        |
| 24 | Indirect analysis of first-line therapy for advanced non-small-cell lung cancer with activating mutations in a Japanese population. <i>Future Oncology</i> , 2021, 17, 103-115.   | 1.1 | 1         |
| 25 | Updated Overall Survival in a Randomized Study Comparing Dacomitinib with Gefitinib as First-Line Treatment in Patients with Advanced Non-Small-Cell Lung Cancer and EGFR-Activating Mutations. <i>Drugs</i> , 2021, 81, 257-266.   | 4.9 | 57        |
| 26 | Predicting osimertinibâ€™treatment outcomes through <i>EGFR</i> mutantâ€™fraction monitoring in the circulating tumor DNA of <i>EGFR</i> T790Mâ€™positive patients with nonêsmall cell lung cancer (WJOG8815L). <i>Molecular Oncology</i> , 2021, 15, 126-137.  | 2.1 | 12        |
| 27 | Clinical Efficacy and Safety of Nivolumab in Japanese Patients With Malignant Pleural Mesothelioma: 3-Year Results of the MERIT Study. <i>JTO Clinical and Research Reports</i> , 2021, 2, 100135.  | 0.6 | 9         |
| 28 | Efficacy of Osimertinib Plus Bevacizumab vs Osimertinib in Patients With <i>EGFR</i> T790Mâ€™Mutated NonêSmall Cell Lung Cancer Previously Treated With Epidermal Growth Factor Receptorâ€™Tyrosine Kinase Inhibitor. <i>JAMA Oncology</i> , 2021, 7, 386.  | 3.4 | 108       |
| 29 | Safety and efficacy of first-line dacomitinib in Asian patients with EGFR mutation-positive non-small cell lung cancer: Results from a randomized, open-label, phase 3 trial (ARCHER 1050). <i>Lung Cancer</i> , 2021, 154, 176-185.  | 0.9 | 18        |
| 30 | Brigatinib in Japanese patients with anaplastic lymphoma kinase (ALK)-positive non-small cell lung cancer (NSCLC): First results from the J-ALTA tyrosine kinase inhibitor (TKI)-naïve expansion cohort.. <i>Journal of Clinical Oncology</i> , 2021, 39, 9042-9042.  | 0.8 | 3         |
| 31 | A randomized phase II study comparing nivolumab (NIVO) with carboplatin-pemetrexed (CbPEM) for patients (pts) with EGFR mutation-positive non-small cell lung cancer (NSCLC) who acquire resistance to tyrosine kinase inhibitors (TKIs) not due to a secondary T790M mutation (WJOG8515L).. <i>Journal of Clinical Oncology</i> , 2021, 39, 9037-9037. | 0.8 | 5         |
| 32 | Challenges in lung cancer multidisciplinary collaboration experienced by specialists in four countries.. <i>Journal of Clinical Oncology</i> , 2021, 39, e23002-e23002.   | 0.8 | 0         |
| 33 | Efficacy and safety of pembrolizumab in patients with advanced mesothelioma in the open-label, single-arm, phase 2 KEYNOTE-158 study. <i>Lancet Respiratory Medicine</i> ,the, 2021, 9, 613-621.  | 5.2 | 44        |
| 34 | Phase 2 Study of YS110, a Recombinant Humanized Anti-CD26 Monoclonal Antibody, in Japanese Patients With Advanced Malignant Pleural Mesothelioma. <i>JTO Clinical and Research Reports</i> , 2021, 2, 100178.   | 0.6 | 4         |
| 35 | Ramucirumab Plus Erlotinib Versus Placebo Plus Erlotinib in Patients With Untreated Metastatic EGFR-Mutated NSCLC: RELAY Japanese Subset. <i>JTO Clinical and Research Reports</i> , 2021, 2, 100171.   | 0.6 | 5         |
| 36 | Implementation of clinical sequencing for molecular profiling in patients with advanced cancer. <i>Cancer Biomarkers</i> , 2021, 31, 119-126.   | 0.8 | 1         |

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 37 | RELAY Subgroup Analyses by EGFR Ex19del and Ex21L858R Mutations for Ramucirumab Plus Erlotinib in Metastatic Nonâ€“Small Cell Lung Cancer. <i>Clinical Cancer Research</i> , 2021, 27, 5258-5271.   | 3.2 | 23        |
| 38 | Pembrolizumab Plus Amrubicin in Patients With Relapsed SCLC: Multi-Institutional, Single-Arm Phase 2 Study. <i>JTO Clinical and Research Reports</i> , 2021, 2, 100184.   | 0.6 | 8         |
| 39 | KRAS Inhibitor Resistance in <i>MET</i> -Amplified <i>KRAS</i> G12C Nonâ€“Small Cell Lung Cancer Induced By RAS- and Nonâ€“RAS-Mediated Cell Signaling Mechanisms. <i>Clinical Cancer Research</i> , 2021, 27, 5697-5707.                                   | 3.2 | 42        |
| 40 | Firstâ€“line pembrolizumab vs chemotherapy in metastatic nonâ€“smallâ€“cell lung cancer: KEYNOTEâ€“024 Japan subset*. <i>Cancer Science</i> , 2021, 112, 5000-5010.   | 1.7 | 6         |
| 41 | Realâ€“world safety of nivolumab in patients with nonâ€“smallâ€“cell lung cancer in Japan: Postmarketing surveillance. <i>Cancer Science</i> , 2021, 112, 4692-4701.  | 1.7 | 14        |
| 42 | Prognostic impact of geriatric assessment in elderly patients with non-small cell lung cancer: an integrated analysis of two randomized phase III trials (JCOG1115-A). <i>Japanese Journal of Clinical Oncology</i> , 2021, 51, 685-692.                    | 0.6 | 1         |
| 43 | Intestinal Microbiota and Gene Expression Reveal Similarity and Dissimilarity Between Immune-Mediated Colitis and Ulcerative Colitis. <i>Frontiers in Oncology</i> , 2021, 11, 763468.  | 1.3 | 10        |
| 44 | Safety and Antitumor Activity of Repeated ASP3026 Administration in Japanese Patients with Solid Tumors: A Phase I Study. <i>Drugs in R and D</i> , 2021, 21, 65-78.  | 1.1 | 1         |
| 45 | Rationale and Design for a Multicenter, Phase II Study of Durvalumab Plus Concurrent Radiation Therapy in Locally Advanced Non-Small Cell Lung Cancer: The DOLPHIN Study (WJOG11619L). <i>Cancer Management and Research</i> , 2021, Volume 13, 9167-9173.  | 0.9 | 5         |
| 46 | Combination therapy with PD-1 or PD-L1 inhibitors for cancer. <i>International Journal of Clinical Oncology</i> , 2020, 25, 818-830.  | 1.0 | 86        |
| 47 | Final progression-free survival results from the J-ALEX study of alectinib versus crizotinib in ALK-positive non-small-cell lung cancer. <i>Lung Cancer</i> , 2020, 139, 195-199.   | 0.9 | 100       |
| 48 | Randomized Phase III Study of Continuation Maintenance Bevacizumab With or Without Pemetrexed in Advanced Nonsquamous Nonâ€“Small-Cell Lung Cancer: COMPASS (WJOG5610L). <i>Journal of Clinical Oncology</i> , 2020, 38, 793-803.                           | 0.8 | 28        |
| 49 | Ramucirumab or placebo plus erlotinib in <i>EGFR</i> -mutated, metastatic nonâ€“smallâ€“cell lung cancer: East Asian subset of RELAY. <i>Cancer Science</i> , 2020, 111, 4510-4525.   | 1.7 | 17        |
| 50 | Severe Immune-Related Hepatitis Treated With Plasma Exchange. <i>Journal of Thoracic Oncology</i> , 2020, 15, e39-e42.  | 0.5 | 3         |
| 51 | Simultaneous targeting of MET overexpression in EGFR mutation-positive non-small cell lung cancer can increase the benefit of EGFR-TKI therapy?. <i>Translational Lung Cancer Research</i> , 2020, 9, 1617-1622.  | 1.3 | 2         |
| 52 | Patient-reported outcomes in RELAY, a phase 3 trial of ramucirumab plus erlotinib versus placebo plus erlotinib in untreated <i>EGFR</i> -mutated metastatic non-small-cell lung cancer. <i>Current Medical Research and Opinion</i> , 2020, 36, 1667-1675. | 0.9 | 11        |
| 53 | Efficacy of Combination Chemotherapy Using a Novel Oral Chemotherapeutic Agent, FTD/TPI, with Ramucirumab Murine Version DC101 in a Mouse Syngeneic Cancer Transplantation Model. <i>Journal of Clinical Medicine</i> , 2020, 9, 4050.                      | 1.0 | 2         |
| 54 | Efficacy and safety of pembrolizumab for the treatment of advanced biliary cancer: Results from the <sc>KEYNOTE</sc>â€“158 and <sc>KEYNOTE</sc>â€“028 studies. <i>International Journal of Cancer</i> , 2020, 147, 2190-2198.                               | 2.3 | 288       |

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 55 | Comparison of Carboplatin Plus Pemetrexed Followed by Maintenance Pemetrexed With Docetaxel Monotherapy in Elderly Patients With Advanced Nonsquamous Non-small Cell Lung Cancer. <i>JAMA Oncology</i> , 2020, 6, e196828.   | 3.4 | 48        |
| 56 | Differential significance of molecular subtypes which were classified into EGFR exon 19 deletion on the first line afatinib monotherapy. <i>BMC Cancer</i> , 2020, 20, 103.  | 1.1 | 14        |
| 57 | Impact of EGFR-TKI Treatment on the Tumor Immune Microenvironment in EGFR Mutation-Positive Non-small Cell Lung Cancer. <i>Clinical Cancer Research</i> , 2020, 26, 2037-2046.   | 3.2 | 142       |
| 58 | Durvalumab With or Without Tremelimumab vs Standard Chemotherapy in First-line Treatment of Metastatic Non-small Cell Lung Cancer. <i>JAMA Oncology</i> , 2020, 6, 661.  | 3.4 | 446       |
| 59 | Safety and efficacy of first-line dacomitinib in Japanese patients with advanced non-small cell lung cancer. <i>Cancer Science</i> , 2020, 111, 1724-1738.   | 1.7 | 20        |
| 60 | NivoCUP: An open-label phase II study on the efficacy of nivolumab in cancer of unknown primary.. <i>Journal of Clinical Oncology</i> , 2020, 38, 106-106.   | 0.8 | 11        |
| 61 | Trastuzumab deruxtecan (T-DXd; DS-8201) in patients with HER2-mutated metastatic non-small cell lung cancer (NSCLC): Interim results of DESTINY-Lung01.. <i>Journal of Clinical Oncology</i> , 2020, 38, 9504-9504.  | 0.8 | 91        |
| 62 | RELAY+: Exploratory study of ramucirumab plus gefitinib in untreated patients (pts) with epidermal growth factor receptor (EGFR)-mutated metastatic non-small cell lung cancer (NSCLC).. <i>Journal of Clinical Oncology</i> , 2020, 38, 9564-9564.                              | 0.8 | 2         |
| 63 | NGSCUP: Phase II trial of site-specific treatment based on gene expression and mutation profiling by next generation sequencing (NGS) for patients (pts) with cancer of unknown primary site (CUP).. <i>Journal of Clinical Oncology</i> , 2020, 38, e15577-e15577.              | 0.8 | 1         |
| 64 | RELAY study of erlotinib (ERL) + ramucirumab (RAM) or placebo (PL) in EGFR-mutated metastatic non-small cell lung cancer (NSCLC): Biomarker analysis using circulating tumor DNA (ctDNA) in Japanese patients (pts).. <i>Journal of Clinical Oncology</i> , 2020, 38, 9527-9527. | 0.8 | 1         |
| 65 | Effects of dose modifications on the safety and efficacy of dacomitinib for EGFR mutation-positive non-small-cell lung cancer. <i>Future Oncology</i> , 2019, 15, 2795-2805.   | 1.1 | 27        |
| 66 | Phase I safety and pharmacokinetics study of rovalpituzumab tesirine in Japanese patients with advanced, recurrent small cell lung cancer. <i>Lung Cancer</i> , 2019, 135, 145-150.  | 0.9 | 18        |
| 67 | Three-year follow-up results from phase II studies of nivolumab in Japanese patients with previously treated advanced non-small cell lung cancer: Pooled analysis of ONO4538-05 and ONO4538-06 studies. <i>Cancer Medicine</i> , 2019, 8, 5183-5193.                             |     | 13        |
| 68 | Ramucirumab plus erlotinib in patients with untreated, EGFR-mutated, advanced non-small-cell lung cancer (RELAY): a randomised, double-blind, placebo-controlled, phase 3 trial. <i>Lancet Oncology</i> , The, 2019, 20, 1655-1669.  | 5.1 | 418       |
| 69 | A comparative study of curated contents by knowledge-based curation system in cancer clinical sequencing. <i>Scientific Reports</i> , 2019, 9, 11340.  | 1.6 | 12        |
| 70 | Clinical and immune profiling for cancer of unknown primary site. , 2019, 7, 251.  |     | 26        |
| 71 | Aberrant HER3 ligand heregulin-expressing head and neck squamous cell carcinoma is resistant to anti-EGFR antibody cetuximab, but not second-generation EGFR-TKI. <i>Oncogenesis</i> , 2019, 8, 54.  | 2.1 | 12        |
| 72 | Clinical Efficacy and Safety of Nivolumab: Results of a Multicenter, Open-label, Single-arm, Japanese Phase II study in Malignant Pleural Mesothelioma (MERIT). <i>Clinical Cancer Research</i> , 2019, 25, 5485-5492.   | 3.2 | 191       |

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 73 | New Era for Next-Generation Sequencing in Japan. <i>Cancers</i> , 2019, 11, 742.  | 1.7 | 22        |
| 74 | Management of common adverse events related to first-line dacomitinib use in <i>EGFR</i> mutation-positive non-small-cell lung cancer: a pooled safety analysis. <i>Future Oncology</i> , 2019, 15, 1481-1491.  | 1.1 | 11        |
| 75 | <scp>KEYNOTE</scp>â€25: Phase 1b study of pembrolizumab in Japanese patients with previously treated programmed death ligand 1â€positive advanced nonâ€smallâ€cell lung cancer. <i>Cancer Science</i> , 2019, 110, 1012-1020.   | 1.7 | 40        |
| 76 | Japanese subgroup analysis of a phase III study of S-1 versus docetaxel in non-small cell lung cancer patients after platinum-based treatment: EAST-LC. <i>International Journal of Clinical Oncology</i> , 2019, 24, 485-493.  | 1.0 | 4         |
| 77 | Sequencing of therapy following first-line afatinib in patients with EGFR mutation-positive non-small cell lung cancer. <i>Lung Cancer</i> , 2019, 132, 126-131.  | 0.9 | 26        |
| 78 | Clinical significance of monitoring EGFR mutation in plasma using multiplexed digital PCR in EGFR mutated patients treated with afatinib (West Japan Oncology Group 8114LTR study). <i>Lung Cancer</i> , 2019, 131, 128-133.  | 0.9 | 18        |
| 79 | Heregulin expression and its clinical implication for patients with EGFR-mutant non-small cell lung cancer treated with EGFR-tyrosine kinase inhibitors. <i>Scientific Reports</i> , 2019, 9, 19501.  | 1.6 | 12        |
| 80 | Osimertinib versus standard-of-care EGFR-TKI as first-line treatment for EGFRm advanced NSCLC: FLAURA Japanese subset. <i>Japanese Journal of Clinical Oncology</i> , 2019, 49, 29-36.  | 0.6 | 101       |
| 81 | First-line onartuzumab plus erlotinib treatment for patients with MET-positive and EGFR mutation-positive non-small-cell lung cancer. <i>Cancer Treatment and Research Communications</i> , 2019, 18, 100113.   | 0.7 | 9         |
| 82 | First- and Second-Generation EGFR-TKIs Are All Replaced to Osimertinib in Chemo-Naive EGFR Mutation-Positive Non-Small Cell Lung Cancer?. <i>International Journal of Molecular Sciences</i> , 2019, 20, 146.   | 1.8 | 118       |
| 83 | Mutational activation of the epidermal growth factor receptor downâ€regulates major histocompatibility complex class I expression via the extracellular signalâ€regulated kinase in nonâ€small cell lung cancer. <i>Cancer Science</i> , 2019, 110, 52-60.   | 1.7 | 31        |
| 84 | U3-1402 sensitizes HER3-expressing tumors to PD-1 blockade by immune activation. <i>Journal of Clinical Investigation</i> , 2019, 130, 374-388.   | 3.9 | 43        |
| 85 | Pembrolizumab (pembro) for advanced biliary adenocarcinoma: Results from the KEYNOTE-028 (KN028) and KEYNOTE-158 (KN158) basket studies.. <i>Journal of Clinical Oncology</i> , 2019, 37, 4079-4079.  | 0.8 | 94        |
| 86 | RELAY: A multinational, double-blind, randomized Phase 3 study of erlotinib (ERL) in combination with ramucirumab (RAM) or placebo (PL) in previously untreated patients with epidermal growth factor receptor mutation-positive (EGFRm) metastatic non-small cell lung cancer (NSCLC).. <i>Journal of Clinical Oncology</i> , 2019, 37, 9000-9000.                                   | 0.8 | 23        |
| 87 | A randomized phase III study of continuous maintenance bevacizumab with or without pemetrexed after induction therapy with carboplatin (Car), pemetrexed (Pem), and bevacizumab (Bev) for advanced non-squamous non-small cell lung cancer (nSQ-NSCLC) without sensitizing EGFR mutations: The COMPASS study (WJOG5610L).. <i>Journal of Clinical Oncology</i> , 2019, 37, 9003-9003. | 0.8 | 3         |
| 88 | Blood tumor mutational burden (bTMB) and tumor PD-L1 as predictive biomarkers of survival in MYSTIC: First-line durvalumab (D) Â± tremelimumab (T) versus chemotherapy (CT) in metastatic (m) NSCLC.. <i>Journal of Clinical Oncology</i> , 2019, 37, 9016-9016.  | 0.8 | 20        |
| 89 | Randomized phase III study comparing carboplatin plus pemetrexed followed by pemetrexed versus docetaxel in elderly patients with advanced non-squamous non-small-cell lung cancer (JCOG1210/WJOG7813L).. <i>Journal of Clinical Oncology</i> , 2019, 37, 9031-9031.  | 0.8 | 8         |
| 90 | Final PFS analysis and safety data from the phase III J-ALEX study of alectinib (ALC) vs. crizotinib (CRZ) in ALK-inhibitor naâ€ve ALK-positive non-small cell lung cancer (ALK+ NSCLC).. <i>Journal of Clinical Oncology</i> , 2019, 37, 9092-9092.   | 0.8 | 14        |

| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 91  | A multicenter, open label, randomized phase III study of atezolizumab with platinum-pemetrexed and with or without bevacizumab for patients with advanced nonsquamous non-small cell lung cancer (WJOG11218L APPLE Study).. Journal of Clinical Oncology, 2019, 37, TPS9125-TPS9125.   | 0.8 | 3         |
| 92  | The impact of sequential therapy of crizotinib followed by alectinib: Real-world data analysis of 840 ALK-inhibitor naïve patients with NSCLC harboring ALK-rearrangement (WJOG9516L).. Journal of Clinical Oncology, 2019, 37, 9038-9038.   | 0.8 | 0         |
| 93  | Phase I study on preliminary safety and efficacy of rovalpituzumab tesirine in Japanese patients (pts) with advanced, recurrent small cell lung cancer (SCLC).. Journal of Clinical Oncology, 2019, 37, 8557-8557.   | 0.8 | 2         |
| 94  | Randomized phase 3 study of maintenance therapy with S-1 plus best supportive care (BSC) versus BSC alone after induction therapy with carboplatin plus S-1 for advanced or relapsed squamous cell lung carcinoma (WJOG7512L).. Journal of Clinical Oncology, 2019, 37, e20531-e20531. | 0.8 | 0         |
| 95  | Phase 3 study of ceritinib vs chemotherapy in ALK-rearranged NSCLC patients previously treated with chemotherapy and crizotinib (ASCEND-5): Japanese subset. Japanese Journal of Clinical Oncology, 2018, 48, 367-375.   | 0.6 | 26        |
| 96  | Real world treatment and outcomes in EGFR mutation-positive non-small cell lung cancer: Long-term follow-up of a large patient cohort. Lung Cancer, 2018, 117, 14-19.  | 0.9 | 63        |
| 97  | Randomized, Double-Blind Phase Ib/III Study of Erlotinib With Ramucirumab or Placebo in Previously Untreated EGFR -Mutant Metastatic Non-“Small-Cell Lung Cancer (RELAY): Phase Ib Results. Clinical Lung Cancer, 2018, 19, 213-220.e4.  | 1.1 | 13        |
| 98  | Analysis of central nervous system efficacy in the J-ALEX study of alectinib versus crizotinib in ALK-positive non-small-cell lung cancer. Lung Cancer, 2018, 121, 37-40.  | 0.9 | 62        |
| 99  | Sterilized talc pleurodesis for malignant pleural effusions: a Phase II study for investigational new drug application in Japan. Japanese Journal of Clinical Oncology, 2018, 48, 376-381.   | 0.6 | 3         |
| 100 | Improvement in Overall Survival in a Randomized Study That Compared Dacomitinib With Gefitinib in Patients With Advanced Non-“Small-Cell Lung Cancer and <i>EGFR</i>-Activating Mutations. Journal of Clinical Oncology, 2018, 36, 2244-2250.  | 0.8 | 361       |
| 101 | CNS Response to Osimertinib Versus Standard Epidermal Growth Factor Receptor Tyrosine Kinase Inhibitors in Patients With Untreated <i>EGFR</i>-Mutated Advanced Non-“Small-Cell Lung Cancer. Journal of Clinical Oncology, 2018, 36, 3290-3297.  | 0.8 | 515       |
| 102 | Induction Chemoradiotherapy (50 Gy), Followed by Resection, for Stage IIIA-N2 Non-Small Cell Lung Cancer. Annals of Thoracic Surgery, 2018, 106, 1018-1024.  | 0.7 | 8         |
| 103 | A randomised phase II trial of S-1 plus cisplatin versus vinorelbine plus cisplatin with concurrent thoracic radiotherapy for unresectable, locally advanced non-small cell lung cancer: WJOG5008L. British Journal of Cancer, 2018, 119, 675-682.                                     | 2.9 | 32        |
| 104 | Prognostic value of Lung Cancer Subscale in older patients with advanced non-small cell lung cancer: An integrated analysis of JCOG0207 and JCOG0803/WJOG4307L (JCOG1414A). Journal of Geriatric Oncology, 2018, 9, 583-588.   | 0.5 | 4         |
| 105 | <sc>ASP</sc>8273 tolerability and antitumor activity in tyrosine kinase inhibitor-naïve Japanese patients with <i>EGFR</i> mutation-positive non-small-cell lung cancer. Cancer Science, 2018, 109, 2532-2538.   | 1.7 | 10        |
| 106 | An open-label feasibility study of nintedanib combined with docetaxel in Japanese patients with locally advanced or metastatic lung adenocarcinoma after failure of first-line chemotherapy. Cancer Chemotherapy and Pharmacology, 2018, 82, 685-694.                                  | 1.1 | 3         |
| 107 | A randomized phase II trial of trastuzumab plus capecitabine versus lapatinib plus capecitabine in patients with HER2-positive metastatic breast cancer previously treated with trastuzumab and taxanes: WJOG6110B/ELTOP. Breast, 2018, 40, 67-75.                                     | 0.9 | 34        |
| 108 | Clinical activity of <sc>ASP</sc>8273 in Asian patients with non-small-cell lung cancer with <i>EGFR</i> activating and T790M mutations. Cancer Science, 2018, 109, 2852-2862.   | 1.7 | 15        |

| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 109 | Dacomitinib (daco) versus gefitinib (gef) for first-line treatment of advanced NSCLC (ARCHER 1050): Final overall survival (OS) analysis.. Journal of Clinical Oncology, 2018, 36, 9004-9004.   | 0.8 | 9         |
| 110 | Clinical characteristics of non-small cell lung cancer harboring mutations in exon 20 of <i>EGFR</i> or <i>HER2</i> . Oncotarget, 2018, 9, 21132-21140.   | 0.8 | 24        |
| 111 | Circulating heregulin level is associated with the efficacy of patritumab combined with erlotinib in patients with non-small cell lung cancer. Lung Cancer, 2017, 105, 1-6.   | 0.9 | 21        |
| 112 | Alectinib versus crizotinib in patients with ALK -positive non-small-cell lung cancer (J-ALEX): an open-label, randomised phase 3 trial. Lancet, The, 2017, 390, 29-39.   | 6.3 | 753       |
| 113 | Phase 1 study of new formulation of patritumab (U3-1287) Process 2, a fully human anti-HER3 monoclonal antibody in combination with erlotinib in Japanese patients with advanced non-small cell lung cancer. Cancer Chemotherapy and Pharmacology, 2017, 79, 489-495.   | 1.1 | 25        |
| 114 | A Randomized Phase II Study Comparing Nivolumab With Carboplatin-Pemetrexed for Patients With EGFR Mutation-Positive Nonsquamous Non-Small-Cell Lung Cancer Who Acquire Resistance to Tyrosine Kinase Inhibitors Not Due to a Secondary T790M Mutation: Rationale and Protocol Design for the WJOG8515L Study. Clinical Lung Cancer, 2017, 18, 719-723. | 1.1 | 13        |
| 115 | Gefitinib or Erlotinib vs Chemotherapy for EGFR Mutation-Positive Lung Cancer: Individual Patient Data Meta-Analysis of Overall Survival. Journal of the National Cancer Institute, 2017, 109, .  | 3.0 | 196       |
| 116 | Ceritinib in patients with advanced, crizotinib-treated, anaplastic lymphoma kinase-rearranged NSCLC: Japanese subset. Japanese Journal of Clinical Oncology, 2017, 47, 618-624.  | 0.6 | 14        |
| 117 | Dacomitinib versus gefitinib as first-line treatment for patients with EGFR-mutation-positive non-small-cell lung cancer (ARCHER 1050): a randomised, open-label, phase 3 trial. Lancet Oncology, The, 2017, 18, 1454-1466.   | 5.1 | 877       |
| 118 | Treatment Rationale and Study Design for the RELAY Study: A Multicenter, Randomized, Double-Blind Study of Erlotinib With Ramucirumab or Placebo in Patients With Epidermal Growth Factor Receptor Mutation-Positive Metastatic Non-Small-Cell Lung Cancer. Clinical Lung Cancer, 2017, 18, 96-99.  | 1.1 | 10        |
| 119 | Three-Year Follow-Up of an Alectinib Phase I/II Study in ALK-Positive Non-Small-Cell Lung Cancer: AF-001JP. Journal of Clinical Oncology, 2017, 35, 1515-1521.  | 0.8 | 63        |
| 120 | Bevacizumab beyond disease progression after first-line treatment with bevacizumab plus chemotherapy in advanced nonsquamous non-small cell lung cancer (WJOG50302) trial. Cancer, 2016, 122, 1050-1059.  | 2.0 | 35        |
| 121 | <i>FGFR</i> gene alterations in lung squamous cell carcinoma are potential targets for the multikinase inhibitor nintedanib. Cancer Science, 2016, 107, 1667-1676.  | 1.7 | 31        |
| 122 | Randomized Phase III Study Comparing Gefitinib With Erlotinib in Patients With Previously Treated Advanced Lung Adenocarcinoma: WJOG 5108L. Journal of Clinical Oncology, 2016, 34, 3248-3257.  | 0.8 | 130       |
| 123 | A randomized, double-blind, phase II study of ramucirumab plus docetaxel vs placebo plus docetaxel in Japanese patients with stage IV non-small cell lung cancer after disease progression on platinum-based therapy. Lung Cancer, 2016, 99, 186-193.   | 0.9 | 88        |
| 124 | Characteristics and overall survival of EGFR mutation-positive non-small cell lung cancer treated with EGFR tyrosine kinase inhibitors: a retrospective analysis for 1660 Japanese patients. Japanese Journal of Clinical Oncology, 2016, 46, 462-467.  | 0.6 | 54        |
| 125 | A phase II study of pemetrexed plus carboplatin followed by maintenance pemetrexed as first-line chemotherapy for elderly patients with advanced non-squamous non-small cell lung cancer. Medical Oncology, 2016, 33, 2.  | 1.2 | 18        |
| 126 | Japanese Society of Clinical Oncology clinical practice guidelines 2010 for antiemesis in oncology: executive summary. International Journal of Clinical Oncology, 2016, 21, 1-12.  | 1.0 | 58        |



| #   | ARTICLE   | IF   | CITATIONS |
|-----|---|------|-----------|
| 127 | Tolerability of Nintedanib (BIBF 1120) in Combination with Docetaxel: A Phase 1 Study in Japanese Patients with Previously Treated Non-Small-Cell Lung Cancer. <i>Journal of Thoracic Oncology</i> , 2015, 10, 346-352.   | 0.5  | 35        |
| 128 | Phase I study of the HER3-targeted antibody patritumab (U3-1287) combined with erlotinib in Japanese patients with non-small cell lung cancer. <i>Lung Cancer</i> , 2015, 88, 275-281.  | 0.9  | 36        |
| 129 | Randomized Phase III Trial Comparing Weekly Docetaxel Plus Cisplatin Versus Docetaxel Monotherapy Every 3 Weeks in Elderly Patients With Advanced Non-Small-Cell Lung Cancer: The Intergroup Trial JCOG0803/WJOG4307L. <i>Journal of Clinical Oncology</i> , 2015, 33, 575-581. | 0.8  | 109       |
| 130 | Multiple regulatory mechanisms of hepatocyte growth factor expression in malignant cells with a short poly(dA) sequence in the HGF gene promoter. <i>Oncology Letters</i> , 2015, 9, 405-410.   | 0.8  | 5         |
| 131 | Gefitinib plus chemotherapy versus placebo plus chemotherapy in EGFR-mutation-positive non-small-cell lung cancer after progression on first-line gefitinib (IMPRESS): a phase 3 randomised trial. <i>Lancet Oncology</i> , The, 2015, 16, 990-998.                             | 5.1  | 353       |
| 132 | Pooled safety analysis of EGFR-TKI treatment for EGFR mutation-positive non-small cell lung cancer. <i>Lung Cancer</i> , 2015, 88, 74-79.   | 0.9  | 157       |
| 133 | A case of severe cerebral embolism after chemotherapy for HER2-positive gastric cancer. <i>BMC Research Notes</i> , 2015, 8, 100.   | 0.6  | 4         |
| 134 | A phase I/IIb study of trametinib (GSK1120212) alone and in combination with gemcitabine in Japanese patients with advanced solid tumors. <i>Investigational New Drugs</i> , 2015, 33, 1058-1067.   | 1.2  | 13        |
| 135 | The pan-HER family tyrosine kinase inhibitor afatinib overcomes HER3 ligand heregulin-mediated resistance to EGFR inhibitors in non-small cell lung cancer. <i>Oncotarget</i> , 2015, 6, 33602-33611.   | 0.8  | 53        |
| 136 | The anti-HER3 antibody patritumab abrogates cetuximab resistance mediated by heregulin in colorectal cancer cells. <i>Oncotarget</i> , 2014, 5, 11847-11856.  | 0.8  | 61        |
| 137 | Targeting MET Amplification as a New Oncogenic Driver. <i>Cancers</i> , 2014, 6, 1540-1552.   | 1.7  | 96        |
| 138 | First-Line Crizotinib versus Chemotherapy in ALK-Positive Lung Cancer. <i>New England Journal of Medicine</i> , 2014, 371, 2167-2177.   | 13.9 | 2,808     |
| 139 | Current status and future perspectives of cooperative study groups for lung cancer in Japan. <i>Respiratory Investigation</i> , 2014, 52, 339-347.  | 0.9  | 3         |
| 140 | Final safety and efficacy of erlotinib in the phase 4 POLARSTAR surveillance study of 10,708 Japanese patients with non-small cell lung cancer. <i>Cancer Science</i> , 2014, 105, 1584-1590.   | 1.7  | 67        |
| 141 | Phase III Study Comparing Amrubicin Plus Cisplatin With Irinotecan Plus Cisplatin in the Treatment of Extensive-Disease Small-Cell Lung Cancer: JCOG 0509. <i>Journal of Clinical Oncology</i> , 2014, 32, 1262-1268.   | 0.8  | 66        |
| 142 | Erlotinib alone or with bevacizumab as first-line therapy in patients with advanced non-squamous non-small-cell lung cancer harbouring EGFR mutations (JO25567): an open-label, randomised, multicentre, phase 2 study. <i>Lancet Oncology</i> , The, 2014, 15, 1236-1244.      | 5.1  | 678       |
| 143 | The expression level of HER3 ligand heregulin mRNA as a predictive biomarker for anti-HER3 antibody patritumab combined with erlotinib in non-small cell lung cancer. <i>Journal of Clinical Oncology</i> , 2014, 32, e19082-e19082.  | 0.8  | 2         |
| 144 | Multiplex genomic profiling of non-small cell lung cancers from the LETS phase III trial of first-line S-1/carboplatin versus paclitaxel/carboplatin: results of a West Japan Oncology Group study. <i>Oncotarget</i> , 2014, 5, 2293-2304.                                     | 0.8  | 32        |

| #   | ARTICLE  | IF   | CITATIONS |
|-----|--|------|-----------|
| 145 | Pemetrexed and carboplatin followed by pemetrexed maintenance therapy in chemo-naïve patients with advanced nonsquamous non-small-cell lung cancer. <i>Investigational New Drugs</i> , 2013, 31, 1275-1282.  | 1.2  | 38        |
| 146 | CH5424802 (RO5424802) for patients with ALK-rearranged advanced non-small-cell lung cancer (AF-001JP study): a single-arm, open-label, phase 1&2 study. <i>Lancet Oncology</i> , The, 2013, 14, 590-598.   | 5.1  | 555       |
| 147 | Crizotinib versus Chemotherapy in Advanced ALK-Positive Lung Cancer. <i>New England Journal of Medicine</i> , 2013, 368, 2385-2394.  | 13.9 | 3,181     |
| 148 | Interstitial Lung Disease Associated with Gefitinib in Japanese Patients with EGFR-mutated Non-small-cell Lung Cancer: Combined Analysis of Two Phase III Trials (NEJ 002 and WJTOG 3405). <i>Japanese Journal of Clinical Oncology</i> , 2013, 43, 664-668.           | 0.6  | 38        |
| 149 | MET amplification as a potential therapeutic target in gastric cancer. <i>Oncotarget</i> , 2013, 4, 9-17.  | 0.8  | 82        |
| 150 | Phase I dose finding study of AZD8931, an inhibitor of EGFR, HER2, and HER3 signaling, alone or in combination with paclitaxel in Japanese patients.. <i>Journal of Clinical Oncology</i> , 2013, 31, e13501-e13501.   | 0.8  | 0         |
| 151 | Antitumor Action of the MET Tyrosine Kinase Inhibitor Crizotinib (PF-02341066) in Gastric Cancer Positive for MET Amplification. <i>Molecular Cancer Therapeutics</i> , 2012, 11, 1557-1564.   | 1.9  | 75        |
| 152 | Postmarketing Surveillance Study of Erlotinib in Japanese Patients With Non-Small-Cell Lung Cancer (NSCLC): An Interim Analysis of 3488 Patients (POLARSTAR). <i>Journal of Thoracic Oncology</i> , 2012, 7, 1296-1303.  | 0.5  | 73        |
| 153 | Perspectives in drug development for cancer therapy in Asia. <i>Chinese Clinical Oncology</i> , 2012, 1, 17.   | 0.4  | 1         |
| 154 | Activation of ERBB2 Signaling Causes Resistance to the EGFR-Directed Therapeutic Antibody Cetuximab. <i>Science Translational Medicine</i> , 2011, 3, 99ra86.  | 5.8  | 543       |
| 155 | MET Tyrosine Kinase Inhibitor Crizotinib (PF-02341066) Shows Differential Antitumor Effects in Non-small Cell Lung Cancer According to MET Alterations. <i>Journal of Thoracic Oncology</i> , 2011, 6, 1624-1631.  | 0.5  | 122       |
| 156 | Phase I Safety, Pharmacokinetic, and Biomarker Study of BIBF 1120, an Oral Triple Tyrosine Kinase Inhibitor in Patients with Advanced Solid Tumors. <i>Molecular Cancer Therapeutics</i> , 2010, 9, 2825-2833.   | 1.9  | 91        |
| 157 | Phase III Trial Comparing Oral S-1 Plus Carboplatin With Paclitaxel Plus Carboplatin in Chemotherapy-Naïve Patients With Advanced Non-Small-Cell Lung Cancer: Results of a West Japan Oncology Group Study. <i>Journal of Clinical Oncology</i> , 2010, 28, 5240-5246. | 0.8  | 161       |
| 158 | Gefitinib versus cisplatin plus docetaxel in patients with non-small-cell lung cancer harbouring mutations of the epidermal growth factor receptor (WJTOG3405): an open label, randomised phase 3 trial. <i>Lancet Oncology</i> , The, 2010, 11, 121-128.              | 5.1  | 3,794     |
| 159 | Interstitial Lung Disease in Japanese Patients with Lung Cancer. <i>American Journal of Respiratory and Critical Care Medicine</i> , 2008, 177, 1348-1357.   | 2.5  | 428       |
| 160 | Efficacy and Safety of Pemetrexed in Combination with Cisplatin for Malignant Pleural Mesothelioma: A Phase I/II Study in Japanese Patients. <i>Japanese Journal of Clinical Oncology</i> , 2008, 38, 339-346.   | 0.6  | 22        |
| 161 | Phase I Dose-escalation and Pharmacokinetic Trial of Lapatinib (GW572016), a Selective Oral Dual Inhibitor of ErbB-1 and -2 Tyrosine Kinases, in Japanese Patients with Solid Tumors. <i>Japanese Journal of Clinical Oncology</i> , 2008, 39, 116-123.                | 0.6  | 18        |
| 162 | Efficacy and Safety of Erlotinib Monotherapy for Japanese Patients with Advanced Non-small Cell Lung Cancer: A Phase II Study. <i>Journal of Thoracic Oncology</i> , 2008, 3, 1439-1445.   | 0.5  | 73        |

| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 163 | Clinical development of EGFR-tyrosine kinase inhibitors in Japan. <i>Cancer Chemotherapy and Pharmacology</i> , 2006, 58, 33-37.  | 1.1 | 1         |
| 164 | Multi-Institutional Randomized Phase II Trial of Gefitinib for Previously Treated Patients With Advanced Nonâ€“Small-Cell Lung Cancer. <i>Journal of Clinical Oncology</i> , 2003, 21, 2237-2246. | 0.8 | 2,822     |
| 165 | Phase I and Pharmacologic Study of Docetaxel and Irinotecan in Advanced Nonâ€“Small-Cell Lung Cancer. <i>Journal of Clinical Oncology</i> , 2000, 18, 2996-3003.                                  | 0.8 | 38        |
| 166 | Two cases of portal-systemic shunts demonstrated by portal venography performed by operative cannulation of ileocolic vein.. <i>Japanese Journal of Medicine</i> , 1983, 22, 31-36.               | 0.1 | 2         |