

# Raja Mokdad-Gargouri

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7812332/publications.pdf>

Version: 2024-02-01

61  
papers

928  
citations

394421

19  
h-index

552781

26  
g-index

62  
all docs

62  
docs citations

62  
times ranked

1568  
citing authors

#	ARTICLE	IF	CITATIONS
1	Inactivation of RASSF1A, RAR $\beta$ 2 and DAP-kinase by promoter methylation correlates with lymph node metastasis in nasopharyngeal carcinoma. <i>Cancer Biology and Therapy</i> , 2009, 8, 444-451.	3.4	48
2	Genotyping of Tunisian hepatitis B virus isolates based on the sequencing of preS2 and S regions. <i>Microbes and Infection</i> , 2000, 2, 607-612.	1.9	39
3	BCL2L12 is a Novel Biomarker for the Prediction of Short-Term Relapse in Nasopharyngeal Carcinoma. <i>Molecular Medicine</i> , 2011, 17, 163-171.	4.4	39
4	PIK3CA amplification is predictive of poor prognosis in Tunisian patients with nasopharyngeal carcinoma. <i>Cancer Science</i> , 2009, 100, 2034-2039.	3.9	37
5	Aberrant methylation of RASSF1A is associated with poor survival in Tunisian breast cancer patients. <i>Journal of Cancer Research and Clinical Oncology</i> , 2010, 136, 203-210.	2.5	36
6	Various 30 and 69bp deletion variants of the Epstein-Barr virus LMP1 may arise by homologous recombination in nasopharyngeal carcinoma of Tunisian patients. <i>Virus Research</i> , 2006, 115, 24-30.	2.2	32
7	Hypermethylation of tumor-related genes in tunisian patients with gastric carcinoma: Clinical and biological significance. <i>Journal of Surgical Oncology</i> , 2011, 103, 687-694.	1.7	32
8	Overexpression of miR-10b in colorectal cancer patients: Correlation with <i>Twist-1</i> and E-cadherin expression. <i>Tumor Biology</i> , 2017, 39, 101042831769591.	1.8	32
9	Epigenetic Alteration of the Wnt Inhibitory Factor-1 Promoter Is Common and Occurs in Advanced Stage of Tunisian Nasopharyngeal Carcinoma. <i>Cancer Investigation</i> , 2010, 28, 896-903.	1.3	30
10	Human p53 induces cell death and downregulates thioredoxin expression in <i>Saccharomyces cerevisiae</i> . <i>FEMS Yeast Research</i> , 2008, 8, 1254-1262.	2.3	28
11	Quantitative expression analysis and prognostic significance of the BCL2-associated Xgene in nasopharyngeal carcinoma: a retrospective cohort study. <i>BMC Cancer</i> , 2013, 13, 293.	2.6	24
12	Translational control of human p53 expression in yeast mediated by 5'-UTR-ORF structural interaction. <i>Nucleic Acids Research</i> , 2001, 29, 1222-1227.	14.5	23
13	Clinical Significance of Epigenetic Inactivation of hMLH1 and BRCA1 in Tunisian Patients with Invasive Breast Carcinoma. <i>Journal of Biomedicine and Biotechnology</i> , 2009, 2009, 1-7.	3.0	23
14	Loss of WIF-1 and Wnt5a expression is related to aggressiveness of sporadic breast cancer in Tunisian patients. <i>Tumor Biology</i> , 2013, 34, 1625-1633.	1.8	23
15	Yeasts as a Tool for Heterologous Gene Expression. <i>Methods in Molecular Biology</i> , 2012, 824, 359-370.	0.9	22
16	Expression of COX-2 and E-cadherin in Tunisian patients with colorectal adenocarcinoma. <i>Acta Histochemica</i> , 2012, 114, 577-581.	1.8	22
17	Over-expression of miR-10b in NPC patients: correlation with LMP1 and Twist1. <i>Tumor Biology</i> , 2015, 36, 3807-3814.	1.8	20
18	Expression and Mutation Pattern of $\beta$ -Catenin and Adenomatous Polyposis Coli in Colorectal Cancer Patients. <i>Archives of Medical Research</i> , 2015, 46, 54-62.	3.3	20

#	ARTICLE	IF	CITATIONS
19	Serine protease inhibitors and human wellbeing interplay: new insights for old friends. <i>PeerJ</i> , 2019, 7, e7224.	2.0	20
20	Hypermethylation of RAR $\beta$ 2 correlates with high COX-2 expression and poor prognosis in patients with colorectal carcinoma. <i>Tumor Biology</i> , 2010, 31, 503-511.	1.8	19
21	Characteristics of epstein barr virus variants associated with gastric carcinoma in Southern Tunisia. <i>Virology Journal</i> , 2011, 8, 500.	3.4	18
22	Extraction and purification of hepatitis B virus-like M particles from a recombinant <i>Saccharomyces cerevisiae</i> strain using alumina powder. <i>Journal of Virological Methods</i> , 2013, 187, 132-137.	2.1	18
23	Aberrant methylation of hMLH1 and p16INK4a in Tunisian patients with sporadic colorectal adenocarcinoma. <i>Bioscience Reports</i> , 2011, 31, 257-264.	2.4	17
24	CpG methylation of ubiquitin carboxyl-terminal hydrolase 1 (UCHL1) and P53 mutation pattern in sporadic colorectal cancer. <i>Tumor Biology</i> , 2016, 37, 1707-1714.	1.8	17
25	Association of FOXA1 and EMT markers (Twist1 and E-cadherin) in breast cancer. <i>Molecular Biology Reports</i> , 2019, 46, 3247-3255.	2.3	17
26	Cloning and characterization of cDNA probes for the analysis of metallothionein gene expression in the Mediterranean bivalves: <i>Ruditapes decussatus</i> and <i>Cerastoderma glaucum</i> . <i>Molecular Biology Reports</i> , 2009, 36, 1007-1014.	2.3	14
27	Expression of HBsAg and preS2-S protein in different yeast based system: A comparative analysis. <i>Protein Expression and Purification</i> , 2009, 66, 131-137.	1.3	14
28	Expression of APC, $\beta$ -catenin and E-cadherin in Tunisian patients with gastric adenocarcinoma: clinical significance. <i>Tumor Biology</i> , 2014, 35, 1775-1783.	1.8	14
29	Downregulation of WIF-1 and Wnt5a in patients with colorectal carcinoma: clinical significance. <i>Tumor Biology</i> , 2014, 35, 7975-7982.	1.8	14
30	Novel and recurrent BRCA1/BRCA2 germline mutations in patients with breast/ovarian cancer: a series from the south of Tunisia. <i>Journal of Translational Medicine</i> , 2021, 19, 108.	4.4	14
31	Overexpression of COX-2 and LMP1 are correlated with lymph node in Tunisian NPC patients. <i>Oral Oncology</i> , 2008, 44, 710-715.	1.5	13
32	Clinicopathological and prognostic significance of p53, Ki-67, and Bcl-2 expression in Tunisian gastric adenocarcinomas. <i>Acta Histochemica</i> , 2014, 116, 1244-1250.	1.8	13
33	CpG methylation of APC promoter 1A in sporadic and familial breast cancer patients. <i>Cancer Biomarkers</i> , 2017, 18, 133-141.	1.7	13
34	High-level expression of human tumour suppressor P53 in the methylotrophic yeast: <i>Pichia pastoris</i> . <i>Protein Expression and Purification</i> , 2007, 54, 283-288.	1.3	12
35	High prevalence of the c.1227_1228dup (p.Glu410GlyfsX43) mutation in Tunisian families affected with MUTYH-associated-polyposis. <i>Familial Cancer</i> , 2012, 11, 503-508.	1.9	11
36	Frequent CpG methylation of ubiquitin carboxyl-terminal hydrolase 1 (UCHL1) in sporadic and hereditary Tunisian breast cancer patients: clinical significance. <i>Medical Oncology</i> , 2013, 30, 418.	2.5	11

#	ARTICLE	IF	CITATIONS
37	Clinical and prognosis value of the CIMP status combined with MLH1 or p16 INK4a methylation in colorectal cancer. <i>Medical Oncology</i> , 2017, 34, 147.	2.5	10
38	FOXA1 Expression in Nasopharyngeal Carcinoma: Association with Clinicopathological Characteristics and EMT Markers. <i>BioMed Research International</i> , 2020, 2020, 1-9.	1.9	10
39	Expression of p16INK4a, Alone or Combined With p53, is Predictive of Better Prognosis in Colorectal Adenocarcinoma in Tunisian Patients. <i>Applied Immunohistochemistry and Molecular Morphology</i> , 2011, 19, 562-568.	1.2	9
40	Cellular localization of human p53 expressed in the yeast <i>Saccharomyces cerevisiae</i> : effect of NLSI deletion. <i>Apoptosis: an International Journal on Programmed Cell Death</i> , 2011, 16, 746-756.	4.9	9
41	Glucose dependant negative translational control of the heterologous expression of the preS2 HBV antigen in yeast. <i>Gene</i> , 2003, 311, 165-170.	2.2	8
42	Methylation status and overexpression of COX-2 in Tunisian patients with ductal invasive breast carcinoma. <i>Tumor Biology</i> , 2011, 32, 461-468.	1.8	8
43	Characterization of C69R variant HBsAg: effect on binding to anti-HBs and the structure of virus-like particles. <i>Archives of Virology</i> , 2015, 160, 2427-2433.	2.1	8
44	Selection of cell death-deficient <i>p53</i> mutants in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 2009, 26, 441-450.	1.7	7
45	RIP140 and LCoR expression in gastrointestinal cancers. <i>Oncotarget</i> , 2017, 8, 111161-111175.	1.8	7
46	A novel pathogenic germline mutation in the adenomatous polyposis coli gene in a Tunisian family with FAP. <i>Familial Cancer</i> , 2011, 10, 567-571.	1.9	5
47	Expression of the human tumor suppressor p53 induces cell death in <i>Pichia pastoris</i> . <i>FEMS Yeast Research</i> , 2012, 12, 2-8.	2.3	5
48	Identification of a Novel Methylated Gene in Nasopharyngeal Carcinoma: TTC40. <i>BioMed Research International</i> , 2014, 2014, 1-7.	1.9	5
49	Overexpression of yeast thioredoxin TRX2 reduces p53-mediated cell death in yeast. <i>Applied Microbiology and Biotechnology</i> , 2015, 99, 8619-8628.	3.6	5
50	Clinical and prognosis relevance of COX-2 expression in Tunisian patients with primary gastric adenocarcinoma. <i>Cancer Biomarkers</i> , 2016, 17, 67-73.	1.7	4
51	Identification of novel candidate genes by exome sequencing in Tunisian familial male breast cancer patients. <i>Molecular Biology Reports</i> , 2020, 47, 6507-6516.	2.3	4
52	Establishment of primary cell culture of <i>Ruditapes decussatus</i> haemocytes for metal toxicity assessment. <i>In Vitro Cellular and Developmental Biology - Animal</i> , 2021, 57, 477-484.	1.5	4
53	Negative/low HER2 expression alone or combined with E-cadherin positivity is predictive of better prognosis in patients with breast carcinoma. <i>Histology and Histopathology</i> , 2012, 27, 377-85.	0.7	4
54	Phage-display screening identifies LMP1-binding peptides targeting the C-terminus region of the EBV oncoprotein. <i>Peptides</i> , 2016, 85, 73-79.	2.4	3

#	ARTICLE	IF	CITATIONS
55	Quantitative measurement of iNOS expression in melanoma, nasopharyngeal, colorectal, and breast tumors of Tunisian patients: comparative study and clinical significance. <i>Tumor Biology</i> , 2016, 37, 5153-5164.	1.8	3
56	Overexpression of the Oncogenic Variant (KLF6-SV1) in Young NPC Patients and Correlation with Lack of E-Cadherin. <i>Analytical Cellular Pathology</i> , 2018, 2018, 1-7.	1.4	3
57	B1.12: a novel peptide interacting with the extracellular loop of the EBV oncoprotein LMP1. <i>Scientific Reports</i> , 2019, 9, 4389.	3.3	3
58	Secreted recombinant P53 protein from <i>Pichia pastoris</i> is a useful antigen for detection of serum p53: autoantibody in patients with advanced colorectal adenocarcinoma. <i>Molecular Biology Reports</i> , 2013, 40, 3865-3872.	2.3	2
59	Lower p66Shc promoter methylation in subjects with chronic renal failure. <i>PLoS ONE</i> , 2021, 16, e0257176.	2.5	2
60	Whole and Purified Aqueous Extracts of <i>Nigella sativa</i> L. Seeds Attenuate Apoptosis and the Overproduction of Reactive Oxygen Species Triggered by p53 Over-Expression in the Yeast <i>Saccharomyces cerevisiae</i> . <i>Cells</i> , 2022, 11, 869.	4.1	1
61	Negative control glucose dependent mediated by the PreS2 region on the translation efficiency of the reporter <i>Sh-bleomycin</i> gene in <i>Saccharomyces cerevisiae</i> . <i>FEMS Yeast Research</i> , 2014, 14, 357-363.	2.3	0