## Juliana Chung-ngor Chan

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/7815248/publications.pdf

Version: 2024-02-01

787 papers 49,265 citations

97 h-index 190 g-index

808 all docs 808 docs citations

808 times ranked 49890 citing authors

#	Article	IF	Citations
1	IDF Diabetes Atlas: Global, regional and country-level diabetes prevalence estimates for 2021 and projections for 2045. Diabetes Research and Clinical Practice, 2022, 183, 109119.	2.8	2,873
2	Type 1 diabetes. Lancet, The, 2014, 383, 69-82.	13.7	1,863
3	Diabetes in Asia. JAMA - Journal of the American Medical Association, 2009, 301, 2129.	7.4	1,674
4	Effects of Once-Weekly Exenatide on Cardiovascular Outcomes in Type 2 Diabetes. New England Journal of Medicine, 2017, 377, 1228-1239.	27.0	1,455
5	A variant in CDKAL1 influences insulin response and risk of type 2 diabetes. Nature Genetics, 2007, 39, 770-775.	21.4	966
6	Genome-wide trans-ancestry meta-analysis provides insight into the genetic architecture of type 2 diabetes susceptibility. Nature Genetics, 2014, 46, 234-244.	21.4	959
7	The genetic architecture of type 2 diabetes. Nature, 2016, 536, 41-47.	27.8	952
8	Variants conferring risk of atrial fibrillation on chromosome 4q25. Nature, 2007, 448, 353-357.	27.8	853
9	Accuracy of Patient Health Questionnaire-9 (PHQ-9) for screening to detect major depression: individual participant data meta-analysis. BMJ: British Medical Journal, 2019, 365, 11476.	2.3	822
10	Sequence variants affecting eosinophil numbers associate with asthma and myocardial infarction. Nature Genetics, 2009, 41, 342-347.	21.4	709
11	Variants in KCNQ1 are associated with susceptibility to type 2 diabetes mellitus. Nature Genetics, 2008, 40, 1092-1097.	21.4	694
12	KDIGO 2020 Clinical Practice Guideline for Diabetes Management in Chronic Kidney Disease. Kidney International, 2020, 98, S1-S115.	5.2	692
13	Two variants on chromosome 17 confer prostate cancer risk, and the one in TCF2 protects against type 2 diabetes. Nature Genetics, 2007, 39, 977-983.	21.4	670
14	Type 2 diabetes in East Asians: similarities and differences with populations in Europe and the United States. Annals of the New York Academy of Sciences, 2013, 1281, 64-91.	3.8	606
15	Plasma glucose levels and diabetes are independent predictors for mortality and morbidity in patients with SARS. Diabetic Medicine, 2006, 23, 623-628.	2.3	604
16	Body mass index, waist circumference and waist:hip ratio as predictors of cardiovascular risk—a review of the literature. European Journal of Clinical Nutrition, 2010, 64, 16-22.	2.9	557
17	Meta-analysis of genome-wide association studies identifies eight new loci for type 2 diabetes in east Asians. Nature Genetics, 2012, 44, 67-72.	21.4	545
18	Effect of Finerenone on Albuminuria in Patients With Diabetic Nephropathy. JAMA - Journal of the American Medical Association, 2015, 314, 884.	7.4	523

#	Article	IF	Citations
19	Cardiovascular outcomes with glucagon-like peptide-1 receptor agonists in patients with type 2 diabetes: a meta-analysis. Lancet Diabetes and Endocrinology,the, 2018, 6, 105-113.	11.4	451
20	A sequence variant in ZFHX3 on 16q22 associates with atrial fibrillation and ischemic stroke. Nature Genetics, 2009, 41, 876-878.	21.4	434
21	Screening diabetic patients for non-alcoholic fatty liver disease with controlled attenuation parameter and liver stiffness measurements: a prospective cohort study. Gut, 2016, 65, 1359-1368.	12.1	386
22	Estimating the sample mean and standard deviation from commonly reported quantiles in meta-analysis. Statistical Methods in Medical Research, 2020, 29, 2520-2537.	1.5	366
23	Diabetes in Asia and the Pacific: Implications for the Global Epidemic. Diabetes Care, 2016, 39, 472-485.	8.6	363
24	Common variants near CAV1 and CAV2 are associated with primary open-angle glaucoma. Nature Genetics, 2010, 42, 906-909.	21.4	357
25	Implication of Genetic Variants Near <i>TCF7L2</i> , <i>SLC30A8</i> , <i>HHEX</i> , <i>CDKAL1</i> , <i cdkal1<="" i="">, <i>CDKAL1</i>, <i cdkal1<="" i="">, <i cdkal1<="" td=""></i></i></i></i></i></i></i></i></i></i></i></i></i></i></i></i></i></i></i></i></i></i></i></i></i></i></i></i></i></i></i></i></i></i></i></i></i></i></i></i></i></i>	0.6	331
26	The Lancet Commission on diabetes: using data to transform diabetes care and patient lives. Lancet, The, 2020, 396, 2019-2082.	13.7	327
27	Prediction of hypertension, diabetes, dyslipidaemia or albuminuria using simple anthropometric indexes in Hong Kong Chinese. International Journal of Obesity, 1999, 23, 1136-1142.	3.4	299
28	Effect of interventions for major depressive disorder and significant depressive symptoms in patients with diabetes mellitus: a systematic review and meta-analysis. General Hospital Psychiatry, 2010, 32, 380-395.	2.4	290
29	Identification of type 2 diabetes loci in 433,540 East Asian individuals. Nature, 2020, 582, 240-245.	27.8	282
30	Multi-ancestry genetic study of type 2 diabetes highlights the power of diverse populations for discovery and translation. Nature Genetics, 2022, 54, 560-572.	21.4	250
31	Multifaceted Determinants for Achieving Glycemic Control. Diabetes Care, 2009, 32, 227-233.	8.6	248
32	Exome sequencing of 20,791Âcases of type 2 diabetes and 24,440Âcontrols. Nature, 2019, 570, 71-76.	27.8	248
33	A genome-wide association study in the Japanese population identifies susceptibility loci for type 2 diabetes at UBE2E2 and C2CD4A-C2CD4B. Nature Genetics, 2010, 42, 864-868.	21.4	245
34	Effects of acarbose on cardiovascular and diabetes outcomes in patients with coronary heart disease and impaired glucose tolerance (ACE): a randomised, double-blind, placebo-controlled trial. Lancet Diabetes and Endocrinology,the, 2017, 5, 877-886.	11.4	245
35	In Utero Exposure to Maternal Hyperglycemia Increases Childhood Cardiometabolic Risk in Offspring. Diabetes Care, 2017, 40, 679-686.	8.6	242
36	Accuracy of the PHQ-2 Alone and in Combination With the PHQ-9 for Screening to Detect Major Depression. JAMA - Journal of the American Medical Association, 2020, 323, 2290.	7.4	242

#	Article	IF	Citations
37	Safety and efficacy of sitagliptin in patients with type 2 diabetes and chronic renal insufficiency. Diabetes, Obesity and Metabolism, 2008, 10, 545-555.	4.4	222
38	Effects of olmesartan on renal and cardiovascular outcomes in type 2 diabetes with overt nephropathy: a multicentre, randomised, placebo-controlled study. Diabetologia, 2011, 54, 2978-2986.	6.3	211
39	Prevalence of Gestational Diabetes Mellitus and Its Risk Factors in Chinese Pregnant Women: A Prospective Population-Based Study in Tianjin, China. PLoS ONE, 2015, 10, e0121029.	2.5	211
40	Metabolic profiles and treatment gaps in young-onset type 2 diabetes in Asia (the JADE programme): a cross-sectional study of a prospective cohort. Lancet Diabetes and Endocrinology,the, 2014, 2, 935-943.	11.4	210
41	Development of the Telemedicine Satisfaction Questionnaire to evaluate patient satisfaction with telemedicine: a preliminary study. Journal of Telemedicine and Telecare, 2003, 9, 46-50.	2.7	202
42	Glomerular Filtration Rate, Cardiorenal End Points, and All-Cause Mortality in Type 2 Diabetic Patients. Diabetes Care, 2006, 29, 2046-2052.	8.6	196
43	Erectile Dysfunction Predicts Coronary Heart Disease in Type 2 Diabetes. Journal of the American College of Cardiology, 2008, 51, 2045-2050.	2.8	193
44	White Blood Cell Count Is Associated With Macro- and Microvascular Complications in Chinese Patients With Type 2 Diabetes. Diabetes Care, 2004, 27, 216-222.	8.6	185
45	Equivalency of the diagnostic accuracy of the PHQ-8 and PHQ-9: a systematic review and individual participant data meta-analysis. Psychological Medicine, 2020, 50, 1368-1380.	4.5	175
46	Whole-genome bisulfite sequencing of multiple individuals reveals complementary roles of promoter and gene body methylation in transcriptional regulation. Genome Biology, 2014, 15, 408.	8.8	173
47	The Reproducibility and Usefulness of the Oral Glucose Tolerance Test in Screening for Diabetes and other Cardiovascular Risk Factors. Annals of Clinical Biochemistry, 1998, 35, 62-67.	1.6	171
48	Factor analysis of the metabolic syndrome: obesity vs insulin resistance as the central abnormality. International Journal of Obesity, 2001, 25, 1782-1788.	3.4	171
49	Association of genetic variation in FTO with risk of obesity and type 2 diabetes with data from 96,551 East and South Asians. Diabetologia, 2012, 55, 981-995.	6.3	171
50	Effectiveness of telephone counselling by a pharmacist in reducing mortality in patients receiving polypharmacy: randomised controlled trial. BMJ: British Medical Journal, 2006, 333, 522.	2.3	168
51	Diabetes in China: a societal solution for a personal challenge. Lancet Diabetes and Endocrinology,the, 2014, 2, 969-979.	11.4	168
52	Lower BMI cut-off value to define obesity in Hong Kong Chinese: an analysis based on body fat assessment by bioelectrical impedance. British Journal of Nutrition, 2001, 85, 239-242.	2.3	164
53	Chinese herbal medicines revisited: a Hong Kong perspective. Lancet, The, 1993, 342, 1532-1534.	13.7	161
54	Metabolic Syndrome Predicts New Onset of Chronic Kidney Disease in 5,829 Patients With Type 2 Diabetes. Diabetes Care, 2008, 31, 2357-2361.	8.6	160

#	Article	IF	CITATIONS
55	Long-term effects of angiotensin-converting enzyme inhibition and metabolic control in hypertensive type 2 diabetic patients. Kidney International, 2000, 57, 590-600.	5.2	158
56	Genome-wide association study identifies three novel loci for type 2 diabetes. Human Molecular Genetics, 2014, 23, 239-246.	2.9	158
57	Sulodexide Fails to Demonstrate Renoprotection in Overt Type 2 Diabetic Nephropathy. Journal of the American Society of Nephrology: JASN, 2012, 23, 123-130.	6.1	151
58	Hepatitis C infection in African Americans: its natural history and histological progression. American Journal of Gastroenterology, 2002, 97, 700-706.	0.4	150
59	Genome-wide association studies in the Japanese population identify seven novel loci for type 2 diabetes. Nature Communications, 2016, 7, 10531.	12.8	149
60	Engineering of Targeted Nanoparticles for Cancer Therapy Using Internalizing Aptamers Isolated by Cell-Uptake Selection. ACS Nano, 2012, 6, 696-704.	14.6	148
61	Associations of Hyperglycemia and Insulin Usage With the Risk of Cancer in Type 2 Diabetes: The Hong Kong Diabetes Registry. Diabetes, 2010, 59, 1254-1260.	0.6	145
62	Association between sleeping hours, working hours and obesity in Hong Kong Chinese: the â€better health for better Hong Kong' health promotion campaign. International Journal of Obesity, 2007, 31, 254-260.	3.4	143
63	Diabetes Mellitus. Stroke, 2013, 44, 1500-1504.	2.0	143
64	Impact of age at type 2 diabetes mellitus diagnosis on mortality and vascular complications: systematic review and meta-analyses. Diabetologia, 2021, 64, 275-287.	6.3	140
65	A Genome-Wide Association Study of Diabetic Kidney Disease in Subjects With Type 2 Diabetes. Diabetes, 2018, 67, 1414-1427.	0.6	136
66	Glucose Intolerance and Cardiometabolic Risk in Children Exposed to Maternal Gestational Diabetes Mellitus in Utero. Pediatrics, 2008, 122, 1229-1234.	2.1	135
67	Associations of sleep duration with obesity and serum lipid profile in children and adolescents. Sleep Medicine, 2011, 12, 659-665.	1.6	133
68	Chinese herbs and warfarin potentiation by â€~Danshen'. Journal of Internal Medicine, 1997, 241, 337-339.	6.0	131
69	Personalized Management of Hyperglycemia in Type 2 Diabetes: Reflections from a Diabetes Care Editors' Expert Forum. Diabetes Care, 2013, 36, 1779-1788.	8.6	130
70	Diabetes and its comorbidities—where East meets West. Nature Reviews Endocrinology, 2013, 9, 537-547.	9.6	124
71	Comparison of enalapril and nifedipine in treating non-insulin dependent diabetes associated with hypertension: one year analysis BMJ: British Medical Journal, 1992, 305, 981-985.	2.3	122
72	Effects of Structured Versus Usual Care on Renal Endpoint in Type 2 Diabetes: The SURE Study. Diabetes Care, 2009, 32, 977-982.	8.6	122

#	Article	IF	CITATIONS
73	Effects of Telephone-Based Peer Support in Patients With Type 2 Diabetes Mellitus Receiving Integrated Care. JAMA Internal Medicine, 2014, 174, 972.	5.1	121
74	Phenotypic Heterogeneity and Associations of Two Aldose Reductase Gene Polymorphisms With Nephropathy and Retinopathy in Type 2 Diabetes. Diabetes Care, 2003, 26, 2410-2415.	8.6	120
75	The Usefulness of the International Diabetes Federation and the National Cholesterol Education Program's Adult Treatment Panel III Definitions of the Metabolic Syndrome in Predicting Coronary Heart Disease in Subjects With Type 2 Diabetes. Diabetes Care, 2007, 30, 1206-1211.	8.6	120
76	Implication of Genetic Variants Near <i>NEGR1</i> , <i>SEC16B</i> , <i>TMEM18</i> , <i>ETV5/DGKG</i> , <i>GNPDA2</i> , <i>LIN7C/BDNF</i> , <i>Nand<i>KCTD15</i>)with Obesity and Type 2 Diabetes in 7705 Chinese. Journal of Clinical Endocrinology and Metabolism, 2010, 95, 2418-2425.</i>	ITCH2	, <i>BCDIN3D/ 120</i>
77	Combined Use of a Fasting Plasma Glucose Concentration and HbA1c or Fructosamine Predicts the Likelihood of Having Diabetes in High-Risk Subjects. Diabetes Care, 1998, 21, 1221-1225.	8.6	118
78	Risk association of HbA $<$ sub $>1csub> variability with chronic kidney disease and cardiovascular disease in type 2 diabetes: prospective analysis of the Hong Kong Diabetes Registry. Diabetes/Metabolism Research and Reviews, 2013, 29, 384-390.$	4.0	118
79	Sonographic measurement of mesenteric fat thickness is a good correlate with cardiovascular risk factors: comparison with subcutaneous and preperitoneal fat thickness, magnetic resonance imaging and anthropometric indexes. International Journal of Obesity, 2003, 27, 1267-1273.	3.4	116
80	Abnormal Liver Function Test Predicts Type 2 Diabetes. Diabetes Care, 2007, 30, 2566-2568.	8.6	116
81	A systematic review on use of Chinese medicine and acupuncture for treatment of obesity. Obesity Reviews, 2012, 13, 409-430.	6.5	114
82	The central roles of obesity-associated dyslipidaemia, endothelial activation and cytokines in the Metabolic Syndrome—an analysis by structural equation modelling. International Journal of Obesity, 2002, 26, 994-1008.	3.4	113
83	Asymmetric dimethylarginine (ADMA): a potential link between endothelial dysfunction and cardiovascular diseases in insulin resistance syndrome?. Diabetologia, 2002, 45, 1609-1616.	6.3	113
84	Waist circumference and body mass index in Chinese children: cutoff values for predicting cardiovascular risk factors. International Journal of Obesity, 2007, 31, 550-558.	3.4	113
85	Prevalence and Clinicopathological Characteristics of Islet Amyloid in Chinese Patients With Type 2 Diabetes. Diabetes, 2003, 52, 2759-2766.	0.6	112
86	Simple anthropometric indexes and cardiovascular risk factors in Chinese. International Journal of Obesity, 1997, 21, 995-1001.	3.4	110
87	Premature Mortality and Comorbidities in Young-onset Diabetes: A 7-Year Prospective Analysis. American Journal of Medicine, 2014, 127, 616-624.	1.5	110
88	Diabetes Management in Chronic Kidney Disease: Synopsis of the 2020 KDIGO Clinical Practice Guideline. Annals of Internal Medicine, 2021, 174, 385-394.	3.9	110
89	Genome-wide Scan for Metabolic Syndrome and Related Quantitative Traits in Hong Kong Chinese and Confirmation of a Susceptibility Locus on Chromosome 1q21-q25. Diabetes, 2004, 53, 2676-2683.	0.6	107
90	Progression of diabetic kidney disease and trajectory of kidney function decline in Chinese patients with Type 2 diabetes. Kidney International, 2019, 95, 178-187.	5.2	105

#	Article	IF	CITATIONS
91	The A1C and ABCD of glycaemia management in type 2 diabetes: a physician's personalized approach. Diabetes/Metabolism Research and Reviews, 2010, 26, 239-244.	4.0	104
92	Sustained Antidiabetic Effects of a Berberine-Containing Chinese Herbal Medicine Through Regulation of Hepatic Gene Expression. Diabetes, 2012, 61, 933-943.	0.6	103
93	Urinary epinephrine and norepinephrine interrelations with obesity, insulin, and the metabolic syndrome in Hong Kong Chinese. Metabolism: Clinical and Experimental, 2001, 50, 135-143.	3.4	102
94	Development and Validation of Stroke Risk Equation for Hong Kong Chinese Patients With Type 2 Diabetes: The Hong Kong Diabetes Registry. Diabetes Care, 2007, 30, 65-70.	8.6	102
95	The prevalence of diabetes mellitus and impaired glucose tolerance among Hong Kong Chinese adults of working age. Diabetes Research and Clinical Practice, 1993, 21, 67-73.	2.8	101
96	Aberrant activation profile of cytokines and mitogen-activated protein kinases in type 2 diabetic patients with nephropathy. Clinical and Experimental Immunology, 2007, 149, 123-131.	2.6	101
97	Development and Validation of a Total Coronary Heart Disease Risk Score in Type 2 Diabetes Mellitus. American Journal of Cardiology, 2008, 101, 596-601.	1.6	101
98	Replication and Identification of Novel Variants at TCF7L2 Associated with Type 2 Diabetes in Hong Kong Chinese. Journal of Clinical Endocrinology and Metabolism, 2007, 92, 3733-3737.	3.6	100
99	A mitochondrial DNA variant at position 16189 is associated with type 2 diabetes mellitus in Asians. Diabetologia, 2008, 51, 602-608.	6.3	100
100	Insulin glargine versus sitagliptin in insulin-naive patients with type 2 diabetes mellitus uncontrolled on metformin (EASIE): a multicentre, randomised open-label trial. Lancet, The, 2012, 379, 2262-2269.	13.7	100
101	Drug-Induced Disorders of Glucose Metabolism. Drug Safety, 1996, 15, 135-157.	3.2	98
102	Burden of Obesity – lessons learnt from Hong Kong Chinese. Obesity Reviews, 2008, 9, 35-40.	6.5	97
103	Glucose Intolerance and Cardiometabolic Risk in Adolescents Exposed to Maternal Gestational Diabetes. Diabetes Care, 2010, 33, 1382-1384.	8.6	97
104	Toxicity of Complementary Therapies: An Eastern Perspective. Journal of Clinical Pharmacology, 2000, 40, 451-456.	2.0	96
105	Renin-Angiotensin System Gene Polymorphisms, Blood Pressure, Dyslipidemia, and Diabetes in Hong Kong Chinese. Diabetes Care, 2001, 24, 356-361.	8.6	96
106	Mesenteric Fat Thickness Is an Independent Determinant of Metabolic Syndrome and Identifies Subjects With Increased Carotid Intima-Media Thickness. Diabetes Care, 2006, 29, 379-384.	8.6	94
107	Development and Validation of an All-Cause Mortality Risk Score in Type 2 Diabetes <subtitle>The Hong Kong Diabetes Registry</subtitle> . Archives of Internal Medicine, 2008, 168, 451.	3.8	94
108	Genome-wide association study in a Chinese population identifies a susceptibility locus for type 2 diabetes at 7q32 near PAX4. Diabetologia, 2013, 56, 1291-1305.	6.3	94

#	Article	IF	CITATIONS
109	Renal risk and renoprotection among ethnic groups with type 2 diabetic nephropathy: A post hoc analysis of RENAAL. Kidney International, 2006, 69, 1675-1682.	5.2	92
110	Co-occurrence of diabetes and depression: Conceptual considerations for an emerging global health challenge. Journal of Affective Disorders, 2012, 142, S56-S66.	4.1	92
111	Diabetes and pregnancy: perspectives from Asia. Diabetic Medicine, 2014, 31, 302-318.	2.3	92
112	Gastrointestinal symptoms in Chinese patients with Type 2 diabetes mellitus. Diabetic Medicine, 1999, 16, 670-674.	2.3	90
113	Weight management and current options in pharmacotherapy: Orlistat and sibutramine. Clinical Therapeutics, 2003, 25, 58-80.	2.5	88
114	High prevalence of metabolic syndrome in Hong Kong Chineseâ€"comparison of three diagnostic criteria. Diabetes Research and Clinical Practice, 2005, 69, 160-168.	2.8	88
115	Phenotypic and genetic clustering of diabetes and metabolic syndrome in Chinese families with type 2 diabetes mellitus. Diabetes/Metabolism Research and Reviews, 2006, 22, 46-52.	4.0	88
116	Up-Regulated Pancreatic Tissue MicroRNA-375 Associates With Human Type 2 Diabetes Through Î <sup>2</sup> -Cell Deficit and Islet Amyloid Deposition. Pancreas, 2010, 39, 843-846.	1.1	88
117	Fundamental Concepts Regarding Testosterone Deficiency and Treatment. Mayo Clinic Proceedings, 2016, 91, 881-896.	3.0	88
118	Phycocyanin protects INS-1E pancreatic beta cells against human islet amyloid polypeptide-induced apoptosis through attenuating oxidative stress and modulating JNK and p38 mitogen-activated protein kinase pathways. International Journal of Biochemistry and Cell Biology, 2009, 41, 1526-1535.	2.8	87
119	Prospective Study on the Incidences of Cardiovascular-Renal Complications in Chinese Patients With Young-Onset Type 1 and Type 2 Diabetes. Diabetes Care, 2014, 37, 149-157.	8.6	87
120	Trends in the incidence of diagnosed diabetes: a multicountry analysis of aggregate data from 22 million diagnoses in high-income and middle-income settings. Lancet Diabetes and Endocrinology,the, 2021, 9, 203-211.	11.4	85
121	Plasma insulin, growth hormone, cortisol, and central obesity among young Chinese type 2 diabetic patients. Diabetes Care, 1999, 22, 1450-1457.	8.6	83
122	Rationale and design of the EXenatide Study of Cardiovascular Event Lowering (EXSCEL) trial. American Heart Journal, 2016, 174, 103-110.	2.7	82
123	Obesity, albuminuria and hypertension among Hong Kong Chinese with non-insulin-dependent diabetes mellitus (NIDDM). Postgraduate Medical Journal, 1993, 69, 204-210.	1.8	81
124	Evidence for DNA Damage as a Biological Link Between Diabetes and Cancer. Chinese Medical Journal, 2015, 128, 1543-1548.	2.3	81
125	Depression in <scp>C</scp> hinese patients with type 2 diabetes: associations with hyperglycemia, hypoglycemia, and poor treatment adherence. Journal of Diabetes, 2015, 7, 800-808.	1.8	81
126	Aspects of Multicomponent Integrated Care Promote Sustained Improvement in Surrogate Clinical Outcomes: A Systematic Review and Meta-analysis. Diabetes Care, 2018, 41, 1312-1320.	8.6	81

#	Article	IF	Citations
127	European Bone Mineral Density Loci Are Also Associated with BMD in East-Asian Populations. PLoS ONE, 2010, 5, e13217.	2.5	81
128	Use of sulphonylurea and cancer in type 2 diabetesâ€"The Hong Kong Diabetes Registry. Diabetes Research and Clinical Practice, 2010, 90, 343-351.	2.8	80
129	Declining Trends of Cardiovascular-Renal Complications and Mortality in Type 2 Diabetes: The Hong Kong Diabetes Database. Diabetes Care, 2017, 40, 928-935.	8.6	80
130	Secular trends in all-cause and cause-specific mortality rates in people with diabetes in Hong Kong, 2001–2016: a retrospective cohort study. Diabetologia, 2020, 63, 757-766.	6.3	80
131	Cigarette smoking is an independent risk factor for type 2 diabetes: a fourâ€year communityâ€based prospective study. Clinical Endocrinology, 2009, 71, 679-685.	2.4	79
132	Diabetes in the Western Pacific Regionâ€"Past, Present and Future. Diabetes Research and Clinical Practice, 2014, 103, 244-255.	2.8	79
133	Measuring depression with CES-D in Chinese patients with type 2 diabetes: the validity and its comparison to PHQ-9. BMC Psychiatry, 2015, 15, 198.	2.6	79
134	From Hong Kong Diabetes Register to JADE Program to RAMP-DM for Data-Driven Actions. Diabetes Care, 2019, 42, 2022-2031.	8.6	79
135	Metabolic and Hemodynamic Effects of Metformin and Glibenclamide in Normotensive NIDDM Patients. Diabetes Care, 1993, 16, 1035-1038.	8.6	77
136	Familial Early-Onset Type 2 Diabetes in Chinese Patients: Obesity and genetics have more significant roles than autoimmunity. Diabetes Care, 2001, 24, 663-671.	8.6	77
137	The associations of body mass index, C-peptide and metabolic status in Chinese Type 2 diabetic patients. Diabetic Medicine, 2004, 21, 349-353.	2.3	77
138	Exendinâ€4 protects pancreatic beta cells from human islet amyloid polypeptideâ€induced cell damage: potential involvement of AKT and mitochondria biogenesis. Diabetes, Obesity and Metabolism, 2010, 12, 815-824.	4.4	77
139	Persistent poor glycaemic control in individuals with type 2 diabetes in developing countries: 12Âyears of real-world evidence of the International Diabetes Management Practices Study (IDMPS). Diabetologia, 2020, 63, 711-721.	6.3	76
140	Diabetes in the Chinese Population and Its Implications for Health Care. Diabetes Care, 1997, 20, 1785-1790.	8.6	74
141	The Trp64Arg polymorphism of the $\hat{l}^2$ 3-adrenergic receptor gene and obesity in Chinese subjects with components of the metabolic syndrome. International Journal of Obesity, 2000, 24, 545-551.	3.4	73
142	Renin Angiotensin Aldosterone System Blockade and Renal Disease in Patients With Type 2 Diabetes: An Asian perspective from the RENAAL study. Diabetes Care, 2004, 27, 874-879.	8.6	73
143	Independent associations between low-density lipoprotein cholesterol and cancer among patients with type 2 diabetes mellitus. Cmaj, 2008, 179, 427-437.	2.0	73
144	Testosterone level in men with typeÂ2 diabetes mellitus and related metabolic effects: A review of current evidence. Journal of Diabetes Investigation, 2015, 6, 112-123.	2.4	73

#	Article	IF	CITATIONS
145	Antibodies to Glutamic Acid Decarboxylase in Young Chinese Diabetic Patients. Annals of Clinical Biochemistry, 1998, 35, 761-767.	1.6	72
146	Diabetes-Related Distress and Physical and Psychological Health in Chinese Type 2 Diabetic Patients. Diabetes Care, 2011, 34, 1094-1096.	8.6	72
147	Advanced liver fibrosis but not steatosis is independently associated with albuminuria in Chinese patients with type 2 diabetes. Journal of Hepatology, 2018, 68, 147-156.	3.7	72
148	Construction of a prediction model for type 2 diabetes mellitus in the Japanese population based on 11 genes with strong evidence of the association. Journal of Human Genetics, 2009, 54, 236-241.	2.3	70
149	Precision medicine in diabetes prevention, classification and management. Journal of Diabetes Investigation, 2018, 9, 998-1015.	2.4	69
150	End-stage renal disease risk equations for Hong Kong Chinese patients with type 2 diabetes: Hong Kong Diabetes Registry. Diabetologia, 2006, 49, 2299-2308.	6.3	68
151	Involvement of mitochondrial dysfunction in human islet amyloid polypeptide-induced apoptosis in INS-1E pancreatic beta cells: An effect attenuated by phycocyanin. International Journal of Biochemistry and Cell Biology, 2011, 43, 525-534.	2.8	67
152	Associations between microRNA (miR-21, 126, 155 and 221), albuminuria and heavy metals in Hong Kong Chinese adolescents. Clinica Chimica Acta, 2012, 413, 1053-1057.	1,1	67
153	The Accuracy of the Patient Health Questionnaire-9 Algorithm for Screening to Detect Major Depression: An Individual Participant Data Meta-Analysis. Psychotherapy and Psychosomatics, 2020, 89, 25-37.	8.8	67
154	The Metabolic Syndrome in Hong Kong Chinese. The interrelationships among its components analyzed by structural equation modeling. Diabetes Care, 1996, 19, 953-959.	8.6	65
155	Common Polymorphisms in MTNR1B, G6PC2 and GCK Are Associated with Increased Fasting Plasma Glucose and Impaired Beta-Cell Function in Chinese Subjects. PLoS ONE, 2010, 5, e11428.	2.5	65
156	Berberine modulates insulin signaling transduction in insulin-resistant cells. Molecular and Cellular Endocrinology, 2010, 317, 148-153.	3.2	65
157	Low HDL Cholesterol, Metformin Use, and Cancer Risk in Type 2 Diabetes. Diabetes Care, 2011, 34, 375-380.	8.6	65
158	Effects of structured care by a pharmacist-diabetes specialist team in patients with Type 2 diabetic nephropathy. American Journal of Medicine, 2005, 118, 1414.e21-1414.e27.	1.5	64
159	Hematocrit, Independent of Chronic Kidney Disease, Predicts Adverse Cardiovascular Outcomes in Chinese Patients With Type 2 Diabetes. Diabetes Care, 2006, 29, 2439-2444.	8.6	64
160	Visceral Fat and Cardiovascular Risk Factors in Chinese NIDDM Patients. Diabetes Care, 1997, 20, 1854-1858.	8.6	63
161	Microvascular and Cardiovascular Outcomes According to Renal Function in Patients Treated With Once-Weekly Exenatide: Insights From the EXSCEL Trial. Diabetes Care, 2020, 43, 446-452.	8.6	63
162	Pancreatic $\hat{l}^2$ cell function and antibodies to glutamic acid decarboxylase (anti-GAD) in Chinese patients with clinical diagnosis of insulin-dependent diabetes mellitus. Diabetes Research and Clinical Practice, 1996, 32, 27-34.	2.8	62

#	Article	IF	CITATIONS
163	Antihypertensive and Anti-Albuminuric Effects of Losartan Potassium and Felodipine in Chinese Elderly Hypertensive Patients with or without IMon-Insulin-Dependent Diabetes mellitus. American Journal of Nephrology, 1997, 17, 72-80.	3.1	62
164	Pathophysiology, phenotypes and management of type 2 diabetes mellitus in Indian and Chinese populations. Nature Reviews Endocrinology, 2022, 18, 413-432.	9.6	62
165	Severe Hypoglycemia Identifies Vulnerable Patients With Type 2 Diabetes at Risk for Premature Death and All-Site Cancer: The Hong Kong Diabetes Registry. Diabetes Care, 2014, 37, 1024-1031.	8.6	61
166	Tumor necrosis factor alpha gene G-308A polymorphism in the metabolic syndrome. Metabolism: Clinical and Experimental, 2000, 49, 1021-1024.	3.4	60
167	Causes of Death in a Contemporary Cohort of Patients With Type 2 Diabetes and Atherosclerotic Cardiovascular Disease: Insights From the TECOS Trial. Diabetes Care, 2017, 40, 1763-1770.	8.6	60
168	Antiâ€inflammatory effects of exendinâ€4, a glucagonâ€like peptideâ€1 analog, on human peripheral lymphocytes in patients with type 2 diabetes. Journal of Diabetes Investigation, 2013, 4, 382-392.	2.4	59
169	Genome-Wide Association Meta-analysis Identifies Novel Variants Associated With Fasting Plasma Glucose in East Asians. Diabetes, 2015, 64, 291-298.	0.6	59
170	The interdisciplinary team in type 2 diabetes management: Challenges and best practice solutions from real-world scenarios. Journal of Clinical and Translational Endocrinology, 2017, 7, 21-27.	1.4	59
171	Increased Risk of Severe Hypoglycemic Events Before and After Cardiovascular Outcomes in TECOS Suggests an At-Risk Type 2 Diabetes Frail Patient Phenotype. Diabetes Care, 2018, 41, 596-603.	8.6	59
172	Serial Transient Elastography Examinations to Monitor Patients With Type 2 Diabetes: A Prospective Cohort Study. Hepatology, 2020, 72, 1230-1241.	7.3	59
173	Interaction Effect of Genetic Polymorphisms in Glucokinase ( <i>GCK</i> ) and Glucokinase Regulatory Protein ( <i>GCKR</i> ) on Metabolic Traits in Healthy Chinese Adults and Adolescents. Diabetes, 2009, 58, 765-769.	0.6	58
174	Genetic Variants of the Protein Kinase $C \cdot \hat{l}^2$ 1 Gene and Development of End-Stage Renal Disease in Patients With Type 2 Diabetes. JAMA - Journal of the American Medical Association, 2010, 304, 881.	7.4	58
175	Angiotensinogen T235 and ACE Insertion/Deletion Polymorphisms Associated With Albuminuria in Chinese Type 2 Diabetic Patients. Diabetes Care, 1998, 21, 431-437.	8.6	57
176	Combined use of fasting plasma glucose and HbA1c predicts the progression to diabetes in Chinese subjects. Diabetes Care, 2000, 23, 1770-1773.	8.6	57
177	The Effect of Orlistat-Induced Weight Loss, Without Concomitant Hypocaloric Diet, on Cardiovascular Risk Factors and Insulin Sensitivity in Young Obese Chinese Subjects With or Without Type 2 Diabetes. Archives of Internal Medicine, 2002, 162, 2428.	3.8	57
178	Genome-wide Scan for Type 2 Diabetes Loci in Hong Kong Chinese and Confirmation of a Susceptibility Locus on Chromosome 1q21-q25. Diabetes, 2004, 53, 1609-1613.	0.6	57
179	The linkage and association of the gene encoding upstream stimulatory factor 1 with type 2 diabetes and metabolic syndrome in the Chinese population. Diabetologia, 2005, 48, 2018-2024.	6.3	57
180	Association of plasminogen activator inhibitor-1 4G/4G genotype and type 2 diabetic nephropathy in Chinese patients. Kidney International, 2000, 57, 632-638.	5.2	56

#	Article	IF	Citations
181	The Complexity of Vascular and Non-Vascular Complications of Diabetes: The Hong Kong Diabetes Registry. Current Cardiovascular Risk Reports, 2011, 5, 230-239.	2.0	56
182	Patients' education, and its impact on care outcomes, resource consumption and working conditions: Data from the International Diabetes Management Practices Study (IDMPS). Diabetes and Metabolism, 2012, 38, 128-134.	2.9	56
183	Effects of Providing Peer Support on Diabetes Management in People With Type 2 Diabetes. Annals of Family Medicine, 2015, 13, S42-S49.	1.9	56
184	BMI and Waist Circumference in Predicting Cardiovascular Risk Factor Clustering in Chinese Adolescents. Obesity, 2007, 15, 494-494.	3.0	55
185	Fat redistribution and adipocyte transformation in uninephrectomized rats. Kidney International, 2008, 74, 467-477.	5.2	55
186	Use of the 1997 American Diabetes Association Diagnostic Criteria for Diabetes in a Hong Kong Chinese Population. Diabetes Care, 1998, 21, 2094-2097.	8.6	54
187	Measuring depressive symptoms using the Patient Health Questionnaire-9 in Hong Kong Chinese subjects with type 2 diabetes. Journal of Affective Disorders, 2013, 151, 660-666.	4.1	54
188	Short-term association between ambient temperature and acute myocardial infarction hospitalizations for diabetes mellitus patients: A time series study. PLoS Medicine, 2018, 15, e1002612.	8.4	54
189	High Prevalence of Insulin Resistance and Metabolic Syndrome in Overweight/Obese Preadolescent Hong Kong Chinese Children Aged 9-12 Years. Diabetes Care, 2003, 26, 250-251.	8.6	53
190	Angiotensin-converting enzyme (ACE) inhibition in type 2, diabetic patients – interaction with ACE insertion/deletion polymorphism. Kidney International, 2006, 69, 1438-1443.	5 <b>.</b> 2	53
191	Bip overexpression, but not CHOP inhibition, attenuates fatty-acid-induced endoplasmic reticulum stress and apoptosis in HepG2 liver cells. Life Sciences, 2010, 87, 724-732.	4.3	53
192	A randomised translational trial of lifestyle intervention using a 3-tier shared care approach on pregnancy outcomes in Chinese women with gestational diabetes mellitus but without diabetes. Journal of Translational Medicine, 2014, 12, 290.	4.4	53
193	Probability of major depression diagnostic classification using semi-structured versus fully structured diagnostic interviews. British Journal of Psychiatry, 2018, 212, 377-385.	2.8	53
194	Excess Burden of Mental Illness and Hospitalization in Young-Onset Type 2 Diabetes. Annals of Internal Medicine, 2019, 170, 145.	3.9	53
195	Abnormal Albuminuria as a Predictor of Mortality and Renal Impairment in Chinese Patients With NIDDM. Diabetes Care, 1995, 18, 1013-1016.	8.6	52
196	Molecular genetics of diabetes mellitus in Chinese subjects: identification of mutations in glucokinase and hepatocyte nuclear factor-1alpha genes in patients with early-onset Type 2 diabetes mellitus/MODY. Diabetic Medicine, 1999, 16, 956-963.	2.3	52
197	Lack of association of angiotensin-converting enzyme (Dd/II) and angiotensinogen M235T gene polymorphism with renal function among Chinese patients with type II diabetes. American Journal of Kidney Diseases, 1999, 33, 1064-1070.	1.9	52
198	Development and validation of a risk score for hospitalization for heart failure in patients with Type 2 Diabetes Mellitus. Cardiovascular Diabetology, 2008, 7, 9.	6.8	52

#	Article	IF	CITATIONS
199	Gestational Diabetes, Maternal Obesity, and the NCD Burden. Clinical Obstetrics and Gynecology, 2013, 56, 633-641.	1.1	52
200	Risk factors in Vâ€shaped risk associations with allâ€cause mortality in type 2 diabetesâ€"The Hong Kong Diabetes Registry. Diabetes/Metabolism Research and Reviews, 2008, 24, 238-246.	4.0	51
201	Pharmacological reduction of NEFA restores the efficacy of incretin-based therapies through GLP-1 receptor signalling in the beta cell in mouse models of diabetes. Diabetologia, 2013, 56, 423-433.	6.3	51
202	An Asian Multicenter Clinical Trial to Assess the Efficacy and Tolerability of Acarbose Compared With Placebo in Type 2 Diabetic Patients Previously Treated With Diet. Diabetes Care, 1998, 21, 1058-1061.	8.6	50
203	A low socio-economic status is an additional risk factor for glucose intolerance in high risk Hong Kong Chinese. European Journal of Epidemiology, 2001, 17, 289-295.	5.7	50
204	Reducing Cardiovascular Risk in Type 2 Diabetes. New England Journal of Medicine, 2003, 348, 457-459.	27.0	50
205	Association of Testosterone, Insulin-Like Growth Factor-I, and C-Reactive Protein with Metabolic Syndrome in Chinese Middle-Aged Men with a Family History of Type 2 Diabetes. Journal of Clinical Endocrinology and Metabolism, 2005, 90, 6418-6423.	3.6	50
206	Chronic hepatitis B viral infection independently predicts renal outcome in type 2 diabetic patients. Diabetologia, 2006, 49, 1777-1784.	6.3	50
207	Protein Kinase Cζ Mediates Insulin-induced Glucose Transport through Actin Remodeling in L6 Muscle Cells. Molecular Biology of the Cell, 2006, 17, 2322-2330.	2.1	50
208	Rationale for and design of the Acarbose Cardiovascular Evaluation (ACE) trial. American Heart Journal, 2014, 168, 23-29.e2.	2.7	50
209	Presence of mitochondrial tRNA Leu(UUR) A to G 3243 mutation in DNA extracted from serum and plasma of patients with type 2 diabetes mellitus. Journal of Clinical Pathology, 2000, 53, 466-469.	2.0	49
210	Association of carotid intima-media thickness with mesenteric, preperitoneal and subcutaneous fat thickness. Atherosclerosis, 2005, 179, 299-304.	0.8	49
211	Mesenteric fat thickness as an independent determinant of fatty liver. International Journal of Obesity, 2006, 30, 787-793.	3.4	49
212	Overweight, family history of diabetes and attending schools of lower academic grading are independent predictors for metabolic syndrome in Hong Kong Chinese adolescents. Archives of Disease in Childhood, 2007, 92, 224-228.	1.9	49
213	Pregnancy and diabetes scenario around the world: China. International Journal of Gynecology and Obstetrics, 2009, 104, S42-5.	2.3	49
214	Baseline characteristics of patients enrolled in the Exenatide Study of Cardiovascular Event Lowering (EXSCEL). American Heart Journal, 2017, 187, 1-9.	2.7	49
215	Secular trends in incidence of type $1$ and type $2$ diabetes in Hong Kong: A retrospective cohort study. PLoS Medicine, 2020, $17$ , e1003052.	8.4	49
216	Determinants of penetrance and variable expressivity in monogenic metabolic conditions across 77,184 exomes. Nature Communications, 2021, 12, 3505.	12.8	49

#	Article	IF	CITATIONS
217	The islet amyloid polypeptide (amylin) gene S20G mutation in Chinese subjects: Evidence for associations with type 2 diabetes and cholesterol levels. Clinical Endocrinology, 2001, 54, 541-546.	2.4	48
218	Prognostic Effect of Insertion/Deletion Polymorphism of the ACE Gene on Renal and Cardiovascular Clinical Outcomes in Chinese Patients With Type 2 Diabetes. Diabetes Care, 2005, 28, 348-354.	8.6	48
219	Effects of Treatment Targets on Subsequent Cardiovascular Events in Chinese Patients With Type 2 Diabetes. Diabetes Care, 2007, 30, 953-959.	8.6	48
220	The Role of Adipocytokines and Neurohormonal Dysregulation in Metabolic Syndrome. Current Diabetes Reviews, 2006, 2, 397-407.	1.3	48
221	Comprehensive risk assessments of diabetic patients from seven Asian countries: The Joint Asia Diabetes Evaluation (JADE) program*. Journal of Diabetes, 2011, 3, 109-118.	1.8	47
222	Addressing different biases in analysing drug use on cancer risk in diabetes in non linical trial settingsâ€"what, why and how?. Diabetes, Obesity and Metabolism, 2012, 14, 579-585.	4.4	47
223	Using a multi-staged strategy based on machine learning and mathematical modeling to predict genotype-phenotype risk patterns in diabetic kidney disease: a prospective case–control cohort analysis. BMC Nephrology, 2013, 14, 162.	1.8	47
224	Sirt3 Deficiency Increased the Vulnerability of Pancreatic Beta Cells to Oxidative Stress-Induced Dysfunction. Antioxidants and Redox Signaling, 2017, 27, 962-976.	5.4	47
225	A Low-Frequency Inactivating <i>AKT2</i> Variant Enriched in the Finnish Population Is Associated With Fasting Insulin Levels and Type 2 Diabetes Risk. Diabetes, 2017, 66, 2019-2032.	0.6	47
226	The impact of maternal gestational weight gain on cardiometabolic risk factors in children. Diabetologia, 2018, 61, 2539-2548.	6.3	47
227	Diabetes in Hong Kong Chinese: evidence for familial clustering and parental effects. Diabetes Care, 2000, 23, 1365-1368.	8.6	46
228	Relationships between the TaqI polymorphism of the dopamine D2 receptor and blood pressure in hyperglycaemic and normoglycaemic Chinese subjects. Clinical Endocrinology, 2001, 55, 605-611.	2.4	46
229	Optimal control allocation in vehicle dynamics control for rollover mitigation. , 2008, , .		46
230	Association between Obesity and Atopy in Chinese Schoolchildren. International Archives of Allergy and Immunology, 2009, 149, 133-140.	2.1	46
231	Association of genetic variants of NOS1AP with type 2 diabetes in a Chinese population. Diabetologia, 2010, 53, 290-298.	6.3	46
232	From design to implementation - The Joint Asia Diabetes Evaluation (JADE) program: A descriptive report of an electronic web-based diabetes management program. BMC Medical Informatics and Decision Making, 2010, 10, 26.	3.0	46
233	Early Identification of Type 2 Diabetes. Diabetes Care, 2011, 34, 244-246.	8.6	46
234	Reducing global diabetes burden by implementing solutions and identifying gaps: a Lancet Commission. Lancet, The, 2016, 387, 1494-1495.	13.7	46

#	Article	IF	Citations
235	Accuracy, precision and user-acceptability of self blood glucose monitoring machines. Diabetes Research and Clinical Practice, 1997, 36, 91-104.	2.8	45
236	The Joint Asia Diabetes Evaluation (JADE) Program: a webâ€based program to translate evidence to clinical practice in Type 2 diabetes. Diabetic Medicine, 2009, 26, 693-699.	2.3	45
237	Glucoseâ€dependent insulinotropic peptide impairs insulin signaling <i>via</i> inducing adipocyte inflammation in glucoseâ€dependent insulinotropic peptide receptorâ€overexpressing adipocytes. FASEB Journal, 2012, 26, 2383-2393.	0.5	45
238	Fatal adult Henochâ€Sch×nlein purpura due to small intestinal infarction. Journal of Internal Medicine, 1992, 232, 181-184.	6.0	44
239	Association between adherence to statin therapy and lipid control in Hong Kong Chinese patients at high risk of coronary heart disease. British Journal of Clinical Pharmacology, 2004, 58, 528-535.	2.4	42
240	Drug-Induced Endocrine and Metabolic Disorders. Drug Safety, 2007, 30, 215-245.	3.2	42
241	Low LDL Cholesterol, Albuminuria, and Statins for the Risk of Cancer in Type 2 Diabetes: The Hong Kong Diabetes Registry. Diabetes Care, 2009, 32, 1826-1832.	8.6	42
242	Amyloid oligomers in diabetic and nondiabetic human pancreas. Translational Research, 2009, 153, 24-32.	5.0	42
243	Renal Outcomes in the EXenatide Study of Cardiovascular Event Lowering (EXSCEL). Diabetes, 2018, 67, .	0.6	42
244	Obesity, independent of insulin resistance, is a major determinant of blood pressure in normoglycemic Hong Kong Chinese. Metabolism: Clinical and Experimental, 2000, 49, 1523-1528.	3.4	41
245	PDZ-domain containing-2 (PDZD2) is a novel factor that affects the growth and differentiation of human fetal pancreatic progenitor cells. International Journal of Biochemistry and Cell Biology, 2008, 40, 789-803.	2.8	41
246	Use of anti-diabetic drugs and glycaemic control in type 2 diabetes—The Hong Kong Diabetes Registry. Diabetes Research and Clinical Practice, 2008, 82, 346-352.	2.8	41
247	Glucose Intolerance and Other Cardiovascular Risk Factors in Chinese Women with a History of Gestational Diabetes Mellitus. Australian and New Zealand Journal of Obstetrics and Gynaecology, 1999, 39, 478-483.	1.0	40
248	Diabetes and cancer: the mechanistic implications of epidemiological analyses from the Hong Kong Diabetes Registry. Diabetes/Metabolism Research and Reviews, 2012, 28, 379-387.	4.0	40
249	Middle Cerebral Artery Stenosis in Type II Diabetic Chinese Patients Is Associated with Conventional Risk Factors but Not with Polymorphisms of the Renin-Angiotensin System Genes. Cerebrovascular Diseases, 2003, 16, 217-223.	1.7	39
250	Smoking and diabetes in Chinese men. Postgraduate Medical Journal, 2001, 77, 240-243.	1.8	38
251	Assessment of glomerular filtration rate in addition to albuminuria is important in managing type II diabetes. Kidney International, 2006, 69, 383-387.	<b>5.2</b>	38
252	Olmesartan Reducing Incidence of Endstage Renal Disease in Diabetic Nephropathy Trial (ORIENT): Rationale and Study Design. Hypertension Research, 2006, 29, 703-709.	2.7	38

#	Article	IF	CITATIONS
253	Interactive effect of retinopathy and macroalbuminuria on all-cause mortality, cardiovascular and renal end points in Chinese patients with TypeÂ2 diabetes mellitus. Diabetic Medicine, 2007, 24, 741-746.	2.3	38
254	Predicting values of lipids and white blood cell count for all-site cancer in type 2 diabetes. Endocrine-Related Cancer, 2008, 15, 597-607.	3.1	38
255	Association between sleep architecture and glucose tolerance in children and adolescents å"¿ç«¥åŠé³å°'å¹´çš"çţço of Diabetes, 2015, 7, 10-15.	e结构ä 1.8	,Žç³–è€é‡ <mark>ç</mark> š
256	Progression of glucose intolerance and cardiometabolic risk factors over a decade in Chinese women with polycystic ovary syndrome: A case-control study. PLoS Medicine, 2019, 16, e1002953.	8.4	38
257	Triglyceride predicts cardiovascular mortality and its relationship with glycaemia and obesity in Chinese type 2 diabetic patients. Diabetes/Metabolism Research and Reviews, 2005, 21, 183-188.	4.0	37
258	Variation Within the Gene Encoding the Upstream Stimulatory Factor 1 Does Not Influence Susceptibility to Type 2 Diabetes in Samples From Populations With Replicated Evidence of Linkage to Chromosome 1q. Diabetes, 2006, 55, 2541-2548.	0.6	37
259	Impacts of chronic kidney disease and albuminuria on associations between coronary heart disease and its traditional risk factors in type 2 diabetic patients – the Hong Kong diabetes registry. Cardiovascular Diabetology, 2007, 6, 37.	6.8	37
260	Ethnic differences in the relationships of anthropometric measures to metabolic risk factors in Asian patients at risk of atherothrombosis. Metabolism: Clinical and Experimental, 2010, 59, 400-408.	3.4	37
261	Association between Physical Activity and Cardiovascular Risk in Chinese Youth Independent of Age and Pubertal Stage. BMC Public Health, 2010, 10, 303.	2.9	37
262	Shortened Leukocyte Telomere Length Is Associated With Glycemic Progression in Type 2 Diabetes: A Prospective and Mendelian Randomization Analysis. Diabetes Care, 2022, 45, 701-709.	8.6	37
263	Trends in all-cause mortality among people with diagnosed diabetes in high-income settings: a multicountry analysis of aggregate data. Lancet Diabetes and Endocrinology,the, 2022, 10, 112-119.	11.4	37
264	Relation between blood pressure and serum concentration of ouabain-like substance in non-insulin-dependent diabetes mellitus. Lancet, The, 1998, 351, 266.	13.7	36
265	Sonographic Measurement of Mesenteric Fat Predicts Presence of Fatty Liver among Subjects with Polycystic Ovary Syndrome. Journal of Clinical Endocrinology and Metabolism, 2011, 96, 799-807.	3.6	36
266	Association between tumour necrosis factor-α G-308A polymorphism and risk of nephropathy in obese Chinese type 2 diabetic patients. Nephrology Dialysis Transplantation, 2005, 20, 2733-2738.	0.7	35
267	Association of lipoprotein lipase S447X, apolipoprotein E exon 4, and apoC3 â^'455T>C polymorphisms on the susceptibility to diabetic nephropathy. Clinical Genetics, 2006, 70, 20-28.	2.0	35
268	Additive Interaction Between the Renin-Angiotensin System and Lipid Metabolism for Cancer in Type 2 Diabetes. Diabetes, 2009, 58, 1518-1525.	0.6	35
269	Lipid control and use of lipid-regulating drugs for prevention of cardiovascular events in Chinese type 2 diabetic patients: a prospective cohort study. Cardiovascular Diabetology, 2010, 9, 77.	6.8	35
270	The Insulin Resistance Syndrome: Mechanisms of Clustering of Cardiovascular Risk. Seminars in Vascular Medicine, 2002, 2, 045-058.	2.1	34

#	Article	IF	Citations
271	Cardiac hypertrophy and remodeling in relation to ACE and angiotensinogen genes genotypes in Chinese dialysis patients. Kidney International, 2003, 63, 1899-1907.	5 <b>.</b> 2	34
272	Metabolic syndrome by the international diabetes federation definition in Hong Kong Chinese. Diabetes Research and Clinical Practice, 2006, 73, 58-64.	2.8	34
273	Common Variation in the LMNA Gene (Encoding Lamin A/C) and Type 2 Diabetes: Association Analyses in 9,518 Subjects. Diabetes, 2007, 56, 879-883.	0.6	34
274	Progression to impaired glucose regulation, diabetes and metabolic syndrome in Chinese women with a past history of gestational diabetes. Diabetes/Metabolism Research and Reviews, 2007, 23, 485-489.	4.0	34
275	Low plasma adiponectin level, white blood cell count and Helicobacter pylori titre independently predict abnormal pancreatic $\hat{l}^2$ -cell function. Diabetes Research and Clinical Practice, 2009, 86, 89-95.	2.8	34
276	Adherence to Oral Hypoglycemic Agents in 26 782 Chinese Patients: A Cohort Study. Journal of Clinical Pharmacology, 2011, 51, 1474-1482.	2.0	34
277	The Asian diabetes phenotypes: Challenges and opportunities. Diabetes Research and Clinical Practice, 2014, 105, 135-139.	2.8	34
278	Use of Net Reclassification Improvement (NRI) Method Confirms The Utility of Combined Genetic Risk Score to Predict Type 2 Diabetes. PLoS ONE, 2013, 8, e83093.	2.5	34
279	Glycaemic control in type 2 diabetes: the impact of body weight, beta-cell function and patient education. QJM - Monthly Journal of the Association of Physicians, 2000, 93, 183-190.	0.5	33
280	Diabetes mellitusâ€"a special medical challenge from a Chinese perspective. Diabetes Research and Clinical Practice, 2001, 54, 19-27.	2.8	33
281	Peripheral vascular disease in Type 2 diabetic Chinese patients: associations with metabolic indices, concomitant vascular disease and genetic factors. Diabetic Medicine, 2003, 20, 988-995.	2.3	33
282	Effect of Angiotensin-Converting Enzyme Inhibition on Survival in 3773 Chinese Type 2 Diabetic Patients. Hypertension, 2004, 44, 294-299.	2.7	33
283	Meta-Analysis of Genome-Wide Linkage Studies of Quantitative Lipid Traits in Families Ascertained for Type 2 Diabetes. Diabetes, 2007, 56, 890-896.	0.6	33
284	Worsening trend of central obesity despite stable or declining body mass index in Hong Kong Chinese between 1996 and 2005. European Journal of Clinical Nutrition, 2010, 64, 549-552.	2.9	33
285	A simple risk score to identify Southern Chinese at high risk for diabetes. Diabetic Medicine, 2010, 27, 644-649.	2.3	33
286	Peri-operative hyperglycemia: a consideration for general surgery?. American Journal of Surgery, 2010, 199, 240-248.	1.8	33
287	Rationale, Design, and Baseline Characteristics of ARTS-DN: A Randomized Study to Assess the Safety and Efficacy of Finerenone in Patients with Type 2 Diabetes Mellitus and a Clinical Diagnosis of Diabetic Nephropathy. American Journal of Nephrology, 2014, 40, 572-581.	3.1	33
288	Quality of care in patients with diabetic kidney disease in Asia: The Joint Asia Diabetes Evaluation ( <scp>JADE</scp> ) Registry. Diabetic Medicine, 2016, 33, 1230-1239.	2.3	33

#	Article	IF	CITATIONS
289	Adverse Reactions to Drugs as a Cause of Admissions to a General Teaching Hospital in Hong Kong. Drug Safety, 1992, 7, 235-240.	3.2	32
290	Implication of fibrate therapy for homocysteine. Lancet, The, 1999, 354, 1208-1209.	13.7	32
291	The lipoprotein lipase gene HindIII polymorphism is associated with lipid levels in early-onset type 2 diabetic patients. Metabolism: Clinical and Experimental, 2003, 52, 338-343.	3.4	32
292	Aberrant Expression of Soluble Co-stimulatory Molecules and Adhesion Molecules in Type 2 Diabetic Patients with Nephropathy. Journal of Clinical Immunology, 2008, 28, 36-43.	3.8	32
293	The problem of obesity among adolescents in Hong Kong: a comparison using various diagnostic criteria. BMC Pediatrics, 2008, 8, 10.	1.7	32
294	Aldose Reductase Genotypes and Cardiorenal Complications. Diabetes Care, 2008, 31, 2148-2153.	8.6	32
295	High risk for cardiovascular disease in Chinese type 2 diabetic patients with major depression—A 7-year prospective analysis of the Hong Kong DiabetesRegistry. Journal of Affective Disorders, 2013, 149, 129-135.	4.1	32
296	Acarbose improves glycemic control in insulin-treated Asian type 2 diabetic patients: Results from a multinational, placebo-controlled study. Diabetes Research and Clinical Practice, 2003, 60, 111-118.	2.8	31
297	Activating Transcription Factor 6 (ATF6) Sequence Polymorphisms in Type 2 Diabetes and Pre-Diabetic Traits. Diabetes, 2007, 56, 856-862.	0.6	31
298	Reference values for serum levels of insulin-like growth factor (IGF-1) and IGF-binding protein 3 (IGFBP-3) and their ratio in Chinese adolescents. Clinical Biochemistry, 2007, 40, 1093-1099.	1.9	31
299	White rice vinegar improves pancreatic beta-cell function and fatty liver in streptozotocin-induced diabetic rats. Acta Diabetologica, 2012, 49, 185-191.	2.5	31
300	Sequence data and association statistics from 12,940 type 2 diabetes cases and controls. Scientific Data, 2017, 4, 170179.	5.3	31
301	Shortened Relative Leukocyte Telomere Length Is Associated With Prevalent and Incident Cardiovascular Complications in Type 2 Diabetes: Analysis From the Hong Kong Diabetes Register. Diabetes Care, 2020, 43, 2257-2265.	8.6	31
302	Obesity, clinical, and genetic predictors for glycemic progression in Chinese patients with type 2 diabetes: A cohort study using the Hong Kong Diabetes Register and Hong Kong Diabetes Biobank. PLoS Medicine, 2020, 17, e1003209.	8.4	31
303	Machine learning risk score for prediction of gestational diabetes in early pregnancy in Tianjin, China. Diabetes/Metabolism Research and Reviews, 2021, 37, e3397.	4.0	31
304	Mitochondrial DNA A3243G mutation in patients with early- or late-onset type 2 diabetes mellitus in Hong Kong Chinese. Clinical Endocrinology, 2000, 52, 557-564.	2.4	30
305	Factors Predicting the Age When Type 2 Diabetes Is Diagnosed in Hong Kong Chinese Subjects. Diabetes Care, 2001, 24, 646-649.	8.6	30
306	Linkage Disequilibrium Mapping of the Replicated Type 2 Diabetes Linkage Signal on Chromosome 1q. Diabetes, 2009, 58, 1704-1709.	0.6	30

#	Article	IF	Citations
307	Association of the PPARG Pro12Ala polymorphism with type 2 diabetes and incident coronary heart disease in a Hong Kong Chinese population. Diabetes Research and Clinical Practice, 2012, 97, 483-491.	2.8	30
308	Associations of uric acid and gammaâ€glutamyltransferase ( <scp>GGT</scp> ) with obesity and components of metabolic syndrome in children and adolescents. Pediatric Obesity, 2013, 8, 351-357.	2.8	30
309	Use of SGLT-2 Inhibitors in Patients with Type 2 Diabetes Mellitus and Abdominal Obesity: An Asian Perspective and Expert Recommendations. Diabetes and Metabolism Journal, 2020, 44, 11.	4.7	30
310	Independent associations of alanine aminotransferase (ALT) levels with cardiovascular risk factor clustering in Chinese adolescents. Journal of Hepatology, 2008, 49, 115-122.	3.7	29
311	Incidence of childhood type 1 diabetes: a worrying trend. Nature Reviews Endocrinology, 2009, 5, 529-530.	9.6	29
312	The Combined Expression of $\langle i \rangle Pdx1 \langle i \rangle$ and $\langle i \rangle MafA \langle i \rangle$ with Either $\langle i \rangle Ngn3 \langle i \rangle$ or $\langle i \rangle NeuroD \langle i \rangle$ Improves the Differentiation Efficiency of Mouse Embryonic Stem Cells into Insulin-Producing Cells. Cell Transplantation, 2013, 22, 147-158.	2.5	29
313	Plasma Levels of Alanine Aminotransferase in the First Trimester Identify High Risk Chinese Women for Gestational Diabetes. Scientific Reports, 2016, 6, 27291.	3.3	29
314	Resource use associated with type 2 diabetes in Africa, the Middle East, South Asia, Eurasia and Turkey: results from the International Diabetes Management Practice Study (IDMPS). BMJ Open Diabetes Research and Care, 2017, 5, e000297.	2.8	29
315	Sex differences in the association between socioeconomic status and diabetes prevalence and incidence in China: cross-sectional and prospective studies of 0.5 million adults. Diabetologia, 2019, 62, 1420-1429.	6.3	29
316	Trends in Glucose-Lowering Drug Use, Glycemic Control, and Severe Hypoglycemia in Adults With Diabetes in Hong Kong, 2002–2016. Diabetes Care, 2020, 43, 2967-2974.	8.6	29
317	The Short Insulin Tolerance Test: Feasibility Study Using Venous Sampling. Diabetic Medicine, 1996, 13, 429-433.	2.3	28
318	Fasting Plasma Glucose as a Screening Test for Diabetes and Its Relationship With Cardiovascular Risk Factors in Hong Kong Chinese. Diabetes Care, 1997, 20, 170-172.	8.6	28
319	Increasing Severity of Cardiovascular Risk Factors With Increasing Middle Cerebral Artery Stenotic Involvement in Type 2 Diabetic Chinese Patients With Asymptomatic Cerebrovascular Disease. Diabetes Care, 2004, 27, 1121-1126.	8.6	28
320	Role of metformin in the initiation of pharmacotherapy for type 2 diabetes: An Asian-Pacific perspective. Diabetes Research and Clinical Practice, 2007, 75, 255-266.	2.8	28
321	US Assessment of Medial Arterial Calcification: A Sensitive Marker of Diabetes-related Microvascular and Macrovascular Complications. Radiology, 2012, 265, 294-302.	7.3	28
322	Chronic Kidney Disease and Associated Cardiovascular Risk Factors in Chinese with Type 2 Diabetes. Diabetes and Metabolism Journal, 2012, 36, 433.	4.7	28
323	Self-Reported Body Weight and Height: An Assessment Tool for Identifying Children with Overweight/Obesity Status and Cardiometabolic Risk Factors Clustering. Maternal and Child Health Journal, 2013, 17, 282-291.	1.5	28
324	Hypoglycaemia, chronic kidney disease and death in type 2 diabetes: the Hong Kong diabetes registry. BMC Endocrine Disorders, 2014, 14, 48.	2,2	28

#	Article	IF	Citations
325	Impact of diabetes education and self-management on the quality of care for people with type 1 diabetes mellitus in the Middle East (the International Diabetes Mellitus Practices Study, IDMPS). Diabetes Research and Clinical Practice, 2019, 147, 29-36.	2.8	28
326	In vivo treatment with glucagon-like peptide $1$ promotes the graft function of fetal islet-like cell clusters in transplanted mice. International Journal of Biochemistry and Cell Biology, 2006, 38, 951-960.	2.8	27
327	Diabetic nephropathyâ€"What are the unmet needs?. Diabetes Research and Clinical Practice, 2008, 82, S15-S20.	2.8	27
328	Additive Interaction of Hyperglycemia and Albuminuria on Risk of Ischemic Stroke in Type 2 Diabetes. Diabetes Care, 2008, 31, 2294-2300.	8.6	27
329	A New Tool to Detect Kidney Disease in Chinese Type 2 Diabetes Patients: Comparison of EZSCAN with Standard Screening Methods. Diabetes Technology and Therapeutics, 2011, 13, 937-943.	4.4	27
330	Healing effect of a twoâ€herb recipe ( <scp>NF3</scp> ) on foot ulcers in <scp>C</scp> hinese patients with diabetes: A randomized doubleâ€blind placeboâ€controlled study (包å«ä,¤§è‰ææ¬ä¸è•çš"é…æ−¹ï¼^NF323-334.	F3ï¥⁄48‱å⁻¹	<sup>1</sup> ä¸å <b>27⁄</b> 2ç³—å°¿ç
331	Unfolded Protein Response Is Required for the Definitive Endodermal Specification of Mouse Embryonic Stem Cells via Smad2 and β-Catenin Signaling. Journal of Biological Chemistry, 2014, 289, 26290-26301.	3.4	27
332	MicroRNA and Diabetic Complications: A Clinical Perspective. Antioxidants and Redox Signaling, 2018, 29, 1041-1063.	5.4	27
333	Age at diagnosis, glycemic trajectories, and responses to oral glucose-lowering drugs in type 2 diabetes in Hong Kong: A population-based observational study. PLoS Medicine, 2020, 17, e1003316.	8.4	27
334	Apolipoprotein M Gene (APOM) Polymorphism Modifies Metabolic and Disease Traits in Type 2 Diabetes. PLoS ONE, 2011, 6, e17324.	2.5	27
335	An insulin receptor gene polymorphism is associated with diastolic blood pressure in Chinese subjects with components of the metabolic syndrome. American Journal of Hypertension, 2000, 13, 745-752.	2.0	26
336	Ethnic differences in the linkage disequilibrium and distribution of single-nucleotide polymorphisms in 35 candidate genes for cardiovascular diseases. Genomics, 2004, 83, 559-565.	2.9	26
337	Prediction of cardiovascular and total mortality in Chinese type 2 diabetic patients by the WHO definition for the metabolic syndrome. Diabetes, Obesity and Metabolism, 2006, 8, 94-104.	4.4	26
338	Pancreatic islet $\hat{l}^2$ -cell deficit and glucose intolerance in rats with uninephrectomy. Cellular and Molecular Life Sciences, 2007, 64, 3119-3128.	5 <b>.</b> 4	26
339	Exendin-4 Improves Blood Glucose Control in Both Young and Aging Normal Non-Diabetic Mice, Possible Contribution of Beta Cell Independent Effects. PLoS ONE, 2011, 6, e20443.	2.5	26
340	Cardiometabolic Risk in Chinese Women with Prior Gestational Diabetes: A 15-Year Follow-Up Study. Gynecologic and Obstetric Investigation, 2012, 73, 168-176.	1.6	26
341	Effects of dual blockade of the renin–angiotensin system on renal and cardiovascular outcomes in type 2 diabetes with overt nephropathy and hypertension in the ORIENT: a post-hoc analysis (ORIENT–Hypertension). Hypertension Research, 2013, 36, 1051-1059.	2.7	26
342	EDN1 Lys198Asn is associated with diabetic retinopathy in type 2 diabetes. Molecular Vision, 2008, 14, 1698-704.	1.1	26

#	Article	IF	Citations
343	Effects of protocol-driven care versus usual outpatient clinic care on survival rates in patients with type 2 diabetes. American Journal of Managed Care, 2003, 9, 606-15.	1.1	26
344	Diabetic Microangiopathic Complications in Young Chinese Diabetic Patients. Journal of Diabetes and Its Complications, 1999, 13, 300-306.	2.3	25
345	Contribution of gene polymorphisms in the renin-angiotensin system to macroangiopathy in patients with diabetic nephropathy. American Journal of Kidney Diseases, 2001, 38, 9-17.	1.9	25
346	Independent predictive roles of eotaxin Ala23Thr, paraoxonaseâ€f2 Ser311Cys and β <sub>3</sub> â€adrenergic receptor Trp64Arg polymorphisms on cardiac disease in Type 2 Diabetesâ€"an 8â€year prospective cohort analysis of 1297 patients. Diabetic Medicine, 2010, 27, 376-383.	2.3	25
347	The association between in utero hyperinsulinemia and adolescent arterial stiffness. Diabetes Research and Clinical Practice, 2012, 95, 169-175.	2.8	25
348	Enhancers and attenuators of risk associations of chronic hepatitis B virus infection with hepatocellular carcinoma in type 2 diabetes. Endocrine-Related Cancer, 2013, 20, 161-171.	3.1	25
349	Nonâ€linear relationship between birthweight and cardiometabolic risk factors in Chinese adolescents and adults. Diabetic Medicine, 2015, 32, 220-225.	2.3	25
350	Graphene oxide $\hat{a}\in$ Polyvinyl alcohol nanocomposite based electrode material for supercapacitors. Journal of Power Sources, 2016, 321, 102-105.	7.8	25
351	Cardiovascular outcomes trials in type 2 diabetes: Time to include young adults. Diabetes, Obesity and Metabolism, 2020, 22, 3-5.	4.4	25
352	The Clinical Utility of SUDOSCAN in Chronic Kidney Disease in Chinese Patients with Type 2 Diabetes. PLoS ONE, 2015, 10, e0134981.	2.5	25
353	Associations between Food Variety and Body Fatness in Hong Kong Chinese Adults. Journal of the American College of Nutrition, 2004, 23, 404-413.	1.8	24
354	Vascular defect beyond the endothelium in type II diabetic patients with overt nephropathy and moderate renal insufficiency. Kidney International, 2006, 70, 711-716.	5.2	24
355	Bayesian analysis of structural equation models with multinomial variables and an application to type 2 diabetic nephropathy. Statistics in Medicine, 2007, 26, 2348-2369.	1.6	24
356	Eczema phenotypes are associated with multiple vitamin <scp>D</scp> pathway genes in <scp>C</scp> hinese children. Allergy: European Journal of Allergy and Clinical Immunology, 2014, 69, 118-124.	5.7	24
357	Additive effects of blood glucose lowering drugs, statins and renin-angiotensin system blockers on all-site cancer risk in patients with type 2 diabetes. BMC Medicine, 2014, 12, 76.	<b>5.</b> 5	24
358	Selenium-enriched Spirulina protects INS-1E pancreatic beta cells from human islet amyloid polypeptide-induced apoptosis through suppression of ROS-mediated mitochondrial dysfunction and PI3/AKT pathway. European Journal of Nutrition, 2015, 54, 509-522.	3.9	24
359	A multicentre demonstration project to evaluate the effectiveness and acceptability of the webâ€based Joint Asia Diabetes Evaluation ( <scp>JADE</scp> ) programme with or without nurse support in Chinese patients with Type 2 diabetes. Diabetic Medicine, 2017, 34, 440-450.	2.3	24
360	Diabetes-Related Complications and Mortality in Patients With Young-Onset Latent Autoimmune Diabetes: A 14-Year Analysis of the Prospective Hong Kong Diabetes Register. Diabetes Care, 2019, 42, 1042-1050.	8.6	24

#	Article	IF	Citations
361	Association of technologically assisted integrated care with clinical outcomes in type 2 diabetes in Hong Kong using the prospective JADE Program: A retrospective cohort analysis. PLoS Medicine, 2020, 17, e1003367.	8.4	24
362	Trends in diabetes-related complications in Hong Kong, 2001–2016: a retrospective cohort study. Cardiovascular Diabetology, 2020, 19, 60.	6.8	24
363	Temporal trends in rates of infection-related hospitalisations in Hong Kong people with and without diabetes, 2001–2016: a retrospective study. Diabetologia, 2021, 64, 109-118.	6.3	24
364	Outcomes of screening for diabetes in high-risk Hong Kong Chinese subjects. Diabetes Care, 2000, 23, 1290-1294.	8.6	23
365	Association of Glomerulopathy With the 5'-End Polymorphism of the Aldose Reductase Gene and Renal Insufficiency in Type 2 Diabetic Patients. Diabetes, 2004, 53, 2984-2991.	0.6	23
366	Albuminuria is a marker of increasing intracranial and extracranial vascular involvement in Type 2 diabetic Chinese patients. Diabetologia, 2004, 47, 1528-1534.	6.3	23
367	Effectiveness of a lifestyle modification programme in weight maintenance in obese subjects after cessation of treatment with Orlistat. Journal of Evaluation in Clinical Practice, 2007, 13, 070728052316007-???.	1.8	23
368	The NCEPâ€ATPIII but not the IDF criteria for the metabolic syndrome identify Type 2 diabetic patients at increased risk of chronic kidney disease. Diabetic Medicine, 2008, 25, 1419-1425.	2.3	23
369	Phenotype–genotype interactions on renal function in type 2 diabetes: an analysis using structural equation modelling. Diabetologia, 2009, 52, 1543-1553.	6.3	23
370	Lack of benefits for prevention of cardiovascular disease with aspirin therapy in type 2 diabetic patients - a longitudinal observational study. Cardiovascular Diabetology, 2009, 8, 57.	6.8	23
371	Evaluation of erectile dysfunction and associated cardiovascular risk using structured questionnaires in Chinese type 2 diabetic men. Journal of Developmental and Physical Disabilities, 2010, 33, 853-860.	3.6	23
372	Reduction and residual proteinuria are therapeutic targets in type 2 diabetes with overt nephropathy: a post hoc analysis (ORIENT-proteinuria). Nephrology Dialysis Transplantation, 2013, 28, 2526-2534.	0.7	23
373	A randomized controlled trial to investigate the impact of a low glycemic index (GI) diet on body mass index in obese adolescents. BMC Public Health, 2014, 14, 180.	2.9	23
374	Drug–subphenotype interactions for cancer in type 2 diabetes mellitus. Nature Reviews Endocrinology, 2015, 11, 372-379.	9.6	23
375	Detecting people at high risk of type 2 diabetes- How do we find them and who should be treated?. Best Practice and Research in Clinical Endocrinology and Metabolism, 2016, 30, 345-355.	4.7	23
376	Gender, diabetes education, and psychosocial factors are associated with persistent poor glycemic control in patients with type 2 diabetes in the <scp>J</scp> oint <scp>A</scp> sia <scp>D</scp> iabetes <scp>E</scp> valuation ( <scp>JADE</scp> ) program. Journal of Diabetes, 2016, 8, 109-119.	1.8	23
377	Glycaemia control and the risk of hospitalisation for infection in patients with type 2 diabetes: Hong Kong Diabetes Registry. Diabetes/Metabolism Research and Reviews, 2017, 33, e2923.	4.0	23
378	Development and validation of an early pregnancy risk score for the prediction of gestational diabetes mellitus in Chinese pregnant women. BMJ Open Diabetes Research and Care, 2020, 8, e000909.	2.8	23

#	Article	IF	CITATIONS
379	SERUM SOLUBLE INTERLEUKIN 2 RECEPTOR IN HYPERTHYROID Graves' DISEASE AND EFFECT OF CARBIMAZOLE THERAPY. Clinical Endocrinology, 1990, 33, 317-321.	2.4	22
380	Association of the POU class $\hat{a} \in f2$ homeobox $\hat{a} \in f1$ gene (POU2F1) with susceptibility to Type $\hat{a} \in f2$ diabetes in Chinese populations. Diabetic Medicine, 2010, 27, 1443-1449.	2.3	22
381	Delivery of integrated diabetes care using logistics and information technology – The Joint Asia Diabetes Evaluation (JADE) program. Diabetes Research and Clinical Practice, 2014, 106, S295-S304.	2.8	22
382	Genetic and clinical variables identify predictors forÂchronic kidney disease in type 2 diabetes. Kidney International, 2016, 89, 411-420.	5.2	22
383	Passive smoking increased risk of gestational diabetes mellitus independently and synergistically with prepregnancy obesity in Tianjin, China. Diabetes/Metabolism Research and Reviews, 2017, 33, e2861.	4.0	22
384	Roles of insulin resistance and beta cell dysfunction in macrosomia among Chinese women with gestational diabetes mellitus. Primary Care Diabetes, 2018, 12, 565-573.	1.8	22
385	Use of sodiumâ€glucose coâ€transporterâ€2 inhibitors in patients with type 2 diabetes mellitus and multiple cardiovascular risk factors: An Asian perspective and expert recommendations. Diabetes, Obesity and Metabolism, 2019, 21, 2354-2367.	4.4	22
386	Young age at diabetes diagnosis amplifies the effect of diabetes duration on risk of chronic kidney disease: a prospective cohort study. Diabetologia, 2021, 64, 1990-2000.	6.3	22
387	Histiocytosis X involving the thyroid and hypothalamus Postgraduate Medical Journal, 1991, 67, 680-682.	1.8	21
388	Herb-induced aconitine poisoning. Lancet, The, 1993, 341, 370-371.	13.7	21
389	The association between dyslipidaemia and obesity in Chinese men after adjustment for insulin resistance. Atherosclerosis, 1998, 138, 153-161.	0.8	21
390	Cutoff values for central obesity in Chinese based on mesenteric fat thickness. Clinical Nutrition, 2009, 28, 679-683.	5.0	21
391	Low triglyceride and nonuse of statins is associated with cancer in type 2 diabetes mellitus. Cancer, 2011, 117, 862-871.	4.1	21
392	Validation of Methods to Control for Immortal Time Bias in a Pharmacoepidemiologic Analysis of Renin^ ^ndash;Angiotensin System Inhibitors in Type 2 Diabetes. Journal of Epidemiology, 2014, 24, 267-273.	2.4	21
393	Prevention and Care Programs Addressing the Growing Prevalence of Diabetes in China. Current Diabetes Reports, 2016, 16, 130.	4.2	21
394	Transancestral fine-mapping of four type 2 diabetes susceptibility loci highlights potential causal regulatory mechanisms. Human Molecular Genetics, 2016, 25, 2070-2081.	2.9	21
395	Interactions between general and central obesity in predicting gestational diabetes mellitus in Chinese pregnant women: A prospective populationâ€based study in Tianjin, China. Journal of Diabetes, 2018, 10, 59-67.	1.8	21
396	Glycaemic responses in Asian and non-Asian people with type 2 diabetes initiating insulin glargine 100†units/mL: A patient-level pooled analysis of 16 randomised controlled trials. Diabetes Research and Clinical Practice, 2018, 135, 199-205.	2.8	21

#	Article	IF	CITATIONS
397	Depressive Symptoms, Co-Morbidities, and Glycemic Control in Hong Kong Chinese Elderly Patients With Type 2 Diabetes Mellitus. Frontiers in Endocrinology, 2018, 9, 261.	3.5	21
398	Middle Cerebral Artery Stenosis Increased the Risk of Vascular Disease Mortality among Type 2 Diabetic Patients. Cerebrovascular Diseases, 2008, 25, 261-267.	1.7	20
399	Effect of interactions between C peptide levels and insulin treatment on clinical outcomes among patients with type 2 diabetes mellitus. Cmaj, 2009, 180, 919-926.	2.0	20
400	Apolipoprotein E polymorphism and expression in type 2 diabetic patients with nephropathy: clinicopathological correlation. Nephrology Dialysis Transplantation, 2009, 24, 1889-1895.	0.7	20
401	Role of lowâ€glycemic index diet in management of childhood obesity. Obesity Reviews, 2011, 12, 492-498.	6.5	20
402	Association between KCNQ1 genetic variants and obesity in Chinese patients with type 2 diabetes. Diabetologia, 2012, 55, 2655-2659.	6.3	20
403	Interactome-transcriptome analysis discovers signatures complementary to GWAS Loci of Type 2 Diabetes. Scientific Reports, 2016, 6, 35228.	3.3	20
404	Type 2 Diabetes Promotes Cell Centrosome Amplification via AKT-ROS-Dependent Signalling of ROCK1 and 14-3-3Ïf. Cellular Physiology and Biochemistry, 2018, 47, 356-367.	1.6	20
405	Circulating branchedâ€chain amino acids and incident heart failure in type 2 diabetes: The Hong Kong Diabetes Register. Diabetes/Metabolism Research and Reviews, 2020, 36, e3253.	4.0	20
406	Use of sodiumâ€glucose coâ€transporterâ€2 inhibitors in <scp>Asian</scp> patients with type 2 diabetes and kidney disease: An <scp>Asian</scp> perspective and expert recommendations. Diabetes, Obesity and Metabolism, 2021, 23, 299-317.	4.4	20
407	Data Resource Profile: The Hong Kong Diabetes Surveillance Database (HKDSD). International Journal of Epidemiology, 2022, 51, e9-e17.	1.9	20
408	A Case of Stevens-Johnson Syndrome, Cholestatic Hepatitis and Haemolytic Anaemia Associated with Use of Mefenamic Acid. Drug Safety, 1991, 6, 230-234.	3.2	19
409	Atrial natriuretic peptide and urinary dopamine output in non-insulin-dependent diabetes mellitus. Clinical Science, 1992, 83, 247-253.	4.3	19
410	Change of glycaemic status in Chinese subjects with impaired fasting glycaemia. Diabetic Medicine, 2001, 18, 745-748.	2.3	19
411	The efficacy and tolerability of fosinopril in Chinese type 2 diabetic patients with moderate renal insufficiency. Diabetes, Obesity and Metabolism, 2006, 8, 342-347.	4.4	19
412	Acute Renal Failure Following Oral Sodium Phosphate Bowel Preparation in Diabetes. Diabetes Care, 2007, 30, 182-183.	8.6	19
413	Risk associations of obesity with sugar-sweetened beverages and lifestyle factors in Chinese: the $\hat{a} \in \mathbb{R}$ Better Health for Better Hong Kong $\hat{a} \in \mathbb{R}$ health promotion campaign. European Journal of Clinical Nutrition, 2010, 64, 1386-1392.	2.9	19
414	Association of statin use and development of renal dysfunction in type 2 diabetesâ€"The Hong Kong Diabetes Registry. Diabetes Research and Clinical Practice, 2010, 88, 227-233.	2.8	19

#	Article	IF	CITATIONS
415	Predictive role of polymorphisms in interleukin-5 receptor alpha-subunit, lipoprotein lipase, integrin A2 and nitric oxide synthase genes on ischemic stroke in type 2 diabetes—An 8-year prospective cohort analysis of 1327 Chinese patients. Atherosclerosis, 2011, 215, 130-135.	0.8	19
416	Maternal history of diabetes is associated with increased cardiometabolic risk in Chinese. Nutrition and Diabetes, 2014, 4, e112-e112.	3.2	19
417	Biologics and biosimilars: what, why and how?. ESMO Open, 2017, 2, e000180.	<b>4.</b> 5	19
418	Effect of Onceâ€Weekly Exenatide on Clinical Outcomes According to Baseline Risk in Patients With Type 2 Diabetes Mellitus: Insights From the EXSCEL Trial. Journal of the American Heart Association, 2018, 7, e009304.	3.7	19
419	Risk-factor profile, drug usage and cardiovascular events within a year in patients with and at high risk of atherothrombosis recruited from Asia as compared with those recruited from non-Asian regions: a substudy of the REduction of Atherothrombosis for Continued Health (REACH) registry. Heart Asia. 2011. 3. 93-8.	1.1	19
420	Validity of Glycated Hemoglobin in Screening and Diagnosing Type 2 Diabetes Mellitus in Chinese Subjects. Korean Journal of Internal Medicine, 2012, 27, 41.	1.7	19
421	Use of Antidiabetic and Antihypertensive Drugs in Hospital and Outpatient Settings in Hong Kong. Annals of Pharmacotherapy, 1996, 30, 232-237.	1.9	18
422	The effect of age on cardiovascular risk factors in Chinese women. International Journal of Cardiology, 1997, 61, 221-227.	1.7	18
423	Drug utilization in a hospital general medical outpatient clinic with particular reference to antihypertensive and antidiabetic drugs. Journal of Clinical Pharmacy and Therapeutics, 1998, 23, 287-294.	1.5	18
424	Sulphonylurea-induced hypoglycaemia in institutionalized elderly in Hong Kong. Diabetic Medicine, 2002, 19, 966-968.	2.3	18
425	Tumour necrosis factor-alpha promoter gene polymorphism at - 308 (genotype AA) in Chinese subjects with Type 2 diabetes. Diabetic Medicine, 2003, 20, 167-168.	2.3	18
426	Carbamazepine and false positive dexamethasone suppression tests for Cushing's syndrome. BMJ: British Medical Journal, 2005, 330, 299-300.	2.3	18
427	Renin angiotensin aldosterone system blockade and renal disease in patients with type 2 diabetes: a subanalysis of Japanese patients from the RENAAL study. Clinical and Experimental Nephrology, 2006, 10, 193-200.	1.6	18
428	Doubling over ten years of central obesity in Hong Kong Chinese working men. Chinese Medical Journal, 2007, 120, 1151-1154.	2.3	18
429	A randomized-controlled trial to investigate the effects of rivoglitazone, a novel PPAR gamma agonist on glucose-lipid control in type 2 diabetes. Diabetes, Obesity and Metabolism, 2011, 13, 806-813.	4.4	18
430	Hong Kong Chinese school children with elevated urine melamine levels: A prospective follow up study. BMC Public Health, 2011, 11, 354.	2.9	18
431	Microtubule network is required for insulin-induced signal transduction and actin remodeling. Molecular and Cellular Endocrinology, 2013, 365, 64-74.	3.2	18
432	Pregnancy outcomes of Chinese women with gestational diabetes mellitus defined by the IADPSG's but not by the 1999 WHO's criteria. Clinical Endocrinology, 2015, 83, 684-693.	2.4	18

#	Article	lF	CITATIONS
433	Sex Differences in Epidemiology and Risk Factors of Acute Coronary Syndrome in Chinese Patients with Type 2 Diabetes: A Long-Term Prospective Cohort Study. PLoS ONE, 2015, 10, e0122031.	2.5	18
434	Borderline ankle–brachial index is associated with increased prevalence of micro- and macrovascular complications in type 2 diabetes: A cross-sectional analysis of 12,772 patients from the Joint Asia Diabetes Evaluation Program. Diabetes and Vascular Disease Research, 2015, 12, 334-341.	2.0	18
435	Effects of blood pressure on renal and cardiovascular outcomes in Asian patients with type 2 diabetes and overt nephropathy: a <i>post hoc</i> analysis (ORIENT-blood pressure). Nephrology Dialysis Transplantation, 2016, 31, 447-454.	0.7	18
436	Development of genome-wide polygenic risk scores for lipid traits and clinical applications for dyslipidemia, subclinical atherosclerosis, and diabetes cardiovascular complications among East Asians. Genome Medicine, 2021, 13, 29.	8.2	18
437	Severe salicylate poisoning associated with the intake of Chinese medicinal oil ( Red Flower Oil'). Australian and New Zealand Journal of Medicine, 1995, 25, 57-57.	0.5	17
438	Risk factors for cataract in Chinese patients with type 2 diabetes: evidence for the influence of the aldose reductase gene. Clinical Genetics, 2002, 59, 356-359.	2.0	17
439	The renoprotective effects of structured care in a clinical trial setting in type 2 diabetic patients with nephropathy. Nephrology Dialysis Transplantation, 2004, 19, 2519-2525.	0.7	17
440	Effects of gender, Helicobacter pylori and hepatitis B virus serology status on cardiovascular and renal complications in Chinese type 2 diabetic patients with overt nephropathy. Diabetes, Obesity and Metabolism, 2004, 6, 223-230.	4.4	17
441	Modified end-stage renal disease risk score for Chinese type 2 diabetic patients—the Hong Kong Diabetes Registry. Diabetologia, 2007, 50, 1348-1350.	6.3	17
442	Higher Islet Amyloid Load in Men Than in Women With Type 2 Diabetes Mellitus. Pancreas, 2008, 37, e68-e73.	1.1	17
443	Expression and subcellular targeting of human insulin-like growth factor binding protein-3 in transgenic tobacco plants. Transgenic Research, 2009, 18, 943-951.	2.4	17
444	Cold-vapour atomic absorption spectrometry underestimates total mercury in blood and urine compared to inductively-coupled plasma mass spectrometry: an important factor for determining mercury reference intervals. Pathology, 2009, 41, 467-472.	0.6	17
445	Diabetic dyslipidaemia in Asian populations in the Western Pacific Region: What we know and don't know. Diabetes Research and Clinical Practice, 2011, 94, 1-13.	2.8	17
446	Serum concentrations of insulin-like growth factor-I, insulin-like growth factor binding protein-3 and cardiovascular risk factors in adolescents. Annals of Clinical Biochemistry, 2011, 48, 263-269.	1.6	17
447	Cardiovascular disease risk factors are highly prevalent in the office-working population of Nanjing in China. International Journal of Cardiology, 2012, 155, 212-216.	1.7	17
448	Circulating LLâ€37 is a biomarker for eczema severity in children. Journal of the European Academy of Dermatology and Venereology, 2012, 26, 518-522.	2.4	17
449	Retinal Information is Independently Associated with Cardiovascular Disease in Patients with Type 2 diabetes. Scientific Reports, 2016, 6, 19053.	3.3	17
450	Low testosterone and clinical outcomes in Chinese men with type 2 diabetes mellitus – Hong Kong Diabetes Registry. Diabetes Research and Clinical Practice, 2017, 123, 97-105.	2.8	17

#	Article	IF	CITATIONS
451	High risk of conversion to diabetes in first-degree relatives of individuals with young-onset type 2 diabetes: a 12-year follow-up analysis. Diabetic Medicine, 2017, 34, 1701-1709.	2.3	17
452	Interactive effects of prepregnancy overweight and gestational diabetes on macrosomia and large for gestational age: A population-based prospective cohort in Tianjin, China. Diabetes Research and Clinical Practice, 2019, 154, 82-89.	2.8	17
453	Plasma fibrinogen concentration in a Chinese population. Atherosclerosis, 1997, 131, 211-217.	0.8	16
454	Young Chinese adults with new onset of diabetic ketoacidosis - clinical course, autoimmune status and progression of pancreatic beta-cell function. Diabetic Medicine, 2000, 17, 295-298.	2.3	16
455	Association of two apolipoprotein Aâ€l gene <i>Msp</i> I polymorphisms with high density lipoprotein (HDL)â€cholesterol levels and indices of obesity in selected healthy Chinese subjects and in patients with earlyâ€onset typeÂ2 diabetes. Clinical Endocrinology, 2003, 59, 442-449.	2.4	16
456	Clinicopathologic characteristics of nodular glomerulosclerosis in Chinese patients with type 2 diabetes. American Journal of Kidney Diseases, 2004, 44, 1039-1049.	1.9	16
457	The preferred magnetic resonance imaging planes in quantifying visceral adipose tissue and evaluating cardiovascular risk. Diabetes, Obesity and Metabolism, 2005, 7, 547-554.	4.4	16
458	Association of Erectile Dysfunction With Cardiovascular Risk Factors and Increasing Existing Vascular Disease in Male Chinese Type 2 Diabetic Patients. Diabetes Care, 2005, 28, 2051-2053.	8.6	16
459	Associations of overweight with insulin resistance, $\hat{l}^2$ -cell function and inflammatory markers in Chinese adolescents. Pediatric Diabetes, 2008, 9, 488-495.	2.9	16
460	Childhood asthma and spirometric indices are associated with polymorphic markers of two vitamin D 25â€hydroxylase genes. Pediatric Allergy and Immunology, 2015, 26, 375-382.	2.6	16
461	Early combination versus initial metformin monotherapy in the management of newly diagnosed type 2 diabetes: An <scp>East Asian</scp> perspective. Diabetes, Obesity and Metabolism, 2021, 23, 3-17.	4.4	16
462	Familial Young-Onset Diabetes, Pre-Diabetes and Cardiovascular Disease Are Associated with Genetic Variants of DACH1 in Chinese. PLoS ONE, 2014, 9, e84770.	2.5	16
463	Use of Continuous Glucose Monitoring in the Assessment and Management of Patients With Diabetes and Chronic Kidney Disease. Frontiers in Endocrinology, 2022, 13, 869899.	3.5	16
464	Pharmacoepidemiology of ACE Inhibitor???Induced Cough. Drug Safety, 1997, 16, 150-151.	3.2	15
465	The mitochondrial tRNALeu(UUR) A to G 3243 mutation is associated with insulin-dependent and non-insulin-dependent diabetes in a Chinese population., 1997, 14, 1026-1031.		15
466	Effects and pharmacokinetics of oral glibenclamide and glipizide in Caucasian and Chinese patients with type-2 diabetes. European Journal of Clinical Pharmacology, 2000, 56, 711-714.	1.9	15
467	C-Reactive Protein and Insulin Resistance in Subjects With Thalassemia Minor and a Family History of Diabetes. Diabetes Care, 2002, 25, 1480-1481.	8.6	15
468	Effects of Obesity on the Conversion from Normal Glucose Tolerance to Diabetes in Hong Kong Chinese. Obesity, 2004, 12, 889-895.	4.0	15

#	Article	IF	CITATIONS
469	Assessing adherence to statin therapy using patient report, pill count, and an electronic monitoring device. American Journal of Health-System Pharmacy, 2005, 62, 411-415.	1.0	15
470	The effect of a one-hour Eastern stress management session on salivary cortisol. Stress and Health, 2006, 22, 45-49.	2.6	15
471	Traditional Chinese Medicine in the Treatment of Diabetes. , 2006, 11, 15-29.		15
472	Low levels of awareness of suboptimal health conditions in a high-risk working population: The "better health for better Hong Kong―health promotion campaign. International Journal of Behavioral Medicine, 2007, 14, 63-69.	1.7	15
473	Renin-angiotensin system activation in renal adipogenesis. American Journal of Physiology - Renal Physiology, 2010, 298, F391-F400.	2.7	15
474	White blood cell count and renin–angiotensin system inhibitors for the risk of cancer in type 2 diabetes. Diabetes Research and Clinical Practice, 2010, 87, 117-125.	2.8	15
475	Genetic variants for type 2 diabetes and new-onset cancer in Chinese with type 2 diabetes. Diabetes Research and Clinical Practice, 2014, 103, 328-337.	2.8	15
476	Telmisartan increases systemic exposure to rosuvastatin after single and multiple doses, and in vitro studies show telmisartan inhibits ABCG2-mediated transport of rosuvastatin. European Journal of Clinical Pharmacology, 2016, 72, 1471-1478.	1.9	15
477	Safety of atorvastatin in Asian patients within clinical trials. Cardiovascular Therapeutics, 2016, 34, 431-440.	2.5	15
478	Exendin-4 Exhibits Enhanced Anti-tumor Effects in Diabetic Mice. Scientific Reports, 2017, 7, 1791.	3.3	15
479	Efficacy and safety of alogliptin in patients with type 2 diabetes mellitus: A multicentre randomized doubleâ€blind placeboâ€controlled Phase 3 study in mainland China, Taiwan, and Hong Kong. Journal of Diabetes, 2017, 9, 386-395.	1.8	15
480	Association between educational level and cardiovascular disease and all-cause mortality in patients with type 2 diabetes: a prospective study in the Joint Asia Diabetes Evaluation Program. Clinical Epidemiology, 2018, Volume 10, 1561-1571.	3.0	15
481	Insights from VERIFY: Early Combination Therapy Provides Better Glycaemic Durability Than a Stepwise Approach in Newly Diagnosed TypeÂ2 Diabetes. Diabetes Therapy, 2020, 11, 2465-2476.	2.5	15
482	Hepatic miR-192-3p reactivation alleviates steatosis by targeting glucocorticoid receptor. JHEP Reports, 2020, 2, 100179.	4.9	15
483	Effects of a Technology-Assisted Integrated Diabetes Care Program on Cardiometabolic Risk Factors Among Patients With Type 2 Diabetes in the Asia-Pacific Region. JAMA Network Open, 2021, 4, e217557.	5.9	15
484	Pharmacodynamics and pharmacokinetics of intravenous glibenclamide in Caucasian and Chinese patients with type-2 diabetes. European Journal of Clinical Pharmacology, 2000, 55, 721-727.	1.9	14
485	Effect of Hepatic Lipase -514C->T Polymorphism and Its Interactions With Apolipoprotein C3 -482C->T and Apolipoprotein E Exon 4 Polymorphisms on the Risk of Nephropathy in Chinese Type 2 Diabetic Patients. Diabetes Care, 2005, 28, 1704-1709.	8.6	14
486	What can we learn from the recent blood glucose lowering megatrials?. Journal of Diabetes Investigation, 2011, 2, 1-5.	2.4	14

#	Article	IF	Citations
487	Associations of pubertal stage and body mass index with cardiometabolic risk in Hong Kong Chinese children: A cross-sectional study. BMC Pediatrics, 2015, 15, 136.	1.7	14
488	Association of self-reported recurrent mild hypoglycemia with incident cardiovascular disease and all-cause mortality in patients with type 2 diabetes. Medicine (United States), 2016, 95, e5183.	1.0	14
489	Evolution of Diabetes Care in Hong Kong: From the Hong Kong Diabetes Register to JADE-PEARL Program to RAMP and PEP Program. Endocrinology and Metabolism, 2018, 33, 17.	3.0	14
490	Realâ€world data reveal unmet clinical needs in insulin treatment in Asian people with type 2 diabetes: the Joint Asia Diabetes Evaluation (JADE) Register. Diabetes, Obesity and Metabolism, 2020, 22, 669-679.	4.4	14
491	Secular trends in rates of hospitalisation for lower extremity amputation and 1Âyear mortality in people with diabetes in Hong Kong, 2001–2016: a retrospective cohort study. Diabetologia, 2020, 63, 2689-2698.	6.3	14
492	High Prevalence of Depressive Symptoms in Patients With Type 1 and Type 2 Diabetes in Developing Countries: Results From the International Diabetes Management Practices Study. Diabetes Care, 2021, 44, 1100-1107.	8.6	14
493	Dexfenfluramine in Obese Chinese NIDDM Patients: A placebo-controlled investigation of the effects on body weight, glycemic control, and cardiovascular risk factors. Diabetes Care, 1997, 20, 1122-1127.	8.6	13
494	Two-hour post-glucose loading plasma glucose is the main determinant for the progression from impaired glucose tolerance to diabetes in Hong Kong Chinese. Diabetes Care, 1999, 22, 2096-2097.	8.6	13
495	Prevalence of gestational diabetes mellitus in Hong Kong based on the 1998 WHO criteria. Diabetic Medicine, 2002, 19, 80-80.	2.3	13
496	Lessons learned from young-onset diabetes in China. Current Diabetes Reports, 2003, 3, 101-107.	4.2	13
497	Effects of body mass index, plasma glucose and cholesterol levels on isolated systolic hypertension. International Journal of Cardiology, 2005, 101, 429-433.	1.7	13
498	Effects of albuminuria and renal dysfunction on development of dyslipidaemia in type 2 diabetes-the Hong Kong Diabetes Registry. Nephrology Dialysis Transplantation, 2008, 23, 2834-2840.	0.7	13
499	Genome-wide linkage scan for factors of metabolic syndrome in a Chinese population. BMC Genetics, 2010, 11, 14.	2.7	13
500	Predictive role of multilocus genetic polymorphisms in cardiovascular disease and inflammation-related genes on chronic kidney disease in Type 2 diabetes—an 8-year prospective cohort analysis of 1163 patients. Nephrology Dialysis Transplantation, 2012, 27, 190-196.	0.7	13
501	Synergistic effects of low LDL cholesterol with other factors for the risk of cancer in type 2 diabetes: the Hong Kong Diabetes Registry. Acta Diabetologica, 2012, 49, 185-193.	2.5	13
502	Hypoglycemia and Comorbidities in Type 2 Diabetes. Current Diabetes Reports, 2015, 15, 80.	4.2	13
503	Genetic effects of multiple asthma loci identified by genomewide association studies on asthma and spirometric indices. Pediatric Allergy and Immunology, 2016, 27, 185-194.	2.6	13
504	Late-Breaking Science Abstracts From the American Heart Association's Scientific Sessions 2017 and Late-Breaking Abstracts in Resuscitation Science From the Resuscitation Science Symposium 2017. Circulation, 2017, 136, e448-e467.	1.6	13

#	Article	IF	Citations
505	A polysaccharide extract from the medicinal plant Maidong inhibits the IKK–NF-κB pathway and IL-1κ–induced islet inflammation and increases insulin secretion. Journal of Biological Chemistry, 2020, 295, 12573-12587.	3.4	13
506	Development and validation of algorithms to classify type $1$ and $2$ diabetes according to age at diagnosis using electronic health records. BMC Medical Research Methodology, 2020, 20, 35.	3.1	13
507	Early combination therapy delayed treatment escalation in newly diagnosed youngâ€onset type 2 diabetes: A subanalysis of the ⟨scp⟩VERIFY⟨/scp⟩ study. Diabetes, Obesity and Metabolism, 2021, 23, 245-251.	4.4	13
508	Long-term metformin use and risk of pneumonia and related death in type 2 diabetes: a registry-based cohort study. Diabetologia, 2021, 64, 1760-1765.	6.3	13
509	Risk associations of long-term HbA1c variability and obesity on cancer events and cancer-specific death in 15,286 patients with diabetes - A prospective cohort study. The Lancet Regional Health - Western Pacific, 2022, 18, 100315.	2.9	13
510	Risk Associations of Glycemic Burden and Obesity With Liver Cancer—A 10‥ear Analysis of 15,280 Patients With Type 2 Diabetes. Hepatology Communications, 2022, 6, 1350-1360.	4.3	13
511	Effect of a Web-Based Management Guide on Risk Factors in Patients With Type 2 Diabetes and Diabetic Kidney Disease. JAMA Network Open, 2022, 5, e223862.	5.9	13
512	Detection of increased serum miR-122-5p and miR-455-3p levels before the clinical diagnosis of liver cancer in people with type 2 diabetes. Scientific Reports, 2021, 11, 23756.	3.3	13
513	Acute Tubular Necrosis Following Endosulphan Insecticide Poisoning. Journal of Toxicology: Clinical Toxicology, 1995, 33, 67-69.	1.5	12
514	Topographical associations between islet endocrine cells and duct epithelial cells in the adult human pancreas. Clinical Endocrinology, 2008, 69, 400-406.	2.4	12
515	Meta-analysis of trial data may support a causal role of hyperglycaemia in cancer. Diabetologia, 2011, 54, 709-710.	6.3	12
516	Use of thiazolidinedione and cancer risk in Type 2 diabetes: The Hong Kong diabetes registry. Diabetes Research and Clinical Practice, 2012, 97, e13-e17.	2.8	12
517	Histopathological Correlations of Islet Amyloidosis With Apolipoprotein E Polymorphisms in Type 2 Diabetic Chinese Patients. Pancreas, 2013, 42, 1129-1137.	1.1	12
518	Self-monitoring of blood glucose (SMBG) and glycaemic control in Cameroon: Results of the International Diabetes Management Practices Study (IDMPS). Diabetes Research and Clinical Practice, 2017, 126, 198-201.	2.8	12
519	The Relationship of Quantitative Retinal Capillary Network to Kidney Function in Type 2 Diabetes. American Journal of Kidney Diseases, 2018, 71, 916-918.	1.9	12
520	Insulin resistance versus $\hat{l}^2$ -cell dysfunction in type 2 diabetes: where public and personalised health meet. Lancet Diabetes and Endocrinology,the, 2020, 8, 92-93.	11.4	12
521	Nonalbuminuric Diabetic Kidney Disease and Risk of All-Cause Mortality and Cardiovascular and Kidney Outcomes in Type 2 Diabetes: Findings From the Hong Kong Diabetes Biobank. American Journal of Kidney Diseases, 2022, 80, 196-206.e1.	1.9	12
522	Associations of the HOMA2â€%B and HOMA2â€IR with progression to diabetes and glycaemic deterioration in young and middleâ€aged Chinese. Diabetes/Metabolism Research and Reviews, 2022, 38, e3525.	4.0	12

#	Article	IF	Citations
523	Glucose-lowering drug use, glycemic outcomes, and severe hypoglycemia: 18-Year trends in OÂ-9 million adults with Diabetes in Hong Kong (2002–2019). The Lancet Regional Health - Western Pacific, 2022, 26, 100509.	2.9	12
524	The Effects of Enalapril and Nifedipine on Carbohydrate and Lipid Metabolism in NIDDM. Diabetes Care, 1994, 17, 859-862.	8.6	11
525	A prescription survey in a hospital hypertension outpatient clinic. British Journal of Clinical Pharmacology, 1997, 44, 577-582.	2.4	11
526	Clinical and biochemical characteristics of type 2 diabetic patients on continuous ambulatory peritoneal dialysis: Relationships with insulin requirement. American Journal of Kidney Diseases, 1999, 34, 514-520.	1.9	11
527	Cardiovascular disease in Chinese type 2 diabetic women is associated with a prolonged QTc interval. International Journal of Cardiology, 2000, 76, 75-80.	1.7	11
528	Obesity, insulin resistance and isolated low high-density-lipoprotein cholesterol in Chinese subjects. Diabetic Medicine, 2001, 18, 663-666.	2.3	11
529	Renin-Angiotensin System Gene Polymorphisms and Retinopathy in Chinese Patients With Type 2 Diabetes. Diabetes Care, 2003, 26, 1643-1644.	8.6	11
530	Effects of chronic hyperglycaemia on incident stroke in Hong Kong Chinese patients with type 2 diabetes. Diabetes/Metabolism Research and Reviews, 2007, 23, 220-226.	4.0	11
531	A case-control study of apoA5 â^1131Tâ†'C polymorphism that examines the role of triglyceride levels in diabetic nephropathy. Journal of Diabetes and Its Complications, 2007, 21, 158-163.	2.3	11
532	Metabolic syndrome by the new IDF criteria in Hong Kong Chinese adolescents and its prediction by using body mass index. Acta Paediatrica, International Journal of Paediatrics, 2008, 97, 1738-1742.	1.5	11
533	Association of <i>early growth responseâ€1</i> gene polymorphisms with total IgE and atopy in asthmatic children. Pediatric Allergy and Immunology, 2009, 20, 142-150.	2.6	11
534	The reproductive and metabolic effect of rosiglitazoneÂon Chinese women with polycystic ovarian syndrome—a double-blind randomized placebo-controlled study. Fertility and Sterility, 2011, 96, 445-451.e1.	1.0	11
535	Copy number variation analysis based on AluScan sequences. Journal of Clinical Bioinformatics, 2014, 4, 15.	1.2	11
536	Long-Term Risk of Cardiovascular Disease among Type 2 Diabetic Patients with Asymptomatic Intracranial Atherosclerosis: A Prospective Cohort Study. PLoS ONE, 2014, 9, e106623.	2.5	11
537	Regular mailing of personalized feedback reports improves glycemic control in diabetes: <scp>A</scp> randomized controlled trial. Journal of Diabetes, 2017, 9, 536-538.	1.8	11
538	Efficacy and Safety of Linagliptin in 2681 Asian Patients Stratified by Age, Obesity, and Renal Function: A Pooled Analysis of Randomized Clinical Trials. Advances in Therapy, 2017, 34, 2150-2162.	2.9	11
539	Indicators of socio-economic status and risk of gestational diabetes mellitus in pregnant women in urban Tianjin, China. Diabetes Research and Clinical Practice, 2018, 144, 192-199.	2.8	11
540	Effects of Lifestyle Intervention of Maternal Gestational Diabetes Mellitus on Offspring Growth Pattern Before Two Years of Age. Diabetes Care, 2021, 44, e42-e44.	8.6	11

#	Article	IF	CITATIONS
541	One in Seven Insulin-Treated Patients in Developing Countries Reported Poor Persistence with Insulin Therapy: Real World Evidence from the Cross-Sectional International Diabetes Management Practices Study (IDMPS). Advances in Therapy, 2021, 38, 3281-3298.	2.9	11
542	Trends in kidney failure and kidney replacement therapy in people with diabetes in Hong Kong, 2002-2015: A retrospective cohort study. The Lancet Regional Health - Western Pacific, 2021, 11, 100165.	2.9	11
543	Guidance on the management of familial hypercholesterolaemia in Hong Kong: an expert panel consensus viewpoint. Hong Kong Medical Journal, 2018, 24, 408-415.	0.1	11
544	Skin autofluorescence is associated with progression of kidney disease in type 2 diabetes: A prospective cohort study from the Hong Kong diabetes biobank. Nutrition, Metabolism and Cardiovascular Diseases, 2022, 32, 436-446.	2.6	11
545	A Clinical Perspective of the Multifaceted Mechanism of Metformin in Diabetes, Infections, Cognitive Dysfunction, and Cancer. Pharmaceuticals, 2022, 15, 442.	3.8	11
546	Relative leucocyte telomere length is associated with incident end-stage kidney disease and rapid decline of kidney function in type 2 diabetes: analysis from the Hong Kong Diabetes Register. Diabetologia, 2022, 65, 375-386.	6.3	11
547	A Chinese Family with Nonâ€insulinâ€dependent Diabetes of Early Onset and Severe Diabetic Complications. Diabetic Medicine, 1990, 7, 211-214.	2.3	10
548	HYPOGLYCAEMIA AND HODGKIN'S DISEASE. British Journal of Haematology, 1990, 76, 434-436.	2.5	10
549	Severe hypoglycaemia in Chinese patients with non-insulin-dependent diabetes treated with insulin or sulphonylureas. Pharmacoepidemiology and Drug Safety, 1992, 1, 207-211.	1.9	10
550	Treatment of Chinese acromegaly with a combination of bromocriptine and octreotide. Australian and New Zealand Journal of Medicine, 2000, 30, 457-461.	0.5	10
551	Increased leptin concentrations and lack of gender difference in Type 2 diabetic patients with nephropathy. Diabetes Research and Clinical Practice, 2004, 64, 93-98.	2.8	10
552	Losartan reduces the costs of diabetic end-stage renal disease: An Asian perspective. Nephrology, 2005, 10, 520-524.	1.6	10
553	<i>PTGDR</i> is not a major candidate gene for asthma and atopy in Chinese children. Pediatric Allergy and Immunology, 2009, 20, 556-562.	2.6	10
554	The pivotal role of protein kinase C zeta (PKCzeta) in insulin- and AMP-activated protein kinase (AMPK)-mediated glucose uptake in muscle cells. Cellular Signalling, 2010, 22, 1513-1522.	3.6	10
555	Development and validation of equations estimating glomerular filtration rates in Chinese patients with type 2 diabetes. Kidney International, 2010, 77, 729-735.	<b>5.</b> 2	10
556	Treatment and Landmark Clinical Trials for Renoprotection. Contributions To Nephrology, 2011, 170, 184-195.	1.1	10
557	Comment: Analyses Using Time-Dependent Pioglitazone Usage in Cox Models May Lead to Wrong Conclusions About Its Association With Cancer. Diabetes Care, 2011, 34, e136-e136.	8.6	10
558	Cadmium and lead in Hong Kong school children. Pathology, 2012, 44, 626-631.	0.6	10

#	Article	IF	Citations
559	Management of Familial Hypercholesterolemia in Hong Kong. Journal of Atherosclerosis and Thrombosis, 2016, 23, 520-531.	2.0	10
560	Diabetes education and health insurance: How they affect the quality of care provided to people with type 1 diabetes in Latin America. Data from the International Diabetes Mellitus Practices Study (IDMPS). Diabetes Research and Clinical Practice, 2019, 147, 47-54.	2.8	10
561	Pancreatic Sirtuin 3 Deficiency Promotes Hepatic Steatosis by Enhancing 5-Hydroxytryptamine Synthesis in Mice With Diet-Induced Obesity. Diabetes, 2021, 70, 119-131.	0.6	10
562	Increased co-expression of PSMA2 and GLP-1 receptor in cervical cancer models in type 2 diabetes attenuated by Exendin-4: A translational case-control study. EBioMedicine, 2021, 65, 103242.	6.1	10
563	Shortened relative leukocyte telomere length is associated with all-cause mortality in type 2 diabetes- analysis from the Hong Kong Diabetes Register. Diabetes Research and Clinical Practice, 2021, 173, 108649.	2.8	10
564	Diabetes, insulin and cancer risk. World Journal of Diabetes, 2012, 3, 60.	3.5	10
565	Anticholinergic poisoning from Chinese herbal medicines. Australian and New Zealand Journal of Medicine, 1994, 24, 317-318.	0.5	9
566	Urinary dopamine and noradrenaline outputs during oral salt loading in healthy Chinese subjects with a family history of hypertension. Autonomic and Autacoid Pharmacology, 1996, 16, 1-6.	0.6	9
567	Single-dose pharmacokinetics of paracetamol and its conjugates in Chinese non-insulin-dependent diabetic patients with renal impairment. European Journal of Clinical Pharmacology, 1997, 52, 285-288.	1.9	9
568	HLA typing and immunological characterization of young-onset diabetes mellitus in a Hong Kong Chinese population. Diabetic Medicine, 2001, 18, 22-28.	2.3	9
569	Identification of the intron 14 splicing defect of the cholesteryl ester transfer protein gene in Hong Kong Chinese. Clinical Genetics, 2001, 59, 287-289.	2.0	9
570	Prognostic role of serum ACE activity on outcome of type 2 diabetic patients on chronic ambulatory peritoneal dialysis. American Journal of Kidney Diseases, 2002, 39, 1054-1060.	1.9	9
571	Considerations on blood glucose management in Type 2 diabetes mellitus. Diabetes/Metabolism Research and Reviews, 2002, 18, 273-285.	4.0	9
572	Assessing adherence to statin therapy using patient report, pill count, and an electronic monitoring device. American Journal of Health-System Pharmacy, 2005, 62, 411-415.	1.0	9
573	Association of Smoking with Increasing Vascular Involvement in Type 2 Diabetic Chinese Patients. Experimental and Clinical Endocrinology and Diabetes, 2006, 114, 301-305.	1.2	9
574	Associations of insulin-like growth factor binding protein-3 gene polymorphisms with IGF-I activity and lipid parameters in adolescents. International Journal of Obesity, 2009, 33, 1446-1453.	3.4	9
575	Atherosclerotic Vascular Disease Rather than Metabolic Syndrome Predicts Ischemic Stroke in Diabetic Patients. Cerebrovascular Diseases, 2010, 30, 374-379.	1.7	9
576	Glucose lowering effect of transgenic human insulin-like growth factor-I from rice: in vitro and in vivostudies. BMC Biotechnology, 2011, 11, 37.	3.3	9

#	Article	IF	Citations
577	Short Body Height and Pre-pregnancy Overweight for Increased Risk of Gestational Diabetes Mellitus: A Population-Based Cohort Study. Frontiers in Endocrinology, 2018, 9, 349.	3.5	9
578	SNPs in PRKCAâ€HIF1Aâ€GLUT1 are associated with diabetic kidney disease in a Chinese Han population with type 2 diabetes. European Journal of Clinical Investigation, 2020, 50, e13264.	3.4	9
579	Skin autofluorescence is associated with higher risk of cardiovascular events in Chinese adults with type 2 diabetes: A prospective cohort study from the Hong Kong Diabetes Biobank. Journal of Diabetes and Its Complications, 2021, 35, 108015.	2.3	9
580	Evaluation of drug usage and expenditure in a hospital diabetes clinic. Journal of Clinical Pharmacy and Therapeutics, 1998, 23, 49-56.	1.5	8
581	A 67-year-old woman with recurrent hypoglycemia: non-islet cell tumour hypoglycemia. Cmaj, 2005, 173, 359-361.	2.0	8
582	Thresholds of risk factors for ischemic stroke in type 2 diabetic patients with and without albuminuria—A non-linear approach. Clinical Neurology and Neurosurgery, 2008, 110, 701-709.	1.4	8
583	Adiposity of the heart revisited: Reversal of dilated cardiomyopathy in a patient with Cushing's syndrome. International Journal of Cardiology, 2011, 151, e22-e23.	1.7	8
584	Interactions between Genetic Variants of FLG and Chromosome 11q13 Locus Determine Susceptibility for Eczema Phenotypes. Journal of Investigative Dermatology, 2012, 132, 1930-1932.	0.7	8
585	Additive effect of aldose reductase Z-4 microsatellite polymorphism and glycaemic control on cataract development in type 2 diabetes. Journal of Diabetes and Its Complications, 2014, 28, 147-151.	2.3	8
586	Early gene–diet interaction between glucokinase regulatory protein (GCKR) polymorphism, vegetable and fish intakes in modulating triglyceride levels in healthy adolescents. Nutrition, Metabolism and Cardiovascular Diseases, 2015, 25, 951-958.	2.6	8
587	ABO blood types and postpartum depression among Chinese women: A prospective cohort study in Tianjin, China. Women and Health, 2018, 58, 685-698.	1.0	8
588	Gender differences in the associations between insomnia and glycemic control in patients with type 2 diabetes: a cross-sectional study. Sleep, 2019, 42, .	1.1	8
589	Interactive effects of testosterone and the androgen receptor CAG repeat length polymorphism on cardiovascularâ€renal events and mortality in men with diabetes. Diabetes/Metabolism Research and Reviews, 2019, 35, e3081.	4.0	8
590	Factors Determining the Blood Pressure Response to Enalapril and Nifedipine in Hypertension Associated With NIDDM. Diabetes Care, 1995, 18, 1001-1006.	8.6	7
591	Supplement to the use of a paired value of fasting plasma glucose and glycated hemoglobin in predicting the likelihood of having diabetes. Diabetes Care, 1998, 21, 2032-2033.	8.6	7
592	Age, body mass index and 2-hour plasma glucose are the major determinants of blood pressure in Chinese women newly diagnosed to have glucose intolerance. International Journal of Cardiology, 1999, 69, 33-39.	1.7	7
593	Association between smoking, pancreatic insulin secretion and insulin resistance in Chinese subjects with or without glucose intolerance. Chinese Medical Journal, 2007, 120, 2233-2237.	2.3	7
594	ORIGINAL ARTICLE: Associations of the growth hormone receptor (⟨i⟩GHR⟨ i⟩) gene polymorphisms with adiposity and IGF†activity in adolescents. Clinical Endocrinology, 2010, 73, 313-322.	2.4	7

#	Article	IF	Citations
595	Diabetes and Noncommunicable Disease. JAMA - Journal of the American Medical Association, 2013, 310, 916.	7.4	7
596	Intrarenal arterial resistance is associated with microvascular complications in Chinese type 2 diabetic patients. Nephrology Dialysis Transplantation, 2013, 28, 651-658.	0.7	7
597	Prediction of women's longâ€ŧerm cardiometabolic risks using glycemic indices during pregnancy. Journal of Obstetrics and Gynaecology Research, 2013, 39, 484-491.	1.3	7
598	Curvilinear associations of sleep patterns during weekdays and weekends with glycemic control in type 2 diabetes: the Hong Kong Diabetes Registry. Acta Diabetologica, 2017, 54, 151-162.	2.5	7
599	Interactions between Prepregnancy Overweight and Passive Smoking for Macrosomia and Large for Gestational Age in Chinese Pregnant Women. Obesity Facts, 2021, 14, 520-530.	3.4	7
600	Genetic Associations of Type 2 Diabetes with Islet Amyloid Polypeptide Processing and Degrading Pathways in Asian Populations. PLoS ONE, 2013, 8, e62378.	2.5	7
601	The inter-relationships between albuminuria, plasma albumin concentration and indices of glycaemic control in non-insulin-dependent diabetes mellitus. Clinica Chimica Acta, 1992, 210, 179-185.	1.1	6
602	Recurrent Glibenclamide-Induced Hypoglycemia: The Importance of Obtaining a Comprehensive Medication History. Annals of Pharmacotherapy, 1994, 28, 119-120.	1.9	6
603	Urinary dopamine and noradrenaline outputs during large acute changes in oral salt intake in healthy Chinese subjects. Autonomic and Autacoid Pharmacology, 1994, 14, 317-323.	0.6	6
604	Association of the D8S282 marker near the lipoprotein lipase gene locus with systolic blood pressure in healthy Chinese subjects. Journal of Hypertension, 2002, 20, 2199-2204.	0.5	6
605	Triglyceride, albuminuria and blood pressure are the major associations of non-fatal cardiovascular disease in Chinese type 2 diabetes. Acta Diabetologica, 2003, 40, 80-84.	2.5	6
606	Genetic variants of hepatocyte nuclear factor- $1\hat{l}^2$ in Chinese young-onset diabetic patients with nephropathy. Journal of Diabetes and Its Complications, 2003, 17, 369-373.	2.3	6
607	Reduction of Total Homocysteine Levels by Oral Folic Acid Fails to Improve Endothelial Function in Children With Chronic Renal Failure. Circulation, 2003, 107, e6-7; author reply e6-7.	1.6	6
608	A novel mutation of SLC22A12 gene causing primary renal hypouricemia in a patient with metabolic syndrome. Clinica Chimica Acta, 2008, 398, 157-158.	1.1	6
609	APOE Genotype-Function Relationship: Evidence of â^'491 A/T Promoter Polymorphism Modifying Transcription Control but Not Type 2 Diabetes Risk. PLoS ONE, 2011, 6, e24669.	2.5	6
610	Cancer Risk in Type 2 Diabetes. Current Diabetes Reports, 2012, 12, 325-328.	4.2	6
611	Selfâ€reported waist circumference: a screening tool for classifying children with overweight/obesity and cardiometabolic risk factor clustering. Pediatric Obesity, 2012, 7, 110-120.	2.8	6
612	Temporal changes in obesity and sleep habits in Hong Kong Chinese school children: a prospective study. Scientific Reports, 2019, 9, 5881.	3.3	6

#	Article	IF	CITATIONS
613	Sudomotor dysfunction independently predicts incident cardiovascular–renal events and all-cause death in type 2 diabetes: the Joint Asia Diabetes Evaluation register. Nephrology Dialysis Transplantation, 2019, 34, 1320-1328.	0.7	6
614	Longâ€term maternal cardiometabolic outcomes 22Âyears after gestational diabetes mellitus. Journal of Diabetes Investigation, 2020, 11, 985-993.	2.4	6
615	<p><em>CYP2C19</em>*<em>2</em> Polymorphism Is Associated with Impaired Oral Clearance of Gliclazide in Healthy Chinese</p> . Pharmacogenomics and Personalized Medicine, 2019, Volume 12, 397-401.	0.7	6
616	Impacts of gestational diabetes on quality of life in Chinese pregnant women in urban Tianjin, China. Primary Care Diabetes, 2020, 14, 425-430.	1.8	6
617	Screening, prevalence, treatment and control of kidney disease in patients with type 1 and type 2 diabetes in low-to-middle-income countries (2005–2017): the International Diabetes Management Practices Study (IDMPS). Diabetologia, 2021, 64, 1246-1255.	6.3	6
618	Association between FGF19, FGF21 and lipocalin-2, and diabetes progression in PCOS. Endocrine Connections, 2021, 10, 1243-1252.	1.9	6
619	Clinical Predictors and Long-term Impact of Acute Kidney Injury on Progression of Diabetic Kidney Disease in Chinese Patients With Type 2 Diabetes. Diabetes, 2022, 71, 520-529.	0.6	6
620	Time-varying risk associations of renin angiotensin system inhibitors with pneumonia and related deaths in a cohort of 252,616 patients with diabetes (2002–2019). Diabetes Research and Clinical Practice, 2022, 185, 109233.	2.8	6
621	The emerging role of incretins and twincretins. Nature Reviews Endocrinology, 2022, 18, 73-74.	9.6	6
622	Internuclear ophthalmoplegia in tuberculous meningitis. Tubercle, 1989, 70, 61-64.	0.6	5
623	17 alpha-Hydroxylase deficiency with persistence of mullerian ducts in a genotypic male and paradoxical aldosterone secretion Postgraduate Medical Journal, 1993, 69, 159-162.	1.8	5
624	A patient compliance survey in a general medical clinic. Journal of Clinical Pharmacy and Therapeutics, 1997, 22, 323-326.	1.5	5
625	Use of indapamide in hospital and community clinics and its effect on plasma potassium in Chinese patients. Journal of Clinical Pharmacy and Therapeutics, 1998, 23, 295-302.	1.5	5
626	Health Hazards of Obesity: An Overview. , 0, , 215-234.		5
627	Effects of systolic and diastolic blood pressures on incident coronary heart disease and allâ $\in$ cause death in Chinese women with Type 2 diabetes: The Hong Kong Diabetes Registry. Journal of Diabetes, 2009, 1, 90-98.	1.8	5
628	Fatty acids inhibit insulin-mediated glucose transport associated with actin remodeling in rat L6 muscle cells. Acta Diabetologica, 2010, 47, 331-339.	2.5	5
629	Hypoadiponectinaemia enhances waist circumference as a predictor of glucose intolerance and clustering of risk factors in Chinese men. Diabetes and Metabolism, 2010, 36, 192-197.	2.9	5
630	Pharmacoepidemiological profiles of oral hypoglycemic agents among 28,773 Chinese patients with diabetes. Diabetes Research and Clinical Practice, 2012, 96, 319-325.	2.8	5

#	Article	IF	Citations
631	Use of laparoscopic sleeve gastrectomy and adjustable gastric banding for suboptimally controlled diabetes in Hong Kong. Diabetes, Obesity and Metabolism, 2012, 14, 372-374.	4.4	5
632	Renin angiotensin system inhibitors may attenuate low LDL cholesterolâ€related cancer risk in type 2 diabetes. Diabetes/Metabolism Research and Reviews, 2014, 30, 415-423.	4.0	5
633	Childhood asthma is associated with polymorphic markers of <i>PROC</i> on 2q14 in addition to 17q21 locus. Pediatric Allergy and Immunology, 2015, 26, 173-180.	2.6	5
634	A Pilot Study to Compare Meal-Triggered Gastric Electrical Stimulation and Insulin Treatment in Chinese Obese Type 2 Diabetes. Diabetes Technology and Therapeutics, 2015, 17, 283-290.	4.4	5
635	Triple combination of insulin glargine, sitagliptin and metformin in type 2 diabetes: The EASIE post-hoc analysis and extension trial. Journal of Diabetes and Its Complications, 2015, 29, 134-141.	2.3	5
636	Diabetes: A Cinderella Subject We Can't Afford to Ignore. PLoS Medicine, 2016, 13, e1002068.	8.4	5
637	Progression to treatment failure among Chinese patients with type 2 diabetes initiated on metformin versus sulphonylurea monotherapy—The Hong Kong Diabetes Registry. Diabetes Research and Clinical Practice, 2016, 112, 57-64.	2.8	5
638	Modifying Effect of Body Mass Index on Survival in Elderly Type 2 Diabetic Patients: Hong Kong Diabetes Registry. Journal of the American Medical Directors Association, 2016, 17, 276.e15-276.e22.	2.5	5
639	Baseline characteristics and temporal differences in Acarbose Cardiovascular Evaluation (ACE) trial participants. American Heart Journal, 2018, 199, 170-175.	2.7	5
640	Insulin glargine compared to neutral protamine Hagedorn (NPH) insulin in patients with type-2 diabetes uncontrolled with oral anti-diabetic agents alone in Hong Kong: a cost-effectiveness analysis. Cost Effectiveness and Resource Allocation, 2019, 17, 13.	1.5	5
641	Cadherinâ€related family member 3 gene impacts childhood asthma in Chinese children. Pediatric Allergy and Immunology, 2020, 31, 133-142.	2.6	5
642	Glycemic Variability and Time in Range During Self-titration of Once Daily Insulin Glargine 300 U/ml Versus Neutral Protamine Hagedorn Insulin in Insulin-naÃ-ve Chinese Type 2 Diabetes Patients. Diabetes Therapy, 2021, 12, 1399-1413.	2.5	5
643	Usefulness of cutâ€off points of International criteria for prediction of postâ€partum diabetes and prediabetes among Chinese women with gestational diabetes. Diabetes/Metabolism Research and Reviews, 2021, 37, e3456.	4.0	5
644	Perinatal famine is associated with excess risk of proliferative retinopathy in patients with type 2 diabetes. Acta Ophthalmologica, 2022, 100, .	1.1	5
645	Drug-Related Hyperglycemia. JAMA - Journal of the American Medical Association, 2002, 287, 714-715.	7.4	5
646	<i>CYP2C19</i> Lossâ€ofâ€function Polymorphisms are Associated with Reduced Risk of Sulfonylurea Treatment Failure in Chinese Patients with Type 2 Diabetes. Clinical Pharmacology and Therapeutics, 2022, 111, 461-469.	4.7	5
647	Human Serum Metabolites as Potential Mediators from Type 2 Diabetes and Obesity to COVID-19 Severity and Susceptibility: Evidence from Mendelian Randomization Study. Metabolites, 2022, 12, 598.	2.9	5
648	No relationship between antibodies to GAD and microangiopathic complications in young Chinese diabetic patients. Diabetes Care, 2000, 23, 1045-1046.	8.6	4

#	Article	IF	CITATIONS
649	Renal Carcinogenesis After Uninephrectomy. Translational Oncology, 2009, 2, 258-263.	3.7	4
650	Are Large Sample Size Studies the Answer to Evaluate Effects of Drug Use in Non–Clinical Trial Settings?. Journal of Clinical Oncology, 2011, 29, 4464-4465.	1.6	4
651	Comment on: Suissa and Azoulay. Metformin and the Risk of Cancer: Time-Related Biases in Observational Studies. Diabetes Care 2012;35:2665–2673. Diabetes Care, 2013, 36, e87-e87.	8.6	4
652	Feature Extraction of Radial Arterial Pulse. , 2014, , .		4
653	How can we optimise diabetes care in real-world practice?. Lancet Diabetes and Endocrinology,the, 2017, 5, 927-929.	11.4	4
654	Explaining the high prevalence of youngâ€onset diabetes among Asians and Indigenous Australians. Medical Journal of Australia, 2017, 207, 331-332.	1.7	4
655	Excess Burden of Mental Illness and Hospitalization in Young-Onset Type 2 Diabetes. Annals of Internal Medicine, 2019, 171, 78.	3.9	4
656	Effects of lifestyle intervention during pregnancy on postpartum diabetes among Chinese women with gestational diabetes. Diabetologia, 2021, 64, 255-258.	6.3	4
657	Migration and diabetes incidence among Chinese adults in Canada, China, Hong Kong, and Taiwan: An international population-based comparative study from 2000 to 2017. Diabetes Research and Clinical Practice, 2021, 180, 109062.	2.8	4
658	Increased Growth of a Newly Established Mouse Epithelial Cell Line Transformed with HPV-16 E7 in Diabetic Mice. PLoS ONE, 2016, 11, e0164490.	2.5	4
659	Poor Glycemic Control in People with T1D and T2Dâ€"Results from the International Diabetes Management Practices Study (IDMPS). Diabetes, 2018, 67, .	0.6	4
660	Metabolic syndrome and type 2 diabetes: the Hong Kong perspective. Clinical Biochemist Reviews, 2005, 26, 51-7.	3.3	4
661	Metformin and the risk of cancer in type 2 diabetes: methodological challenges and perspectives. Annals of Translational Medicine, 2014, 2, 52.	1.7	4
662	Autotaxin signaling facilitates $\hat{l}^2$ cell dedifferentiation and dysfunction induced by Sirtuin 3 deficiency. Molecular Metabolism, 2022, 60, 101493.	6.5	4
663	Integratome analysis of adipose tissues reveals abnormal epigenetic regulation of adipogenesis, inflammation, and insulin signaling in obese individuals with type 2 diabetes. Clinical and Translational Medicine, $2021, 11, e596$ .	4.0	4
664	Assessment of Bidirectional Relationships between Leisure Sedentary Behaviors and Neuropsychiatric Disorders: A Two-Sample Mendelian Randomization Study. Genes, 2022, 13, 962.	2.4	4
665	Acute confusion, Chinese herbal medicines and tuberculous meningitis. Australian and New Zealand Journal of Medicine, 1994, 24, 590-591.	0.5	3
666	Use of a paired value of fasting plasma glucose and glycated hemoglobin in predicting the likelihood of diabetes in a community. Diabetes Care, 1999, 22, 1908-1909.	8.6	3

#	Article	IF	CITATIONS
667	Recurrent Hypoglycaemia in a Patient with Metastatic Pancreatic Carcinoma. PLoS Medicine, 2006, 3, e331.	8.4	3
668	<i>CHIA</i> confers susceptibility to childhood eczema. British Journal of Dermatology, 2010, 163, 1360-1362.	1.5	3
669	Glucose intolerance and cardiovascular risk factors in Hong Kong: Data from two occupation-based cross-sectional surveys. Diabetes Research and Clinical Practice, 2010, 90, 222-227.	2.8	3
670	Associations of intra-renal arterial resistance index with chronic kidney disease and carotid intima-media thickness in type 2 diabetes mellitus. Diabetes Research and Clinical Practice, 2011, 92, e37-e40.	2.8	3
671	HUS and diabetic nephropathy. Nephrology Dialysis Transplantation, 2012, 27, ii11-ii13.	0.7	3
672	Health-Related Quality of Life in Chinese Patients with Type 2 Diabetes: An Analysis of the Joint Asia Diabetes Evaluation (JADE) Program. Journal of Diabetes & Metabolism, 2014, 05, .	0.2	3
673	International Diabetes Management Practice Study (Idmps): Resource Use Associated With Type 2 Diabetes In Africa, Middle East, South Asia, Eurasia and Turkey. Value in Health, 2015, 18, A619.	0.3	3
674	Type 2 Diabetes in Turkey – A Cost of Illness Study. Value in Health, 2016, 19, A669.	0.3	3
675	Crossâ€sectional survey of biosimilar insulin utilization in Asia: The Joint Asia Diabetes Evaluation Program. Journal of Diabetes Investigation, 2018, 9, 1312-1322.	2.4	3
676	Regional evidence and international recommendations to guide lipid management in Asian patients with type 2 diabetes with special reference to renal dysfunction. Journal of Diabetes, 2018, 10, 200-212.	1.8	3
677	Associations of Renal Augmented Velocity Index with Arterial Stiffness, Carotid Intima–Media Thickness and Blood Pressure, in Comparison with Renal Resistive Index. Ultrasound in Medicine and Biology, 2021, 47, 1279-1288.	1.5	3
678	Phenotypic and Genetic Heterogeneity in a Thai Glucokinase MODY Family Reveals the Complexity of Young-Onset Diabetes. Frontiers in Endocrinology, 2021, 12, 690343.	3.5	3
679	Diabetes in China and the Western Pacific Region. , 2017, , 63-83.		3
680	Monitoring may need to be prolonged in patients given warfarin and amiodarone. BMJ: British Medical Journal, 1996, 313, 301-302.	2.3	3
681	Sodium-glucose co-transporter-2 inhibitors: know the patient and the drugs., 2019, 25, 268-270.		3
682	Reasons for Discontinuation of Insulin Therapyâ€"Results from the International Diabetes Management Practices Study (IDMPS). Diabetes, 2018, 67, 1026-P.	0.6	3
683	Nonachievement of Glycemic Targetâ€"Results from the International Diabetes Management Practices Study (IDMPS). Diabetes, 2018, 67, 1030-P.	0.6	3
684	External validation of a shortened screening tool using individual participant data meta-analysis: A case study of the Patient Health Questionnaire-Dep-4. Methods, 2022, 204, 300-311.	3.8	3

#	Article	IF	CITATIONS
685	Continuous Glucose Monitoring Metrics in the Assessment of Glycemia in Moderate-to-Advanced CKD in Diabetes. Kidney International Reports, 2022, 7, 1354-1363.	0.8	3
686	DIDMOAD syndrome in a Chinese male with HLA DR7 DRw12. Postgraduate Medical Journal, 1987, 63, 1109-1109.	1.8	2
687	Hyperfibrinogenaemia did not improve after treating hyperglycaemia in Chinese type 2 diabetic patients. Annals of Clinical Biochemistry, 2000, 37, 655-661.	1.6	2
688	Renal kallikrein-kinin system, but not renal dopamine system, mediates the natriuretic response to intravenous saline infusion in healthy Chinese subjects. Autonomic and Autacoid Pharmacology, 2000, 20, 37-45.	0.6	2
689	Long-term effects of hormone replacement therapy. Lancet, The, 2003, 361, 254.	13.7	2
690	Antihypertensive Therapy and Incidence of Type 2 Diabetes in an Elderly Cohort: Response to Padwal et al Diabetes Care, 2005, 28, 762-762.	8.6	2
691	To: Holstein A, Stumvoll M (2005) Contraindications can damage your health—is metformin a case in point? Diabetologia 48:2454–2459. Diabetologia, 2006, 49, 1127-1128.	6.3	2
692	The Authors??? Reply. Drug Safety, 2007, 30, 728-729.	3.2	2
693	Riskâ€benefit analysis of use of statins for primary prevention of cardiovascular disease in subjects without diabetes. Journal of Diabetes Investigation, 2013, 4, 344-346.	2.4	2
694	Epidemiology/Genetics. Diabetes, 2013, 62, A365-A438.	0.6	2
695	Conceptual Perspectives on the Co-Occurrence of Mental and Physical Disease: Diabetes and Depression as a Model. Key Issues in Mental Health, 2015, , 1-14.	0.6	2
696	A proof-of-concept study to evaluate the efficacy and safety of BTI320 on post-prandial hyperglycaemia in Chinese subjects with pre-diabetes. BMC Endocrine Disorders, 2018, 18, 59.	2.2	2
697	Augmented Velocity Index: A New Doppler Index Associated with Arterial Stiffness. Ultrasound in Medicine and Biology, 2019, 45, 2747-2757.	1.5	2
698	Tackling obesity during the COVIDâ€19 pandemic. Diabetes/Metabolism Research and Reviews, 2021, 37, e3393.	4.0	2
699	Investigating the role of $\langle i \rangle$ dachshund $b \langle i \rangle$ in the development of the pancreatic islet in zebrafish. Journal of Diabetes Investigation, 2021, 12, 710-727.	2.4	2
700	No Improvement in Glycemic Control or Rates of Diabetes-Related Complications for People with Type 2 Diabetesâ€"Results from 10 Years of the International Diabetes Management Practices Study (IDMPS). Diabetes, 2018, 67, .	0.6	2
701	Factors Associated with Achievement of Glycemic Goal in Africaâ€"Results from the International Diabetes Management Practices Study (IDMPS). Diabetes, 2018, 67, 2399-PUB.	0.6	2
702	Comment on Dawed et al. Genome-Wide Meta-Analysis Identifies Genetic Variants Associated With Glycemic Response to Sulfonylureas. Diabetes Care 2021;44:2673–2682. Diabetes Care, 2022, 45, e80-e81.	8.6	2

#	Article	IF	Citations
703	Implementation of Precision Genetic Approaches for Type 1 and 2 Diabetes., 2022,, 111-129.		2
704	Combined associations of family history and self-management with age at diagnosis and cardiometabolic risk in 86,931 patients with type 2 diabetes: Joint Asia Diabetes Evaluation (JADE) Register from 11 countries. BMC Medicine, 2022, 20, .	5 <b>.</b> 5	2
705	Postpartum Hypertension, Bromocriptine and Phenylpropanolamine. Drug Investigation, 1994, 8, 254-256.	0.6	1
706	Phenotypic Heterogeneity and Associations of Two Aldose Reductase Gene Polymorphisms With Nephropathy and Retinopathy in Type 2 Diabetes: Response to Ng et al Diabetes Care, 2004, 27, 290-290.	8.6	1
707	P-85 Metabolic syndrome identifies chronic kidney disease in Chinese patients with Type 2 diabetes â€" a prospective study. Diabetes Research and Clinical Practice, 2008, 79, S86.	2.8	1
708	$\langle i \rangle$ PHF11 $\langle  i \rangle$ is not a major candidate gene for asthma or eczema in Chinese children. Pediatric Pulmonology, 2010, 45, 890-897.	2.0	1
709	Fine-scale stratification analysis of Hong Kong Chinese population. , 2010, , .		1
710	PO064 METABOLIC PROFILE OF WOMEN WITH DIABETES TYPE 2 WITH AND WITHOUT A HISTORY OF GESTATIONAL DIABETES MELLITUS – HONG KONG JOINT ASIA DIABETES EVALUATION (JADE). Diabetes Research and Clinical Practice, 2014, 106, S77.	2.8	1
711	Variable selection and prediction of clinical outcome with multiply-imputed data via Bayesian model averaging. , 2016, , .		1
712	Cost of Type 2 Diabetes in Urban Indian Population. Value in Health, 2016, 19, A670.	0.3	1
713	Comparison of glycemic control in Asian and non-Asian T2D patients initiating insulin glargine 100 U/mL as add-on therapy to OADs. Diabetes Research and Clinical Practice, 2016, 120, S114-S115.	2.8	1
714	A patient-centric approach to optimise insulin therapy in Asia. Journal of Diabetes and Its Complications, 2016, 30, 973-980.	2.3	1
715	Augmented Velocity Index - A New Doppler Index to Reflect Arterial Stiffness. Ultrasound in Medicine and Biology, 2017, 43, S245.	1.5	1
716	Category A: Eâ€Poster Presentation: Contraception and Fertility Control. BJOG: an International Journal of Obstetrics and Gynaecology, 2018, 125, 26-28.	2.3	1
717	Within-trial cost-effectiveness of lifestyle intervention using a 3-tier shared care approach for pregnancy outcomes in Chinese women with gestational diabetes. PLoS ONE, 2020, 15, e0237738.	2.5	1
718	Rationale for glycemic control to improve cardiovascular outcome: Lessons from east and west. Journal of Diabetes, 2021, 13, 698-700.	1.8	1
719	Data on diabetes-specific distress are needed to improve the quality of diabetes care – Authors' reply. Lancet, The, 2021, 397, 2150.	13.7	1
720	Frequency of Diabetes-Related Complications in Type 1 and Type 2 Diabetes—Results from the International Diabetes Management Practices Study (IDMPS). Diabetes, 2018, 67, .	0.6	1

#	Article	IF	Citations
721	Methodological challenges to control for immortal time bias in addressing drug effects in type 2 diabetes. World Journal of Methodology, 2015, 5, 122.	3.5	1
722	Integrated Diabetes Care in Hong Kong: From Research to Practice to Policy., 2017,, 65-85.		1
723	Management of People with Diabetes in Africaâ€"Results from the International Diabetes Management Practices Study (IDMPS). Diabetes, 2018, 67, .	0.6	1
724	Neuronal Dysfunction Is Linked to the Famine-Associated Risk of Proliferative Retinopathy in Patients With Type 2 Diabetes. Frontiers in Neuroscience, 2022, $16$ , .	2.8	1
725	Albuminuria, Insulin Resistance and Dyslipidaemia in Chinese Patients with Nonâ€insulinâ€dependent Diabetes (NIDDM). Diabetic Medicine, 1996, 13, 150-155.	2.3	1
726	Predictors and Effects of Participation in Peer Support: A Prospective Structural Equation Modeling Analysis. Annals of Behavioral Medicine, 0, , .	2.9	1
727	Acute Tubular Necrosis Following Endosulfan Insecticide Poisoning: Author's Reply. Journal of Toxicology: Clinical Toxicology, 1995, 33, 377-378.	1.5	O
728	Age-Related Changes in Albumin Adjusted Serum Calcium in Hong Kong Chinese. Annals of Clinical Biochemistry, 1997, 34, 571-573.	1.6	О
729	Evaluation of Losartan Usage in a Regional Hospital in Hong Kong. Journal of Pharmacy Technology, 1999, 15, 170-175.	1.0	O
730	Screening for and treatment of left-ventricular abnormalities in diabetes mellitus. Lancet, The, 2002, 360, 1251-1252.	13.7	0
731	PCV4 VARIABLE PATIENT COMPLIANCE WITH STATINS AND ASSOCIATED LIPID CONTROL AMONG CHINESE PATIENTS WITH HIGH RISK FOR CORONARY HEART DISEASE. Value in Health, 2004, 7, 318-319.	0.3	O
732	PDB7 THE METABOLIC EFFECTS OF ORLISTAT AND ROSIGLITAZONE ON INSULIN ACTION IN A GROUP OF CHINESE PATIENTS AFFECTED BYTHE METABOLIC SYNDROME. Value in Health, 2005, 8, A156.	0.3	О
733	Association Between Obesity and Serum IgE Measurements in a Community Cohort of Chinese Schoolchildren. Journal of Allergy and Clinical Immunology, 2008, 121, S207-S207.	2.9	O
734	CD2-B Emerging treatment for the unmet needs In Type 2 diabetes. Diabetes Research and Clinical Practice, 2008, 79, S8-S9.	2.8	0
735	DWP2-1 Cardio-metabolic-inflammatory risk predictors for heart failure in Type 2 diabetes — the Hong Kong Diabetes Registry. Diabetes Research and Clinical Practice, 2008, 79, S22.	2.8	O
736	CD3-2 Alanine aminotransferase (ALT) levels â€" a marker of cardiovascular risk factors clustering in adolescents. Diabetes Research and Clinical Practice, 2008, 79, S34.	2.8	0
737	Diabetes in Asian Immigrant Populations—Reply. JAMA - Journal of the American Medical Association, 2009, 302, 1646.	7.4	O
738	Response to †Lipid disorders in experimental chronic kidney disease: a role for SREBPs'. Kidney International, 2009, 75, 338-339.	5.2	0

#	Article	IF	CITATIONS
739	ANALYSIS OF SAFETY OF ATORVASTATIN IN ASIAN PATIENTS IN CLINICAL TRIALS. Heart, 2012, 98, E316.3-E316.	2.9	0
740	Type 2 Diabetes. , 2012, , 273-282.		0
741	PO150 PROTECTIVE ROLE OF SIRTUIN 3 (SIRT3) IN HIGH FAT DIET (HFD) INDUCED IMPAIRMENT OF PANCREATIC BETA CELLS' SURVIVAL AND FUNCTIONS. Diabetes Research and Clinical Practice, 2014, 106, S123.	2.8	0
742	PO106 USING NURSES AND INFORMATION TECHNOLOGY TO ENGAGE PATIENTS WITH DIABETES IN THE INITIAL FOLLOW UP PERIOD IMPROVES GLYCAEMIC CONTROL: INSIGHTS FROM THE PHILIPPINES JADE PROGRAM. Diabetes Research and Clinical Practice, 2014, 106, S99.	2.8	0
743	PO408 A QUALITY IMPROVEMENT PROGRAM TO DETECT DEPRESSION USING THE 9-ITEM PATIENT HEALTH QUESTIONNAIRE IN CHINESE WITH TYPE 2 DIABETES. Diabetes Research and Clinical Practice, 2014, 106, S253-S254.	2.8	0
744	PO363 UNRAVELLING PATHWAYS CONTRIBUTING TO THE ASSOCIATION BETWEEN BIRTHWEIGHT AND PREDIABETES: AN ANALYSIS USING STRUCTURAL EQUATION MODELING. Diabetes Research and Clinical Practice, 2014, 106, S232.	2.8	0
745	OP54 EFFECT OF HIGH GLUCOSE ON GENE EXPRESSION IN HCT116 COLORECTAL CANCER CELLS. Diabetes Research and Clinical Practice, 2014, 106, S28.	2.8	0
746	PO124 CLINICAL FACTORS ASSOCIATED WITH SELF-REPORTED HYPOGLYCAEMIA IN VIETNAMESE PATIENTS WITH DIABETES MELLITUS: ANALYSIS OF THE JOINT ASIA DIABETES EVALUATION (JADE) PROGRAM. Diabetes Research and Clinical Practice, 2014, 106, S111.	2.8	0
747	OP34 A MODIFIED RISK EQUATION FOR DEVELOPMENT OF CORONARY HEART DISEASE IN HONG KONG CHINESE WITH TYPE 2 DIABETES. Diabetes Research and Clinical Practice, 2014, 106, S17.	2.8	0
748	OP1 PEER LEADERS ACCELERATED TRAINING INITIATIVE TO UNLEASH POTENTIAL OF MENTORSHIP (PLATINUM) PROGRAM: A 4-YEAR OBSERVATIONAL STUDY ON THE EFFECTS OF PROVIDING PEER SUPPORT IN PEOPLE WITH TYPE 2 DIABETES. Diabetes Research and Clinical Practice, 2014, 106, S1.	2.8	0
749	PO172 PROGRESSION TO TREATMENT FAILURE AMONG CHINESE PATIENTS WITH TYPE 2 DIABETES INITIATED ON METFORMIN VERSUS SULPHONYLUREA MONOTHERAPY – THE HONG KONG DIABETES REGISTRY. Diabetes Research and Clinical Practice, 2014, 106, S135.	2.8	0
750	Impact of Hypoglycaemic Events on Healthcare Resource Use in Type 2 Diabetes Patients. Value in Health, 2016, 19, A683.	0.3	0
751	Impact of Diabetes Management on Diabetes Control and Resource Consumption in Type 2 Diabetic Patients. Value in Health, 2016, 19, A683.	0.3	0
752	Individualized blood pressure targets in Asian diabetic patients with or without nephropathy. Diabetes Research and Clinical Practice, 2016, 120, S12.	2.8	0
753	Effectiveness of peer leaders in diabetes self-management support. Diabetes Research and Clinical Practice, 2016, 120, S28-S29.	2.8	0
754	Risks of progression to end-stage renal disease among type 2 diabetic patients with albuminuric and non-albuminuric chronic kidney disease. Diabetes Research and Clinical Practice, 2016, 120, S46.	2.8	0
755	Association between GWAS-identified variants with CKD in Chinese with Type 2 Diabetes: The Hong Kong Diabetes Registry. Diabetes Research and Clinical Practice, 2016, 120, S49.	2.8	0
756	Genome-wide association study in Chinese identifies new susceptibility loci associated with chronic kidney disease in type 2 diabetes. Diabetes Research and Clinical Practice, 2016, 120, S49-S50.	2.8	0

#	Article	IF	CITATIONS
757	Association between CAG repeat length polymorphism of androgen receptor gene, cardio-metabolic risk factors and clinical outcomes in Chinese men. Diabetes Research and Clinical Practice, 2016, 120, S84-S85.	2.8	O
758	Associations between insomnia and glycemic control in Hong Kong Chinese patients with type 2 diabetes. Diabetes Research and Clinical Practice, 2016, 120, S158.	2.8	0
759	Association between human adenovirus-36 infection, obesity and glycemia in prospective follow-up of Hong Kong Chinese school children. Diabetes Research and Clinical Practice, 2016, 120, S184.	2.8	0
760	Rare variants in ABCG2 identified by sequencing may influence the lipid response to rosuvastatin treatment. Atherosclerosis, 2016, 252, e202-e203.	0.8	0
761	Genetic variants and lipid traits in the Hong Kong Chinese patients with type 2 diabetes. Atherosclerosis, 2016, 252, e148.	0.8	O
762	Severe hypertriglycerideamia related to a novel mutation in TRIB1 and a known variant in APOA5. Atherosclerosis, 2017, 263, e230.	0.8	0
763	Severe familial hypercholesterolaemia in two brothers with three heterozygous coding mutations in LDLR. Atherosclerosis, 2017, 263, e230.	0.8	0
764	Cost-Effectiveness Study To Evaluate Insulin Glargine Compared With NPH Insulin In Patients With Type 2 Diabetes Uncontrolled With Oral Anti-Diabetic Agents In Hong Kong. Value in Health, 2017, 20, A480.	0.3	0
765	Detection of Increased Serum miR-122-5p and miR-455-3p Levels Before the Clinical Diagnosis of Liver Cancer in People With Type 2 Diabetes. SSRN Electronic Journal, 0, , .	0.4	0
766	Association of hip fractures with cardiometabolicâ€renal risk factors in Southern Chinese patients with type 2 diabetes – the Hong Kong Diabetes Register. Journal of Diabetes Investigation, 2021, 12, 1739-1748.	2.4	0
767	IDDF2021-ABS-0138â€Circulating non-coding transcripts serving as biomarkers for diabetic liver steatosis. , 2021, , .		0
768	The diabetic wound healing effect of a two-herb formula and its mechanisms of action. Planta Medica, 2012, 78, .	1.3	0
769	Development of Prediction Models under Multiple Imputation for Coronary Heart Disease in Type 2 Diabetes Mellitus., 2013,,.		0
770	Effect of CYP2C19 *2 and *3 variants on sulphonylurea monotherapy treatment failure in Chinese patients with Type 2 diabetes. Proceedings for Annual Meeting of the Japanese Pharmacological Society, 2018, WCP2018, PO3-14-18.	0.0	0
771	High Serum Branched-Chain Amino Acids Level Independently Predicts Incident Heart Failure—The Hong Kong Diabetes Register. Diabetes, 2018, 67, 455-P.	0.6	0
772	Frequency of High Blood Pressure and Dyslipidemia in T1D and T2Dâ€"Results from the International Diabetes Management Practices Study (IDMPS). Diabetes, 2018, 67, 621-P.	0.6	0
773	Identification of Patient Profiles with Better Glycemic Outcomes Using Machine Learningâ€"Results from the International Diabetes Management Practices Study (IDMPS). Diabetes, 2018, 67, 2398-PUB.	0.6	0
774	Prospective Study on Incidence of Cardiovascular-Renal Diseases, Severe Hypoglycaemia, and Death in Chinese with Latent Autoimmune Diabetes in Adults. Diabetes, 2018, 67, 1590-P.	0.6	0

#	Article	IF	CITATIONS
775	Circulating MicroRNAs Associated with Incident End-Stage Renal Disease in Chinese with Type 2 Diabetes. Diabetes, 2018, 67, .	0.6	O
776	Burden of Hospitalization in Young-Onset Type 2 Diabetes—The Hong Kong Diabetes Register. Diabetes, 2018, 67, 1649-P.	0.6	0
777	Title is missing!. , 2020, 17, e1003316.		O
778	Title is missing!. , 2020, 17, e1003316.		0
779	Title is missing!. , 2020, 17, e1003316.		O
780	Title is missing!. , 2020, 17, e1003316.		0
781	Title is missing!. , 2020, 17, e1003052.		O
782	Title is missing!. , 2020, 17, e1003052.		0
783	Title is missing!. , 2020, 17, e1003052.		O
784	Title is missing!. , 2020, 17, e1003052.		0
785	Title is missing!. , 2020, 17, e1003052.		O
786	Legacy effect of high glucose on promoting survival of HCT116 colorectal cancer cells by reducing endoplasmic reticulum stress response American Journal of Cancer Research, 2021, 11, 6004-6023.	1.4	0
787	Glycated haemoglobin and cardiovascular risk factors in Chinese subjects with normal glucose tolerance. Diabetic Medicine, 1998, 15, 573-578.	2.3	O