

Bo Dai

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7843006/publications.pdf>

Version: 2024-02-01

125
papers

3,321
citations

172457

29
h-index

206112

48
g-index

132
all docs

132
docs citations

132
times ranked

5237
citing authors

#	ARTICLE	IF	CITATIONS
1	Low-cost thermophoretic profiling of extracellular-vesicle surface proteins for the early detection and classification of cancers. <i>Nature Biomedical Engineering</i> , 2019, 3, 183-193.	22.5	324
2	Constitutively Active AR-V7 Plays an Essential Role in the Development and Progression of Castration-Resistant Prostate Cancer. <i>Scientific Reports</i> , 2015, 5, 7654.	3.3	140
3	Identification and Validation of Stromal Immunity Predict Survival and Benefit from Adjuvant Chemotherapy in Patients with Muscle-Invasive Bladder Cancer. <i>Clinical Cancer Research</i> , 2018, 24, 3069-3078.	7.0	124
4	Tumor-associated Macrophage-derived Interleukin-23 Interlinks Kidney Cancer Glutamine Addiction with Immune Evasion. <i>European Urology</i> , 2019, 75, 752-763.	1.9	123
5	Label-Free Isolation and mRNA Detection of Circulating Tumor Cells from Patients with Metastatic Lung Cancer for Disease Diagnosis and Monitoring Therapeutic Efficacy. <i>Analytical Chemistry</i> , 2015, 87, 11893-11900.	6.5	101
6	Acral Melanoma in Chinese: A Clinicopathological and Prognostic Study of 142 cases. <i>Scientific Reports</i> , 2016, 6, 31432.	3.3	92
7	Inactivation of the AMPK-GATA3-ECHS1 Pathway Induces Fatty Acid Synthesis That Promotes Clear Cell Renal Cell Carcinoma Growth. <i>Cancer Research</i> , 2020, 80, 319-333.	0.9	90
8	Retinoic Acid-Related Orphan Receptor C Regulates Proliferation, Glycolysis, and Chemoresistance via the PD-L1/ITGB6/STAT3 Signaling Axis in Bladder Cancer. <i>Cancer Research</i> , 2019, 79, 2604-2618.	0.9	87
9	Intratumoral TIGIT ⁺ CD8 ⁺ T-cell infiltration determines poor prognosis and immune evasion in patients with muscle-invasive bladder cancer. , 2020, 8, e000978.		81
10	Tumor-infiltrating CD39 ⁺ CD8 ⁺ T cells determine poor prognosis and immune evasion in clear cell renal cell carcinoma patients. <i>Cancer Immunology, Immunotherapy</i> , 2020, 69, 1565-1576.	4.2	72
11	Tumor stroma-infiltrating mast cells predict prognosis and adjuvant chemotherapeutic benefits in patients with muscle invasive bladder cancer. <i>Oncimmunology</i> , 2018, 7, e1474317.	4.6	61
12	Blockade of DC-SIGN ⁺ Tumor-Associated Macrophages Reactivates Antitumor Immunity and Improves Immunotherapy in Muscle-Invasive Bladder Cancer. <i>Cancer Research</i> , 2020, 80, 1707-1719.	0.9	61
13	Long non-coding RNA LOC572558 inhibits bladder cancer cell proliferation and tumor growth by regulating the AKT-MDM2-p53 signaling axis. <i>Cancer Letters</i> , 2016, 380, 369-374.	7.2	60
14	MicroRNA-302a Suppresses Tumor Cell Proliferation by Inhibiting AKT in Prostate Cancer. <i>PLoS ONE</i> , 2015, 10, e0124410.	2.5	58
15	Predicting Regional Lymph Node Metastasis in Chinese Patients With Penile Squamous Cell Carcinoma: The Role of Histopathological Classification, Tumor Stage and Depth of Invasion. <i>Journal of Urology</i> , 2006, 176, 1431-1435.	0.4	46
16	Activation of the mammalian target of rapamycin signalling pathway in prostate cancer and its association with patient clinicopathological characteristics. <i>BJU International</i> , 2009, 104, 1009-1016.	2.5	45
17	SPOP promotes ATF2 ubiquitination and degradation to suppress prostate cancer progression. <i>Journal of Experimental and Clinical Cancer Research</i> , 2018, 37, 145.	8.6	43
18	GLUT1 is an AR target contributing to tumor growth and glycolysis in castration-resistant and enzalutamide-resistant prostate cancers. <i>Cancer Letters</i> , 2020, 485, 45-55.	7.2	42

#	ARTICLE	IF	CITATIONS
19	Long noncoding RNA expression signatures of bladder cancer revealed by microarray. <i>Oncology Letters</i> , 2014, 7, 1197-1202.	1.8	41
20	Germline DNA Repair Gene Mutation Landscape in Chinese Prostate Cancer Patients. <i>European Urology</i> , 2019, 76, 280-283.	1.9	41
21	Low TIM3 expression indicates poor prognosis of metastatic prostate cancer and acts as an independent predictor of castration resistant status. <i>Scientific Reports</i> , 2017, 7, 8869.	3.3	40
22	Relationship between circumcision and human papillomavirus infection: a systematic review and meta-analysis. <i>Asian Journal of Andrology</i> , 2017, 19, 125.	1.6	39
23	Primary invasive carcinoma associated with penoscrotal extramammary Paget's disease: a clinicopathological analysis of 56 cases. <i>BJU International</i> , 2015, 115, 153-160.	2.5	36
24	Low pretreatment serum total testosterone is associated with a high incidence of Gleason score 8-10 disease in prostatectomy specimens: data from ethnic Chinese patients with localized prostate cancer. <i>BJU International</i> , 2012, 110, E667-72.	2.5	35
25	CCR8 blockade primes anti-tumor immunity through intratumoral regulatory T cells destabilization in muscle-invasive bladder cancer. <i>Cancer Immunology, Immunotherapy</i> , 2020, 69, 1855-1867.	4.2	35
26	Identification and validation of dichotomous immune subtypes based on intratumoral immune cells infiltration in clear cell renal cell carcinoma patients. , 2020, 8, e000447.		35
27	Tumor-associated macrophages expressing galectin-9 identify immunoevasive subtype muscle-invasive bladder cancer with poor prognosis but favorable adjuvant chemotherapeutic response. <i>Cancer Immunology, Immunotherapy</i> , 2019, 68, 2067-2080.	4.2	34
28	Impact of preoperative 5 α -reductase inhibitors on perioperative blood loss in patients with benign prostatic hyperplasia: a meta-analysis of randomized controlled trials. <i>BMC Urology</i> , 2015, 15, 47.	1.4	33
29	Evaluation of fine particles in surgical smoke from an urologist's operating room by time and by distance. <i>International Urology and Nephrology</i> , 2015, 47, 1671-1678.	1.4	33
30	Tumor Infiltrating Mast Cells (TIMs) Confers a Marked Survival Advantage in Nonmetastatic Clear-Cell Renal Cell Carcinoma. <i>Annals of Surgical Oncology</i> , 2017, 24, 1435-1442.	1.5	33
31	Analysis of KIT expression and gene mutation in human acral melanoma: with a comparison between primary tumors and corresponding metastases/recurrences. <i>Human Pathology</i> , 2013, 44, 1472-1478.	2.0	31
32	Targeting CPT1B as a potential therapeutic strategy in castration-resistant and enzalutamide-resistant prostate cancer. <i>Prostate</i> , 2020, 80, 950-961.	2.3	31
33	Combination of circulating tumor cell enumeration and tumor marker detection in predicting prognosis and treatment effect in metastatic castration-resistant prostate cancer. <i>Oncotarget</i> , 2015, 6, 41825-41836.	1.8	29
34	Identification and validation of an eight-gene expression signature for predicting high Fuhrman grade renal cell carcinoma. <i>International Journal of Cancer</i> , 2017, 140, 1199-1208.	5.1	29
35	Evaluation of ^{99m} Tc-labeled PSMA-SPECT/CT imaging in prostate cancer patients who have undergone biochemical relapse. <i>Asian Journal of Andrology</i> , 2017, 19, 267.	1.6	29
36	Stromal LAG-3 ⁺ cells infiltration defines poor prognosis subtype muscle-invasive bladder cancer with immunoevasive contexture. , 2020, 8, e000651.		29

#	ARTICLE	IF	CITATIONS
37	Basal Cell Carcinoma of the Scrotum: Clinicopathologic Analysis of 10 Cases. <i>Dermatologic Surgery</i> , 2012, 38, 783-790.	0.8	27
38	Phosphorylated 4EBP1 is associated with tumor progression and poor prognosis in Xp11.2 translocation renal cell carcinoma. <i>Scientific Reports</i> , 2016, 6, 23594.	3.3	27
39	Diagnosis of adults Xp11.2 translocation renal cell carcinoma by immunohistochemistry and FISH assays: clinicopathological data from ethnic Chinese population. <i>Scientific Reports</i> , 2016, 6, 21677.	3.3	26
40	Norcantharidin induces autophagy-related prostate cancer cell death through Beclin-1 upregulation by miR-129-5p suppression. <i>Tumor Biology</i> , 2016, 37, 15643-15648.	1.8	26
41	Prognostic value of D-lactate dehydrogenase in patients with clear cell renal cell carcinoma. <i>Oncology Letters</i> , 2018, 16, 866-874.	1.8	26
42	Upregulation of COL6A1 is predictive of poor prognosis in clear cell renal cell carcinoma patients. <i>Oncotarget</i> , 2015, 6, 27378-27387.	1.8	26
43	The Oncogenic Role of COL23A1 in Clear Cell Renal Cell Carcinoma. <i>Scientific Reports</i> , 2017, 7, 9846.	3.3	25
44	Renal cell carcinoma histological subtype distribution differs by age, gender, and tumor size in coastal Chinese patients. <i>Oncotarget</i> , 2017, 8, 71797-71804.	1.8	25
45	Chemokine Receptors CXCR4 and CXCR7 are Associated with Tumor Aggressiveness and Prognosis in Extramammary Paget Disease. <i>Journal of Cancer</i> , 2017, 8, 2471-2477.	2.5	22
46	PD-L1 expression in tumour-infiltrating lymphocytes is a poor prognostic factor for primary acral melanoma patients. <i>Histopathology</i> , 2018, 73, 386-396.	2.9	22
47	SOX2 and SOX12 are predictive of prognosis in patients with clear cell renal cell carcinoma. <i>Oncology Letters</i> , 2018, 15, 4564-4570.	1.8	22
48	Intratumoral IL22-producing cells define immunoevasive subtype muscle-invasive bladder cancer with poor prognosis and superior nivolumab responses. <i>International Journal of Cancer</i> , 2020, 146, 542-552.	5.1	22
49	Spiradenocarcinoma, cylindrocarcinoma and spiradenocylindrocarcinoma: a clinicopathological study of nine cases. <i>Histopathology</i> , 2014, 65, 658-666.	2.9	21
50	PD-L1 expression in Xp11.2 translocation renal cell carcinoma: Indicator of tumor aggressiveness. <i>Scientific Reports</i> , 2017, 7, 2074.	3.3	21
51	Serum testosterone level predicts the effective time of androgen deprivation therapy in metastatic prostate cancer patients. <i>Asian Journal of Andrology</i> , 2017, 19, 178.	1.6	21
52	CHEK2 mutation and risk of prostate cancer: a systematic review and meta-analysis. <i>International Journal of Clinical and Experimental Medicine</i> , 2015, 8, 15708-15.	1.3	21
53	Prognostic significance of the TREK-1 K2P potassium channels in prostate cancer. <i>Oncotarget</i> , 2015, 6, 18460-18468.	1.8	20
54	Expression of Dicer and Its Related MiRNAs in the Progression of Prostate Cancer. <i>PLoS ONE</i> , 2015, 10, e0120159.	2.5	19

#	ARTICLE	IF	CITATIONS
55	Individualized prostate biopsy strategy for Chinese patients with different prostate-specific antigen levels. <i>Asian Journal of Andrology</i> , 2008, 10, 325-331.	1.6	18
56	Serum Adiponectin Level May be an Independent Predictor of Clear Cell Renal Cell Carcinoma. <i>Journal of Cancer</i> , 2016, 7, 1340-1346.	2.5	18
57	Pathological Features of Localized Prostate Cancer in China: A Contemporary Analysis of Radical Prostatectomy Specimens. <i>PLoS ONE</i> , 2015, 10, e0121076.	2.5	18
58	Comprehensive Analysis of <i>BAP1</i> Somatic Mutation in Clear Cell Renal Cell Carcinoma to Explore Potential Mechanisms <i>in Silico</i> . <i>Journal of Cancer</i> , 2018, 9, 4108-4116.	2.5	17
59	Extended versus non-extended lymphadenectomy during radical cystectomy for patients with bladder cancer: a meta-analysis of the effect on long-term and short-term outcomes. <i>World Journal of Surgical Oncology</i> , 2019, 17, 225.	1.9	17
60	Prognostic Value of Germline DNA Repair Gene Mutations in De Novo Metastatic and Castration-Sensitive Prostate Cancer. <i>Oncologist</i> , 2020, 25, e1042-e1050.	3.7	17
61	Forkhead box series expression network is associated with outcome of clear cell renal cell carcinoma. <i>Oncology Letters</i> , 2018, 15, 8669-8680.	1.8	16
62	Development and External Validation of a Novel 12-Gene Signature for Prediction of Overall Survival in Muscle-Invasive Bladder Cancer. <i>Frontiers in Oncology</i> , 2019, 9, 856.	2.8	16
63	Phenotypes of circulating tumour cells predict time to castration resistance in metastatic castration-sensitive prostate cancer. <i>BJU International</i> , 2019, 124, 258-267.	2.5	16
64	Assessment of survival of patients with metastatic clear cell renal cell carcinoma after radical cytoreductive nephrectomy versus no surgery: a SEER analysis. <i>International Braz J Urol: Official Journal of the Brazilian Society of Urology</i> , 2015, 41, 288-295.	1.5	15
65	Prognostic factors in primary anorectal melanoma: a clinicopathological study of 60 cases in China. <i>Human Pathology</i> , 2018, 79, 77-85.	2.0	15
66	B4GALT1 expression predicts prognosis and adjuvant chemotherapy benefits in muscle-invasive bladder cancer patients. <i>BMC Cancer</i> , 2018, 18, 590.	2.6	15
67	CCR5 blockade inflames antitumor immunity in BAP1-mutant clear cell renal cell carcinoma. , 2020, 8, e000228.		15
68	The Value of 99mTc-PSMA SPECT/CT-Guided Surgery for Identifying and Locating Lymph Node Metastasis in Prostate Cancer Patients. <i>Annals of Surgical Oncology</i> , 2019, 26, 653-659.	1.5	14
69	Poliovirus receptor CD155 is up-regulated in muscle-invasive bladder cancer and predicts poor prognosis. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2020, 38, 41.e11-41.e18.	1.6	14
70	NR1H3 Expression is a Prognostic Factor of Overall Survival for Patients with Muscle-Invasive Bladder Cancer. <i>Journal of Cancer</i> , 2017, 8, 852-860.	2.5	13
71	Expression of ARID1B Is Associated With Poor Outcomes and Predicts the Benefit from Adjuvant Chemotherapy in Bladder Urothelial Carcinoma. <i>Journal of Cancer</i> , 2017, 8, 3490-3497.	2.5	13
72	Identification and validation of poor prognosis immunoevasive subtype of muscle-invasive bladder cancer with tumor-infiltrating podoplanin ⁺ cell abundance. <i>Oncolmmunology</i> , 2020, 9, 1747333.	4.6	13

#	ARTICLE	IF	CITATIONS
73	Inherited Mutations in Chinese Men With Prostate Cancer. <i>Journal of the National Comprehensive Cancer Network: JNCCN</i> , 2022, 20, 54-62.	4.9	13
74	Preferentially expressed antigen in melanoma immunohistochemistry as an adjunct for differential diagnosis in acral lentiginous melanoma and acral nevi. <i>Human Pathology</i> , 2022, 120, 9-17.	2.0	13
75	Functional variants of the 5-methyltetrahydrofolate-homocysteine methyltransferase gene significantly increase susceptibility to prostate cancer: Results from an ethnic Han Chinese population. <i>Scientific Reports</i> , 2016, 6, 36264.	3.3	12
76	Low serum prostate-specific antigen level predicts poor outcomes in patients with primary neuroendocrine prostate cancer. <i>Prostate</i> , 2019, 79, 1563-1571.	2.3	12
77	PBRM1 regulates proliferation and the cell cycle in renal cell carcinoma through a chemokine/chemokine receptor interaction pathway. <i>PLoS ONE</i> , 2017, 12, e0180862.	2.5	12
78	High NUCB2 expression level represents an independent negative prognostic factor in Chinese cohorts of non-metastatic clear cell renal cell carcinoma patients. <i>Oncotarget</i> , 2017, 8, 35244-35254.	1.8	11
79	NUDT expression is predictive of prognosis in patients with clear cell renal cell carcinoma. <i>Oncology Letters</i> , 2017, 14, 6121-6128.	1.8	10
80	Identification and validation of an 18-gene signature highly-predictive of bladder cancer metastasis. <i>Scientific Reports</i> , 2018, 8, 374.	3.3	10
81	PAK1 expression determines poor prognosis and immune evasion in metastatic renal cell carcinoma patients. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2020, 38, 293-304.	1.6	10
82	Conditional survival among patients with adrenal cortical carcinoma determined using a national population-based surveillance, epidemiology, and end results registry. <i>Oncotarget</i> , 2015, 6, 44955-44962.	1.8	10
83	National Comprehensive Cancer Network (NCCN) risk classification in predicting biochemical recurrence after radical prostatectomy: a retrospective cohort study in Chinese prostate cancer patients. <i>Asian Journal of Andrology</i> , 2018, 20, 551.	1.6	9
84	Optical biopsy of bladder cancer using confocal laser endomicroscopy. <i>International Urology and Nephrology</i> , 2019, 51, 1473-1479.	1.4	9
85	Development and validation of a robust multigene signature as an aid to predict early relapse in stage I-III clear cell and papillary renal cell cancer. <i>Journal of Cancer</i> , 2020, 11, 997-1007.	2.5	9
86	Clinicopathological, immunohistochemical and fluorescence in situ hybridisation features of early subungual melanoma: an analysis of 65 cases. <i>Histopathology</i> , 2021, 78, 717-726.	2.9	9
87	Prognostic value of granulocyte colony-stimulating factor in patients with non-metastatic clear cell renal cell carcinoma. <i>Oncotarget</i> , 2017, 8, 69961-69971.	1.8	9
88	Factors influencing biochemical recurrence in patients who have received salvage radiotherapy after radical prostatectomy: a systematic review and meta-analysis. <i>Asian Journal of Andrology</i> , 2017, 19, 493.	1.6	9
89	External validation and newly development of a nomogram to predict overall survival of abiraterone-treated, castration-resistant patients with metastatic prostate cancer. <i>Asian Journal of Andrology</i> , 2018, 20, 184.	1.6	9
90	Human epidermal growth factor receptor type 2 protein expression in Chinese metastatic prostate cancer patients correlates with cancer specific survival and increases after exposure to hormonal therapy. <i>Asian Journal of Andrology</i> , 2008, 10, 701-709.	1.6	8

#	ARTICLE	IF	CITATIONS
91	Prognosis of rare pathological primary urethral carcinoma. <i>Cancer Management and Research</i> , 2018, Volume 10, 6815-6822.	1.9	8
92	Poor clinical outcomes and immunoevasive contexture in interleukin-9 abundant muscle-invasive bladder cancer. <i>International Journal of Cancer</i> , 2020, 147, 3539-3549.	5.1	8
93	Intravesical Recurrence After Radical Nephroureterectomy of Upper Urinary Tract Urothelial Carcinoma: A Large Population-Based Investigation of Clinicopathologic Characteristics and Survival Outcomes. <i>Frontiers in Surgery</i> , 2021, 8, 590448.	1.4	8
94	Oral etoposide and oral prednisone for the treatment of castration resistant prostate cancer. <i>Kaohsiung Journal of Medical Sciences</i> , 2014, 30, 82-85.	1.9	7
95	MTHFR c.677C>T Inhibits Cell Proliferation and Decreases Prostate Cancer Susceptibility in the Han Chinese Population in Shanghai. <i>Scientific Reports</i> , 2016, 6, 36290.	3.3	7
96	Identification and validation of an excellent prognosis subtype of muscle-invasive bladder cancer patients with intratumoral CXCR5 ⁺ CD8 ⁺ T cell abundance. <i>Oncolmmunology</i> , 2020, 9, 1810489.	4.6	7
97	Association of glutathione S-transferase T1 and M1 polymorphisms with prostate cancer susceptibility in populations of Asian descent: a meta-analysis. <i>Oncotarget</i> , 2015, 6, 35843-35850.	1.8	7
98	Outcomes of patients with lymph node metastasis treated with radical prostatectomy and adjuvant androgen deprivation therapy in a Chinese population: results from a cohort study. <i>World Journal of Surgical Oncology</i> , 2015, 13, 172.	1.9	6
99	High CXC chemokine receptor 1 level represents an independent negative prognosticator in non-metastatic clear-cell renal cell carcinoma patients. <i>Oncolmmunology</i> , 2017, 6, e1359450.	4.6	6
100	The Prognostic Value of Programmed Death-Ligand 1 in a Chinese Cohort With Clear Cell Renal Cell Carcinoma. <i>Frontiers in Oncology</i> , 2019, 9, 879.	2.8	6
101	Prognostic Value of Local Treatment in Prostate Cancer Patients With Different Metastatic Sites: A Population Based Retrospective Study. <i>Frontiers in Oncology</i> , 2020, 10, 527952.	2.8	6
102	Tumor-infiltrating IL-17A ⁺ cells determine favorable prognosis and adjuvant chemotherapeutic response in muscle-invasive bladder cancer. <i>Oncolmmunology</i> , 2020, 9, 1747332.	4.6	6
103	Kinetics of testosterone recovery in clinically localized prostate cancer patients treated with radical prostatectomy and subsequent short-term adjuvant androgen deprivation therapy. <i>Asian Journal of Andrology</i> , 2013, 15, 466-470.	1.6	6
104	Eosinophil percentage elevation as a prognostic factor for overall survival in patients with metastatic renal cell carcinoma treated with tyrosine kinase inhibitor. <i>Oncotarget</i> , 2016, 7, 68943-68953.	1.8	6
105	Clinical activity of abiraterone plus prednisone in docetaxel-naïve and docetaxel-resistant Chinese patients with metastatic castration-resistant prostate cancer. <i>Asian Journal of Andrology</i> , 2019, 21, 131.	1.6	6
106	Presence of CD133 ⁺ positive circulating tumor cells predicts worse progression-free survival in patients with metastatic castration-sensitive prostate cancer. <i>International Journal of Urology</i> , 2022, 29, 383-389.	1.0	6
107	Retrograde radical cystectomy and consequent peritoneal cavity reconstruction benefits localized male bladder cancer: results from a cohort study. <i>World Journal of Surgical Oncology</i> , 2015, 13, 132.	1.9	5
108	Evaluation of the major changes in eighth edition of the American Joint Committee on Cancer pathological staging for prostate cancer treated with prostatectomy. <i>PLoS ONE</i> , 2017, 12, e0187887.	2.5	5

#	ARTICLE	IF	CITATIONS
109	Evaluation of clinical staging of the American Joint Committee on Cancer (eighth edition) for prostate cancer. <i>World Journal of Urology</i> , 2018, 36, 769-774.	2.2	5
110	Diagnostic Performance of Confocal Laser Endomicroscopy for the Detection of Bladder Cancer: Systematic Review and Meta-Analysis. <i>Urologia Internationalis</i> , 2020, 104, 523-532.	1.3	5
111	The Effect of an Information Support Program on Self-Efficacy of Prostate Cancer Patients during Hormonal Therapy. <i>Asia-Pacific Journal of Oncology Nursing</i> , 2021, 8, 639-652.	1.6	5
112	Clinical significance of TMPRSS4 in prostate cancer. <i>International Journal of Clinical and Experimental Pathology</i> , 2014, 7, 8053-8.	0.5	5
113	PTEN genomic deletion defines favorable prognostic biomarkers in localized prostate cancer: a systematic review and meta-analysis. <i>International Journal of Clinical and Experimental Medicine</i> , 2015, 8, 5430-7.	1.3	5
114	High IL-23+ cells infiltration correlates with worse clinical outcomes and abiraterone effectiveness in patients with prostate cancer. <i>Asian Journal of Andrology</i> , 2022, 24, 147.	1.6	5
115	Prognostic and Predictive Value of O6-methylguanine Methyltransferase for Chemotherapy in Patients with Muscle-Invasive Bladder Cancer. <i>Annals of Surgical Oncology</i> , 2018, 25, 342-348.	1.5	4
116	Intratumoral CCR5 ⁺ neutrophils identify immunogenic subtype muscle-invasive bladder cancer with favorable prognosis and therapeutic responses. <i>Oncolmmunology</i> , 2020, 9, 1802176.	4.6	4
117	Development of a preliminary nomogram to predict progression of bone scan for castration-resistant prostate cancer. <i>OncoTargets and Therapy</i> , 2015, 8, 713.	2.0	3
118	PCA3 rs544190G>A and prostate cancer risk in an eastern Chinese population. <i>International Braz J Urol: Official Journal of the Brazilian Society of Urology</i> , 2018, 44, 500-505.	1.5	3
119	Development and validation of a mitochondrial metabolism-associated nomogram for prediction of prognosis in clear cell renal cell carcinoma. <i>Clinical and Translational Medicine</i> , 2020, 10, e120.	4.0	3
120	GPR160 is a potential biomarker associated with prostate cancer. <i>Signal Transduction and Targeted Therapy</i> , 2021, 6, 241.	17.1	3
121	Prognostic value of PTEN in de novo diagnosed metastatic prostate cancer. <i>Asian Journal of Andrology</i> , 2022, 24, 50.	1.6	2
122	A Germline Variant at 8q24 Contributes to the Serum p2PSA Level in a Chinese Prostate Biopsy Cohort. <i>Frontiers in Oncology</i> , 2021, 11, 753920.	2.8	2
123	High expression of F2RL3 correlates with aggressive features and poor survival in clear cell renal cell carcinoma. <i>Journal of Cancer</i> , 2018, 9, 3400-3406.	2.5	1
124	Prognostic Value of an Immunohistochemical Signature in Patients With Bladder Cancer Undergoing Radical Cystectomy. <i>Frontiers in Oncology</i> , 2021, 11, 641385.	2.8	1
125	Prognosis of the Metachronous and Synchronous Bilateral Renal Cancer and Second Primary Cancer After the Bilateral Renal Cancer: a Population-Based Analysis. <i>SN Comprehensive Clinical Medicine</i> , 2019, 1, 900-904.	0.6	0