

Pedro F Mateos

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7960624/publications.pdf>

Version: 2024-02-01

130
papers

6,552
citations

53794

45
h-index

71685

76
g-index

141
all docs

141
docs citations

141
times ranked

4516
citing authors

#	ARTICLE	IF	CITATIONS
1	Yield response of common bean to co-inoculation with Rhizobium and Pseudomonas endophytes and microscopic evidence of different colonised spaces inside the nodule. <i>European Journal of Agronomy</i> , 2021, 122, 126187.	4.1	15
2	Identification of Canola Roots Endophytic Bacteria and Analysis of Their Potential as Biofertilizers for Canola Crops with Special Emphasis on Sporulating Bacteria. <i>Agronomy</i> , 2021, 11, 1796.	3.0	15
3	Role of QseG membrane protein in beneficial enterobacterial interactions with plants and <i>Mesorhizobia</i> . <i>Journal of Plant Interactions</i> , 2021, 16, 510-521.	2.1	2
4	Selection of the Root Endophyte <i>Pseudomonas brassicacearum</i> CDVBN10 as Plant Growth Promoter for <i>Brassica napus</i> L. <i>Crops. Agronomy</i> , 2020, 10, 1788.	3.0	24
5	Búsqueda de bacterias productoras de antibióticos a partir del culturoma rizosférico. <i>FarmaJournal</i> , 2020, 5, 43-50.	0.0	0
6	The N-fixing legume <i>Periandra mediterranea</i> constrains the invasion of an exotic grass (<i>Melinis</i>) in a semi-arid region. <i>Journal of Ecology</i> , 2020, 108, 1010-1020.	3.3	10
7	Legumes display common and host-specific responses to the rhizobial cellulase CelC2 during primary symbiotic infection. <i>Scientific Reports</i> , 2019, 9, 13907.	3.3	8
8	Genome Insights into the Novel Species <i>Microvirga brassicacearum</i> , a Rapeseed Endophyte with Biotechnological Potential. <i>Microorganisms</i> , 2019, 7, 354.	3.6	30
9	Heterologous expression of <i>nifA</i> or <i>nodD</i> genes improves chickpea- <i>Mesorhizobium</i> symbiotic performance. <i>Plant and Soil</i> , 2019, 436, 607-621.	3.7	7
10	<i>Phaseolus vulgaris</i> is nodulated by the symbiovar <i>viciae</i> of several genospecies of <i>Rhizobium laguerreae</i> complex in a Spanish region where <i>Lens culinaris</i> is the traditionally cultivated legume. <i>Systematic and Applied Microbiology</i> , 2019, 42, 240-247.	2.8	22
11	Heterologous Expression of Rhizobial CelC2 Cellulase Impairs Symbiotic Signaling and Nodulation in <i>Medicago truncatula</i> . <i>Molecular Plant-Microbe Interactions</i> , 2018, 31, 568-575.	2.6	9
12	Probiotic activities of <i>Rhizobium laguerreae</i> on growth and quality of spinach. <i>Scientific Reports</i> , 2018, 8, 295.	3.3	50
13	A conserved ϵ -proteobacterial small RNA contributes to osmoadaptation and symbiotic efficiency of rhizobia on legume roots. <i>Environmental Microbiology</i> , 2017, 19, 2661-2680.	3.8	27
14	<i>Mesorhizobium</i> bacterial strains isolated from the legume <i>Lotus corniculatus</i> are an alternative source for the production of polyhydroxyalkanoates (PHAs) to obtain bioplastics. <i>Environmental Science and Pollution Research</i> , 2017, 24, 17436-17445.	5.3	5
15	Recent Advances in the Active Biomolecules Involved in Rhizobia-Legume Symbiosis. , 2017, , 45-74.		7
16	Bacterial Probiotics: A Truly Green Revolution. , 2017, , 131-162.		14
17	Invasion of the Brazilian campo rupestre by the exotic grass <i>Melinis minutiflora</i> is driven by the high soil N availability and changes in the N cycle. <i>Science of the Total Environment</i> , 2017, 577, 202-211.	8.0	24
18	<i>Bacillus terrae</i> sp. nov. isolated from <i>Cistus ladanifer</i> rhizosphere soil. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2017, 67, 1478-1481.	1.7	12

#	ARTICLE	IF	CITATIONS
19	Mesorhizobium helmanticense sp. nov., isolated from Lotus corniculatus nodules. International Journal of Systematic and Evolutionary Microbiology, 2017, 67, 2301-2305.	1.7	21
20	Analysis and effect of the use of biofertilizers on <i>Trifolium rubens</i> L., a preferential attention species in Castile and Leon, Spain, with the aim of increasing the plants conservation status. AIMS Microbiology, 2017, 3, 733-746.	2.2	7
21	Rhizobium Symbiotic Enzyme Cellulase CelC2: Properties and Applications. , 2016, , 81-89.		2
22	The Symbiotic Performance of Chickpea Rhizobia Can Be Improved by Additional Copies of the clpB Chaperone Gene. PLoS ONE, 2016, 11, e0148221.	2.5	14
23	Rhizobial Biofertilizers for Ornamental Plants. , 2016, , 13-21.		3
24	Analysis of the PGPB Potential of Bacterial Endophytes Associated with Maize. , 2016, , 23-35.		5
25	Effective Colonization of Spinach Root Surface by Rhizobium. , 2016, , 109-122.		8
26	Paenibacillus periandrae sp. nov., isolated from nodules of Periandra mediterranea. International Journal of Systematic and Evolutionary Microbiology, 2016, 66, 1838-1843.	1.7	16
27	Plants Probiotics as a Tool to Produce Highly Functional Fruits: The Case of Phyllobacterium and Vitamin C in Strawberries. PLoS ONE, 2015, 10, e0122281.	2.5	106
28	Pseudomonas coleopterorum sp. nov., a cellulase-producing bacterium isolated from the bark beetle Hylesinus fraxini. International Journal of Systematic and Evolutionary Microbiology, 2015, 65, 2852-2858.	1.7	50
29	Rhizobium cellulosityticum as a co-inoculant enhances Phaseolus vulgaris grain yield under greenhouse conditions. Symbiosis, 2015, 67, 135-141.	2.3	11
30	Rhizobium as plant probiotic for strawberry production under microcosm conditions. Symbiosis, 2015, 67, 25-32.	2.3	18
31	The high diversity of Lotus corniculatus endosymbionts in soils of northwest Spain. Symbiosis, 2015, 67, 11-20.	2.3	16
32	Revision of the taxonomic status of the species Rhizobium lupini and reclassification as Bradyrhizobium lupini comb. nov.. International Journal of Systematic and Evolutionary Microbiology, 2015, 65, 1213-1219.	1.7	52
33	Cicer canariense, an endemic legume to the Canary Islands, is nodulated in mainland Spain by fast-growing strains from symbiovar trifolii phylogenetically related to Rhizobium leguminosarum. Systematic and Applied Microbiology, 2015, 38, 346-350.	2.8	8
34	Revision of the taxonomic status of type strains of Mesorhizobium loti and reclassification of strain USDA 3471T as the type strain of Mesorhizobium erdmanii sp. nov. and ATCC 33669T as the type strain of Mesorhizobium jarvisii sp. nov.. International Journal of Systematic and Evolutionary Microbiology, 2015, 65, 1703-1708.	1.7	47
35	MALDI-TOF mass spectrometry as a tool for differentiation of Bradyrhizobium species: Application to the identification of Lupinus nodulating strains. Systematic and Applied Microbiology, 2013, 36, 565-571.	2.8	21
36	Use of <i>Rhizobium leguminosarum</i> as a potential biofertilizer for <i>Lactuca sativa</i> and <i>Daucus carota</i> crops. Journal of Plant Nutrition and Soil Science, 2013, 176, 876-882.	1.9	99

#	ARTICLE	IF	CITATIONS
37	A ClpB Chaperone Knockout Mutant of <i>Mesorhizobium ciceri</i> Shows a Delay in the Root Nodulation of Chickpea Plants. <i>Molecular Plant-Microbe Interactions</i> , 2012, 25, 1594-1604.	2.6	23
38	Role of Rhizobium endoglucanase CelC2 in cellulose biosynthesis and biofilm formation on plant roots and abiotic surfaces. <i>Microbial Cell Factories</i> , 2012, 11, 125.	4.0	86
39	Rhizobium Promotes Non-Legumes Growth and Quality in Several Production Steps: Towards a Biofertilization of Edible Raw Vegetables Healthy for Humans. <i>PLoS ONE</i> , 2012, 7, e38122.	2.5	155
40	Nodulation in <i>Dimorphandra wilsonii</i> Rizz. (Caesalpinioideae), a Threatened Species Native to the Brazilian Cerrado. <i>PLoS ONE</i> , 2012, 7, e49520.	2.5	38
41	Development of Functional Symbiotic White Clover Root Hairs and Nodules Requires Tightly Regulated Production of Rhizobial Cellulase CelC2. <i>Molecular Plant-Microbe Interactions</i> , 2011, 24, 798-807.	2.6	31
42	The celC gene, a new phylogenetic marker useful for taxonomic studies in Rhizobium. <i>Systematic and Applied Microbiology</i> , 2011, 34, 393-399.	2.8	13
43	MALDI-TOF Mass Spectrometry Is a Fast and Reliable Platform for Identification and Ecological Studies of Species from Family Rhizobiaceae. <i>PLoS ONE</i> , 2011, 6, e20223.	2.5	94
44	Strains nodulating <i>Lupinus albus</i> on different continents belong to several new chromosomal and symbiotic lineages within Bradyrhizobium. <i>Antonie Van Leeuwenhoek</i> , 2010, 97, 363-376.	1.7	48
45	Analysis of core genes supports the reclassification of strains <i>Agrobacterium radiobacter</i> K84 and <i>Agrobacterium tumefaciens</i> AKE10 into the species <i>Rhizobium rhizogenes</i> . <i>Systematic and Applied Microbiology</i> , 2010, 33, 247-251.	2.8	48
46	Key Molecules Involved in Beneficial Infection Process in Rhizobia—“Legume Symbiosis.”, 2010, , 55-80.		7
47	<i>Bradyrhizobium pachyrhizi</i> sp. nov. and <i>Bradyrhizobium jicamae</i> sp. nov., isolated from effective nodules of <i>Pachyrhizus erosus</i> . <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2009, 59, 1929-1934.	1.7	127
48	Rhizobia from Lanzarote, the Canary Islands, That Nodulate <i>Phaseolus vulgaris</i> Have Characteristics in Common with <i>Sinorhizobium meliloti</i> Isolates from Mainland Spain. <i>Applied and Environmental Microbiology</i> , 2009, 75, 2354-2359.	3.1	40
49	<i>Acinetobacter</i> strains IH9 and OC11, two rhizospheric phosphate solubilizing isolates able to promote plant growth, constitute a new genomovar of <i>Acinetobacter calcoaceticus</i> . <i>Systematic and Applied Microbiology</i> , 2009, 32, 334-341.	2.8	20
50	The analysis of core and symbiotic genes of rhizobia nodulating <i>Vicia</i> from different continents reveals their common phylogenetic origin and suggests the distribution of <i>Rhizobium leguminosarum</i> strains together with <i>Vicia</i> seeds. <i>Archives of Microbiology</i> , 2009, 191, 659-668.	2.2	49
51	Phenotypic, genotypic, and symbiotic diversities in strains nodulating clover in different soils in Spain. <i>Canadian Journal of Microbiology</i> , 2009, 55, 1207-1216.	1.7	25
52	Stable low molecular weight RNA profiling showed variations within <i>Sinorhizobium meliloti</i> and <i>Sinorhizobium medicae</i> nodulating different legumes from the alfalfa cross-inoculation group. <i>FEMS Microbiology Letters</i> , 2008, 282, 273-281.	1.8	10
53	Revision of the taxonomic status of the species <i>Rhizobium leguminosarum</i> (Frank 1879) Frank 1889AL, <i>Rhizobium phaseoli</i> Dangeard 1926AL and <i>Rhizobium trifolii</i> Dangeard 1926AL. <i>R. trifolii</i> is a later synonym of <i>R. leguminosarum</i> . Reclassification of the strain <i>R. leguminosarum</i> DSM 30132 (=NCIMB) Tj ETQq1 1 01784314 rgs1 /Overle 2008, 58, 2484-2490.	2.8	151
54	<i>Cohnella phaseoli</i> sp. nov., isolated from root nodules of <i>Phaseolus coccineus</i> in Spain, and emended description of the genus <i>Cohnella</i> . <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2008, 58, 1855-1859.	1.7	67

#	ARTICLE	IF	CITATIONS
55	<i>Saccharibacillus sacchari</i> gen. nov., sp. nov., isolated from sugar cane. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2008, 58, 1850-1854.	1.7	35
56	<i>Rhizobium</i> cellulase CelC2 is essential for primary symbiotic infection of legume host roots. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2008, 105, 7064-7069.	7.1	119
57	<i>Alcanivorax balearicus</i> sp. nov., isolated from Lake Martel. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2007, 57, 1331-1335.	1.7	35
58	<i>Ochrobactrum cytisi</i> sp. nov., isolated from nodules of <i>Cytisus scoparius</i> in Spain. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2007, 57, 784-788.	1.7	138
59	Reclassification of <i>Pseudomonas aurantiaca</i> as a synonym of <i>Pseudomonas chlororaphis</i> and proposal of three subspecies, <i>P. chlororaphis</i> subsp. <i>chlororaphis</i> subsp. nov., <i>P. chlororaphis</i> subsp. <i>aureofaciens</i> subsp. nov., comb. nov. and <i>P. chlororaphis</i> subsp. <i>aurantiaca</i> subsp. nov., comb. nov.. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2007, 57, 1286-1290.	1.7	99
60	Characterization of a strain of <i>Pseudomonas fluorescens</i> that solubilizes phosphates in vitro and produces high antibiotic activity against several microorganisms. , 2007, , 265-268.		4
61	Two strains isolated from tumours of <i>Prunus persica</i> are able to solubilize phosphate in vitro. , 2007, , 347-349.		1
62	Phosphate solubilizing rhizobia originating from <i>Medicago</i> , <i>Melilotus</i> and <i>Trigonella</i> grown in a Spanish soil. , 2007, , 149-156.		3
63	Biodiversity of populations of phosphate solubilizing rhizobia that nodulates chickpea in different Spanish soils. , 2007, , 23-33.		8
64	Characterization of xylanolytic bacteria present in the bract phyllosphere of the date palm <i>Phoenix dactylifera</i> . <i>Letters in Applied Microbiology</i> , 2007, 44, 181-187.	2.2	97
65	Strains of <i>Mesorhizobium amorphae</i> and <i>Mesorhizobium tianshanense</i> , carrying symbiotic genes of common chickpea endosymbiotic species, constitute a novel biovar (<i>ciceri</i>) capable of nodulating <i>Cicer arietinum</i> . <i>Letters in Applied Microbiology</i> , 2007, 44, 412-418.	2.2	92
66	Genetic characterization of fast-growing rhizobia able to nodulate <i>Prosopis alba</i> in North Spain. <i>FEMS Microbiology Letters</i> , 2007, 277, 210-216.	1.8	40
67	<i>Rhizobium cellulosityticum</i> sp. nov., isolated from sawdust of <i>Populus alba</i> . <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2007, 57, 844-848.	1.7	80
68	Biodiversity of populations of phosphate solubilizing rhizobia that nodulates chickpea in different Spanish soils. <i>Plant and Soil</i> , 2006, 287, 23-33.	3.7	104
69	A new approach for separating low-molecular-weight RNA molecules by staircase electrophoresis in non-sequencing gels. <i>Electrophoresis</i> , 2006, 27, 1732-1738.	2.4	1
70	<i>Photobacterium halotolerans</i> sp. nov., isolated from Lake Martel in Spain. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2006, 56, 1067-1071.	1.7	37
71	<i>Paenibacillus cellulosityticus</i> sp. nov., a cellulolytic and xylanolytic bacterium isolated from the bract phyllosphere of <i>Phoenix dactylifera</i> . <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2006, 56, 2777-2781.	1.7	46
72	The Coexistence of Symbiosis and Pathogenicity-Determining Genes in <i>Rhizobium rhizogenes</i> Strains Enables Them to Induce Nodules and Tumors or Hairy Roots in Plants. <i>Molecular Plant-Microbe Interactions</i> , 2005, 18, 1325-1332.	2.6	71

#	ARTICLE	IF	CITATIONS
73	Application of horizontal staircase electrophoresis in agarose minigels to the random intergenic spacer analysis of clinical samples. <i>Electrophoresis</i> , 2005, 26, 4402-4410.	2.4	13
74	<i>Marteella mediterranea</i> gen. nov., sp. nov., a novel β -proteobacterium isolated from a subterranean saline lake. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2005, 55, 955-959.	1.7	46
75	<i>Micromonospora mirobrigensis</i> sp. nov.. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2005, 55, 877-880.	1.7	66
76	<i>Paenibacillus rhizosphaerae</i> sp. nov., isolated from the rhizosphere of <i>Cicer arietinum</i> . <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2005, 55, 1305-1309.	1.7	28
77	<i>Phyllobacterium trifolii</i> sp. nov., nodulating <i>Trifolium</i> and <i>Lupinus</i> in Spanish soils. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2005, 55, 1985-1989.	1.7	143
78	<i>Paenibacillus xylanilyticus</i> sp. nov., an airborne xylanolytic bacterium. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2005, 55, 405-408.	1.7	65
79	Nodulation of <i>Lupinus albus</i> by Strains of <i>Ochrobactrum lupini</i> sp. nov. <i>Applied and Environmental Microbiology</i> , 2005, 71, 1318-1327.	3.1	219
80	<i>Paenibacillus phyllosphaerae</i> sp. nov., a xylanolytic bacterium isolated from the phyllosphere of <i>Phoenix dactylifera</i> . <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2005, 55, 743-746.	1.7	54
81	<i>Xylanibacterium ulmi</i> gen. nov., sp. nov., a novel xylanolytic member of the family Promicromonosporaceae. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2004, 54, 557-561.	1.7	38
82	<i>Bradyrhizobium betae</i> sp. nov., isolated from roots of <i>Beta vulgaris</i> affected by tumour-like deformations. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2004, 54, 1271-1275.	1.7	115
83	<i>Pseudomonas lutea</i> sp. nov., a novel phosphate-solubilizing bacterium isolated from the rhizosphere of grasses. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2004, 54, 847-850.	1.7	59
84	<i>Cellulomonas xylanilytica</i> sp. nov., a cellulolytic and xylanolytic bacterium isolated from a decayed elm tree. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2004, 54, 533-536.	1.7	43
85	<i>Agromyces ulmi</i> sp. nov., a xylanolytic bacterium isolated from <i>Ulmus nigra</i> in Spain. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2004, 54, 1987-1990.	1.7	40
86	<i>Mycobacterium psychrotolerans</i> sp. nov., isolated from pond water near a uranium mine. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2004, 54, 1459-1463.	1.7	29
87	Identification of Fast-Growing Rhizobia Nodulating Tropical Legumes from Puerto Rico as <i>Rhizobium gallicum</i> and <i>Rhizobium tropici</i> . <i>Systematic and Applied Microbiology</i> , 2004, 27, 469-477.	2.8	46
88	<i>Microbacterium ulmi</i> sp. nov., a xylanolytic, phosphate-solubilizing bacterium isolated from sawdust of <i>Ulmus nigra</i> . <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2004, 54, 513-517.	1.7	32
89	Identification of microorganisms by PCR amplification and sequencing of a universal amplified ribosomal region present in both prokaryotes and eukaryotes. <i>Journal of Microbiological Methods</i> , 2004, 56, 413-426.	1.6	37
90	Genomic fingerprinting of <i>Frankia</i> strains by PCR-based techniques. Assessment of a primer based on the sequence of 16S rRNA gene of <i>Escherichia coli</i> . <i>Plant and Soil</i> , 2003, 254, 115-123.	3.7	6

#	ARTICLE	IF	CITATIONS
91	Description of <i>Devosia neptuniae</i> sp. nov. that Nodulates and Fixes Nitrogen in Symbiosis with <i>Neptunia natans</i> , an Aquatic Legume from India. <i>Systematic and Applied Microbiology</i> , 2003, 26, 47-53.	2.8	170
92	<i>Pseudomonas rhizosphaerae</i> sp. nov., a novel species that actively solubilizes phosphate in vitro. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2003, 53, 2067-2072.	1.7	90
93	<i>Xylanimonas cellulositytica</i> gen. nov., sp. nov., a xylanolytic bacterium isolated from a decayed tree (<i>Ulmus nigra</i>). <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2003, 53, 99-103.	1.7	88
94	Genomic fingerprinting of <i>Frankia</i> strains by PCR-based techniques. Assessment of a primer based on the sequence of 16S rRNA gene of <i>Escherichia coli</i> . , 2003, , 115-123.		0
95	A New Species of <i>Devosia</i> That Forms a Unique Nitrogen-Fixing Root-Nodule Symbiosis with the Aquatic Legume <i>Neptunia natans</i> (L.f.) Druce. <i>Applied and Environmental Microbiology</i> , 2002, 68, 5217-5222.	3.1	277
96	Title is missing!. <i>European Journal of Plant Pathology</i> , 2002, 108, 179-184.	1.7	42
97	<i>Rhizobium sullae</i> sp. nov. (formerly ' <i>Rhizobium hedysari</i> '), the root-nodule microsymbiont of <i>Hedysarum coronarium</i> L.. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2002, 52, 1267-1276.	1.7	31
98	<i>Rhizobium sullae</i> sp. nov. (formerly ' <i>Rhizobium hedysari</i> '), the root-nodule microsymbiont of <i>Hedysarum coronarium</i> L. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2002, 52, 1267-1276.	1.7	70
99	An effective, rapid and simple method for total RNA extraction from bacteria and yeast. <i>Journal of Microbiological Methods</i> , 2001, 47, 59-63.	1.6	42
100	Growth promotion of chickpea and barley by a phosphate solubilizing strain of <i>Mesorhizobium mediterraneum</i> under growth chamber conditions. <i>Soil Biology and Biochemistry</i> , 2001, 33, 103-110.	8.8	256
101	Growth promotion of common bean (<i>Phaseolus vulgaris</i> L.) by a strain of <i>Burkholderia cepacia</i> under growth chamber conditions. <i>Soil Biology and Biochemistry</i> , 2001, 33, 1927-1935.	8.8	80
102	A two primers random amplified polymorphic DNA procedure to obtain polymerase chain reaction fingerprints of bacterial species. <i>Electrophoresis</i> , 2001, 22, 1086-1089.	2.4	86
103	YeastIdent-Food/ProleFood, a new system for the identification of food yeasts based on physiological and biochemical tests. <i>Food Microbiology</i> , 2001, 18, 637-646.	4.2	15
104	Title is missing!. <i>European Journal of Plant Pathology</i> , 2001, 107, 931-938.	1.7	11
105	Cellulase isoenzyme profiles in <i>Frankia</i> strains belonging to different cross-inoculation groups. <i>Plant and Soil</i> , 2001, 229, 35-39.	3.7	15
106	Analysis of Stable Low Molecular Weight (LMW) RNA Profiles of Hydrocarbon Metabolizing Bacteria by Staircase Electrophoresis. <i>Systematic and Applied Microbiology</i> , 2001, 24, 290-293.	2.8	1
107	Stable Low Molecular Weight RNA Analyzed by Staircase Electrophoresis, a Molecular Signature for Both Prokaryotic and Eukaryotic Microorganisms. <i>Systematic and Applied Microbiology</i> , 2001, 24, 490-499.	2.8	19
108	Characterization of Rhizobial Isolates of <i>Phaseolus vulgaris</i> by Staircase Electrophoresis of Low-Molecular-Weight RNA. <i>Applied and Environmental Microbiology</i> , 2001, 67, 1008-1010.	3.1	54

#	ARTICLE	IF	CITATIONS
109	The beneficial plant growth-promoting association of <i>Rhizobium leguminosarum</i> bv. <i>trifolii</i> with rice roots. <i>Functional Plant Biology</i> , 2001, 28, 845.	2.1	116
110	<i>Mesorhizobium chacoense</i> sp. nov., a novel species that nodulates <i>Prosopis alba</i> in the Chaco Arido region (Argentina).. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2001, 51, 1011-1021.	1.7	100
111	Erosion of root epidermal cell walls by <i>Rhizobium</i> polysaccharide-degrading enzymes as related to primary host infection in the <i>Rhizobium</i> –legume symbiosis. <i>Canadian Journal of Microbiology</i> , 2001, 47, 475-487.	1.7	42
112	Erosion of root epidermal cell walls by <i>Rhizobium</i> polysaccharide-degrading enzymes as related to primary host infection in the <i>Rhizobium</i> –legume symbiosis. <i>Canadian Journal of Microbiology</i> , 2001, 47, 475-487.	1.7	38
113	Restriction Fragment Length Polymorphism Analysis of 16S rDNA and Low Molecular Weight RNA Profiling of Rhizobial Isolates from Shrubby Legumes Endemic to the Canary Islands. <i>Systematic and Applied Microbiology</i> , 2000, 23, 418-425.	2.8	36
114	Title is missing!. <i>European Journal of Plant Pathology</i> , 2000, 106, 789-793.	1.7	19
115	Staircase electrophoresis profiles of stable low-molecular-weight RNA—a new technique for yeast fingerprinting.. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2000, 50, 917-923.	1.7	17
116	Symbiotic characteristics and selection of autochthonous strains of <i>Sinorhizobium meliloti</i> populations in different soils. <i>Soil Biology and Biochemistry</i> , 1999, 31, 1039-1047.	8.8	16
117	Analysis of LMW RNA Profiles of <i>Frankia</i> Strains by Staircase Electrophoresis. <i>Systematic and Applied Microbiology</i> , 1998, 21, 539-545.	2.8	17
118	Analysis of Stable Low-Molecular-Weight RNA Profiles of Members of the Family <i>Rhizobiaceae</i> . <i>Applied and Environmental Microbiology</i> , 1998, 64, 1555-1559.	3.1	28
119	Title is missing!. <i>Plant and Soil</i> , 1997, 194, 99-114.	3.7	289
120	Enhancement of resolution of low molecular weight RNA profiles by staircase electrophoresis. <i>Electrophoresis</i> , 1997, 18, 1909-1911.	2.4	35
121	Natural endophytic association between <i>Rhizobium leguminosarum</i> bv. <i>trifolii</i> and rice roots and assessment of its potential to promote rice growth. , 1997, , 99-114.		19
122	Physiological and biochemical characterization of <i>Trichoderma harzianum</i> , a biological control agent against soilborne fungal plant pathogens. <i>Applied and Environmental Microbiology</i> , 1997, 63, 3189-3198.	3.1	126
123	Influence of the symbiotic plasmid (pSym) on cellulase production by <i>Rhizobium leguminosarum</i> bv. <i>trifolii</i> ANU843. <i>Soil Biology and Biochemistry</i> , 1996, 28, 131-133.	8.8	11
124	Cell-bound cellulase and polygalacturonase production by <i>Rhizobium</i> and <i>Bradyrhizobium</i> species. <i>Soil Biology and Biochemistry</i> , 1996, 28, 917-921.	8.8	37
125	Subnanomolar concentrations of membrane chitolipooligosaccharides from <i>Rhizobium leguminosarum</i> biovar <i>trifolii</i> are fully capable of eliciting symbiosis-related responses on white clover. <i>Plant and Soil</i> , 1996, 186, 93-98.	3.7	15
126	Defective Infection and Nodulation of Clovers by Exopolysaccharide Mutants of <i>Rhizobium leguminosarum</i> bv. <i>trifolii</i> . <i>Functional Plant Biology</i> , 1996, 23, 285.	2.1	51

#	ARTICLE	IF	CITATIONS
127	The unique root-nodule symbiosis between Rhizobium and the aquatic legume, <i>Neptunia natans</i> (L. f.) Druce. <i>Planta</i> , 1995, 196, 311-320.	3.2	82
128	Direct in situ identification of cellulose microfibrils associated with <i>Rhizobium leguminosarum</i> biovar <i>trifolii</i> attached to the root epidermis of white clover. <i>Canadian Journal of Microbiology</i> , 1995, 41, 202-207.	1.7	26
129	Attenuation of Symbiotic Effectiveness by <i>Rhizobium meliloti</i> SAF22 Related to the Presence of a Cryptic Plasmid. <i>Applied and Environmental Microbiology</i> , 1995, 61, 2033-2036.	3.1	20
130	Cell-associated pectinolytic and cellulolytic enzymes in <i>Rhizobium leguminosarum</i> biovar <i>trifolii</i> . <i>Applied and Environmental Microbiology</i> , 1992, 58, 1816-1822.	3.1	148