
Kentaro Kohagura

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/7964568/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	An association between uric acid levels and renal arteriolopathy in chronic kidney disease: a biopsy-based study. Hypertension Research, 2013, 36, 43-49.	2.7	116
2	Nonproteinuric Versus Proteinuric Phenotypes in Diabetic Kidney Disease: A Propensity Score–Matched Analysis of a Nationwide, Biopsy-Based Cohort Study. Diabetes Care, 2019, 42, 891-902.	8.6	77
3	Nationwide multicentre kidney biopsy study of Japanese patients with type 2 diabetes. Nephrology Dialysis Transplantation, 2018, 33, 138-148.	0.7	62
4	Inflammation as a Risk of Developing Chronic Kidney Disease in Rheumatoid Arthritis. PLoS ONE, 2016, 11, e0160225.	2.5	42
5	Hyperuricemia predicts future metabolic syndrome: a 4-year follow-up study of a large screened cohort in Okinawa, Japan. Hypertension Research, 2014, 37, 232-238.	2.7	41
6	Associations between serum uric acid levels and the incidence of hypertension and metabolic syndrome: a 4-year follow-up study of a large screened cohort in Okinawa, Japan. Hypertension Research, 2015, 38, 213-218.	2.7	39
7	Prevalence of anemia according to stage of chronic kidney disease in a large screening cohort of Japanese. Clinical and Experimental Nephrology, 2009, 13, 614-620.	1.6	32
8	Effects of xanthine oxidase inhibitors on renal function and blood pressure in hypertensive patients with hyperuricemia. Hypertension Research, 2016, 39, 593-597.	2.7	29
9	Chronic kidney disease, inflammation, and cardiovascular disease risk in rheumatoid arthritis. Journal of Cardiology, 2018, 71, 277-283.	1.9	29
10	Clinicopathological analysis of biopsy-proven diabetic nephropathy based on the Japanese classification of diabetic nephropathy. Clinical and Experimental Nephrology, 2018, 22, 570-582.	1.6	28
11	Plasma Aldosterone in Hypertensive Patients on Chronic Hemodialysis: Distribution, Determinants and Impact on Survival. Hypertension Research, 2006, 29, 597-604.	2.7	17
12	Modification of the impact of hypertension on proteinuria by renal arteriolar hyalinosis in nonnephrotic chronic kidney disease. Journal of Hypertension, 2016, 34, 2274-2279.	0.5	16
13	Clinicopathological features of fast eGFR decliners among patients with diabetic nephropathy. BMJ Open Diabetes Research and Care, 2020, 8, e001157.	2.8	16
14	Proteinuria and decreased body mass index as a significant risk factor in developing end-stage renal disease. Clinical and Experimental Nephrology, 2008, 12, 363-369.	1.6	15
15	Olmesartan clinical trial in Okinawan patients under OKIDS (OCTOPUS) study: design and methods. Clinical and Experimental Nephrology, 2009, 13, 145-151.	1.6	14
16	Nationwide multicenter kidney biopsy study of Japanese patients with hypertensive nephrosclerosis. Clinical and Experimental Nephrology, 2018, 22, 629-637.	1.6	14
17	Serum hemoglobin concentration and risk of renal function decline in early stages of diabetic kidney disease: a nationwide, biopsy-based cohort study. Nephrology Dialysis Transplantation, 2022, 37, 489-497.	0.7	14
18	Luseogliflozin, a sodium-glucose cotransporter 2 inhibitor, preserves renal function irrespective of acute changes in the estimated glomerular filtration rate in Japanese patients with type 2 diabetes. Hypertension Research, 2020, 43, 876-883.	2.7	13

Kentaro Kohagura

#	Article	IF	CITATIONS
19	Augmented Association Between Blood Pressure and Proteinuria in Hyperuricemic Patients With Nonnephrotic Chronic Kidney Disease. American Journal of Hypertension, 2018, 31, 480-485.	2.0	7
20	Hyponatremia and mortality among very elderly residents in a geriatric health service facility. Clinical and Experimental Nephrology, 2018, 22, 1404-1410.	1.6	7
21	Add-On Effect of Angiotensin Receptor Blockade (Candesartan) on Clinical Remission in Active IgA Nephropathy Patients Treated with Steroid Pulse Therapy and Tonsillectomy: a Randomized, Parallel-Group Comparison Trial. Kidney and Blood Pressure Research, 2018, 43, 780-792.	2.0	6
22	rHuEPO Dose Inversely Correlated with the Number of Circulating CD34+ Cells in Maintenance Hemodialysis Patients. Nephron Clinical Practice, 2008, 108, c41-c46.	2.3	5
23	Hypertriglyceridemia accompanied by increased serum complement component 3 and proteinuria in non-nephrotic chronic kidney disease. Clinical and Experimental Nephrology, 2014, 18, 453-460.	1.6	4
24	The Association between Glomerular Diameter and Secondary Focal Segmental Glomerulosclerosis in Chronic Kidney Disease. Kidney and Blood Pressure Research, 2021, 46, 433-440.	2.0	4
25	A high normal ankle-brachial index is associated with biopsy-proven severe renal small artery intimal thickening and impaired renal function in chronic kidney disease. Hypertension Research, 2020, 43, 929-937.	2.7	3
26	Age-related Changes in Renal Arterio-Arteriolosclerosis in Kidney Disease:ÂRenal Biopsy-based Study. Kidney International Reports, 2022, 7, 2101-2104.	0.8	3
27	Amplified Association Between Blood Pressure and Albuminuria in Overweight Patients With Biopsy-Proven Hypertensive Nephrosclerosis. American Journal of Hypertension, 2019, 32, 486-491.	2.0	2
28	Therapeutic drug monitoring in peritoneal dialysis: A case of nontuberculous mycobacterium catheterâ€related infection treated with amikacin. Clinical Case Reports (discontinued), 2020, 8, 995-998.	0.5	2
29	Two-year longitudinal trajectory patterns of albuminuria and subsequent rates of end-stage kidney disease and all-cause death: a nationwide cohort study of biopsy-proven diabetic kidney disease. BMJ Open Diabetes Research and Care, 2021, 9, e002241.	2.8	2
30	Incidental detection of Corynebacterium jeikeium endocarditis via regular blood examination in an afebrile hemodialysis patient. CEN Case Reports, 2020, 9, 220-224.	0.9	1
31	Understanding the Complex Interaction Between Uric Acid and Hypertension. American Journal of Hypertension, 2020, 33, 822-824.	2.0	1
32	Association of urinary angiotensinogen with renal arteriolar remodeling in chronic kidney disease. Journal of Hypertension, 2022, 40, 650-657.	0.5	1
33	Response to: The Contribution of Nutrition to the Protective Value of High Plasma Aldosterone Concentrations in Hemodialysis Patients. Hypertension Research, 2007, 30, 752.	2.7	Ο
34	Long-term efficacy of tonsillectomy as a treatment for IgA nephropathy. Journal of Japan Society of Immunology & Allergology in Otolaryngology, 2018, 36, 1-6.	0.0	0
35	1. Pathophysiology and Therapeutic Strategies of Nephrosclerosis in the Ageing Society, Japan. The Journal of the Japanese Society of Internal Medicine, 2016, 105, 811-817.	0.0	0
36	Changes in serum concentration of rilpivirine in an HIV-infected patient treated with a combination therapy of hemodialysis and peritoneal dialysis. Renal Replacement Therapy, 2020, 6, .	0.7	0