

Kentaro Kohagura

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7964568/publications.pdf>

Version: 2024-02-01

36
papers

662
citations

623734

14
h-index

580821

25
g-index

36
all docs

36
docs citations

36
times ranked

962
citing authors

#	ARTICLE	IF	CITATIONS
1	Serum hemoglobin concentration and risk of renal function decline in early stages of diabetic kidney disease: a nationwide, biopsy-based cohort study. <i>Nephrology Dialysis Transplantation</i> , 2022, 37, 489-497.	0.7	14
2	Association of urinary angiotensinogen with renal arteriolar remodeling in chronic kidney disease. <i>Journal of Hypertension</i> , 2022, 40, 650-657.	0.5	1
3	Age-related Changes in Renal Arterio-Arteriosclerosis in Kidney Disease: Renal Biopsy-based Study. <i>Kidney International Reports</i> , 2022, 7, 2101-2104.	0.8	3
4	The Association between Glomerular Diameter and Secondary Focal Segmental Glomerulosclerosis in Chronic Kidney Disease. <i>Kidney and Blood Pressure Research</i> , 2021, 46, 433-440.	2.0	4
5	Two-year longitudinal trajectory patterns of albuminuria and subsequent rates of end-stage kidney disease and all-cause death: a nationwide cohort study of biopsy-proven diabetic kidney disease. <i>BMJ Open Diabetes Research and Care</i> , 2021, 9, e002241.	2.8	2
6	A high normal ankle-brachial index is associated with biopsy-proven severe renal small artery intimal thickening and impaired renal function in chronic kidney disease. <i>Hypertension Research</i> , 2020, 43, 929-937.	2.7	3
7	Clinicopathological features of fast eGFR decliners among patients with diabetic nephropathy. <i>BMJ Open Diabetes Research and Care</i> , 2020, 8, e001157.	2.8	16
8	Luseogliflozin, a sodium-glucose cotransporter 2 inhibitor, preserves renal function irrespective of acute changes in the estimated glomerular filtration rate in Japanese patients with type 2 diabetes. <i>Hypertension Research</i> , 2020, 43, 876-883.	2.7	13
9	Incidental detection of <i>Corynebacterium jeikeium</i> endocarditis via regular blood examination in an afebrile hemodialysis patient. <i>CEN Case Reports</i> , 2020, 9, 220-224.	0.9	1
10	Therapeutic drug monitoring in peritoneal dialysis: A case of nontuberculous mycobacterium catheter-related infection treated with amikacin. <i>Clinical Case Reports (discontinued)</i> , 2020, 8, 995-998.	0.5	2
11	Understanding the Complex Interaction Between Uric Acid and Hypertension. <i>American Journal of Hypertension</i> , 2020, 33, 822-824.	2.0	1
12	Changes in serum concentration of rilpivirine in an HIV-infected patient treated with a combination therapy of hemodialysis and peritoneal dialysis. <i>Renal Replacement Therapy</i> , 2020, 6, .	0.7	0
13	Nonproteinuric Versus Proteinuric Phenotypes in Diabetic Kidney Disease: A Propensity Score-Matched Analysis of a Nationwide, Biopsy-Based Cohort Study. <i>Diabetes Care</i> , 2019, 42, 891-902.	8.6	77
14	Amplified Association Between Blood Pressure and Albuminuria in Overweight Patients With Biopsy-Proven Hypertensive Nephrosclerosis. <i>American Journal of Hypertension</i> , 2019, 32, 486-491.	2.0	2
15	Augmented Association Between Blood Pressure and Proteinuria in Hyperuricemic Patients With Nonnephrotic Chronic Kidney Disease. <i>American Journal of Hypertension</i> , 2018, 31, 480-485.	2.0	7
16	Nationwide multicentre kidney biopsy study of Japanese patients with type 2 diabetes. <i>Nephrology Dialysis Transplantation</i> , 2018, 33, 138-148.	0.7	62
17	Clinicopathological analysis of biopsy-proven diabetic nephropathy based on the Japanese classification of diabetic nephropathy. <i>Clinical and Experimental Nephrology</i> , 2018, 22, 570-582.	1.6	28
18	Chronic kidney disease, inflammation, and cardiovascular disease risk in rheumatoid arthritis. <i>Journal of Cardiology</i> , 2018, 71, 277-283.	1.9	29

#	ARTICLE	IF	CITATIONS
19	Nationwide multicenter kidney biopsy study of Japanese patients with hypertensive nephrosclerosis. <i>Clinical and Experimental Nephrology</i> , 2018, 22, 629-637.	1.6	14
20	Long-term efficacy of tonsillectomy as a treatment for IgA nephropathy. <i>Journal of Japan Society of Immunology & Allergology in Otolaryngology</i> , 2018, 36, 1-6.	0.0	0
21	Add-On Effect of Angiotensin Receptor Blockade (Candesartan) on Clinical Remission in Active IgA Nephropathy Patients Treated with Steroid Pulse Therapy and Tonsillectomy: a Randomized, Parallel-Group Comparison Trial. <i>Kidney and Blood Pressure Research</i> , 2018, 43, 780-792.	2.0	6
22	Hyponatremia and mortality among very elderly residents in a geriatric health service facility. <i>Clinical and Experimental Nephrology</i> , 2018, 22, 1404-1410.	1.6	7
23	Modification of the impact of hypertension on proteinuria by renal arteriolar hyalinosis in nonnephrotic chronic kidney disease. <i>Journal of Hypertension</i> , 2016, 34, 2274-2279.	0.5	16
24	Effects of xanthine oxidase inhibitors on renal function and blood pressure in hypertensive patients with hyperuricemia. <i>Hypertension Research</i> , 2016, 39, 593-597.	2.7	29
25	Inflammation as a Risk of Developing Chronic Kidney Disease in Rheumatoid Arthritis. <i>PLoS ONE</i> , 2016, 11, e0160225.	2.5	42
26	1. Pathophysiology and Therapeutic Strategies of Nephrosclerosis in the Ageing Society, Japan. <i>The Journal of the Japanese Society of Internal Medicine</i> , 2016, 105, 811-817.	0.0	0
27	Associations between serum uric acid levels and the incidence of hypertension and metabolic syndrome: a 4-year follow-up study of a large screened cohort in Okinawa, Japan. <i>Hypertension Research</i> , 2015, 38, 213-218.	2.7	39
28	Hyperuricemia predicts future metabolic syndrome: a 4-year follow-up study of a large screened cohort in Okinawa, Japan. <i>Hypertension Research</i> , 2014, 37, 232-238.	2.7	41
29	Hypertriglyceridemia accompanied by increased serum complement component 3 and proteinuria in non-nephrotic chronic kidney disease. <i>Clinical and Experimental Nephrology</i> , 2014, 18, 453-460.	1.6	4
30	An association between uric acid levels and renal arteriopathy in chronic kidney disease: a biopsy-based study. <i>Hypertension Research</i> , 2013, 36, 43-49.	2.7	116
31	Olmesartan clinical trial in Okinawan patients under OKIDS (OCTOPUS) study: design and methods. <i>Clinical and Experimental Nephrology</i> , 2009, 13, 145-151.	1.6	14
32	Prevalence of anemia according to stage of chronic kidney disease in a large screening cohort of Japanese. <i>Clinical and Experimental Nephrology</i> , 2009, 13, 614-620.	1.6	32
33	Proteinuria and decreased body mass index as a significant risk factor in developing end-stage renal disease. <i>Clinical and Experimental Nephrology</i> , 2008, 12, 363-369.	1.6	15
34	rHuEPO Dose Inversely Correlated with the Number of Circulating CD34+ Cells in Maintenance Hemodialysis Patients. <i>Nephron Clinical Practice</i> , 2008, 108, c41-c46.	2.3	5
35	Response to: The Contribution of Nutrition to the Protective Value of High Plasma Aldosterone Concentrations in Hemodialysis Patients. <i>Hypertension Research</i> , 2007, 30, 752.	2.7	0
36	Plasma Aldosterone in Hypertensive Patients on Chronic Hemodialysis: Distribution, Determinants and Impact on Survival. <i>Hypertension Research</i> , 2006, 29, 597-604.	2.7	17