

Sharee N Light

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7976434/publications.pdf>

Version: 2024-02-01

14
papers

899
citations

1040056

9
h-index

1125743

13
g-index

14
all docs

14
docs citations

14
times ranked

1430
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | The Combined Use of Neuropsychiatric and Neuropsychological Assessment Tools to Make a Differential Dementia Diagnosis in the Presence of "Long-Haul" COVID-19. <i>Case Reports in Neurology</i> , 2022, 14, 130-148. | 0.7 | 3 |
| 2 | Empathy for joy recruits a broader prefrontal network than empathy for sadness and is predicted by executive functioning.. <i>Neuropsychology</i> , 2021, 35, 90-102. | 1.3 | 13 |
| 3 | Measuring change in anhedonia using the "Happy Faces" task pre- to post-repetitive transcranial magnetic stimulation (rTMS) treatment to left dorsolateral prefrontal cortex in Major Depressive Disorder (MDD): relation to empathic happiness. <i>Translational Psychiatry</i> , 2019, 9, 217. | 4.8 | 15 |
| 4 | The Measurement of Positive Valence Forms of Empathy and Their Relation to Anhedonia and Other Depressive Symptomatology. <i>Frontiers in Psychology</i> , 2019, 10, 815. | 2.1 | 7 |
| 5 | The Heterogeneity of Empathy: Possible Treatment for Anhedonia?. <i>Frontiers in Psychiatry</i> , 2019, 10, 185. | 2.6 | 0 |
| 6 | Fronto-striatal activity predicts anhedonia and positive empathy subtypes. <i>Brain Imaging and Behavior</i> , 2019, 13, 1554-1565. | 2.1 | 6 |
| 7 | "Top-Down" Mu-Opioid System Function in Humans: Mu-Opioid Receptors in Ventrolateral Prefrontal Cortex Mediate the Relationship Between Hedonic Tone and Executive Function in Major Depressive Disorder. <i>Journal of Neuropsychiatry and Clinical Neurosciences</i> , 2017, 29, 357-364. | 1.8 | 18 |
| 8 | Patterns of frontoparietal activation as a marker for unsuccessful visuospatial processing in healthy aging. <i>Brain Imaging and Behavior</i> , 2016, 10, 686-696. | 2.1 | 15 |
| 9 | Electromyographically assessed empathic concern and empathic happiness predict increased prosocial behavior in adults. <i>Biological Psychology</i> , 2015, 104, 116-129. | 2.2 | 52 |
| 10 | Relationships Between Changes in Sustained Fronto-Striatal Connectivity and Positive Affect in Major Depression Resulting From Antidepressant Treatment. <i>American Journal of Psychiatry</i> , 2013, 170, 197-206. | 7.2 | 140 |
| 11 | Reduced Right Ventrolateral Prefrontal Cortex Activity While Inhibiting Positive Affect Is Associated with Improvement in Hedonic Capacity After 8 Weeks of Antidepressant Treatment in Major Depressive Disorder. <i>Biological Psychiatry</i> , 2011, 70, 962-968. | 1.3 | 82 |
| 12 | Reduced capacity to sustain positive emotion in major depression reflects diminished maintenance of fronto-striatal brain activation. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2009, 106, 22445-22450. | 7.1 | 383 |
| 13 | Empathy Is Associated With Dynamic Change in Prefrontal Brain Electrical Activity During Positive Emotion in Children. <i>Child Development</i> , 2009, 80, 1210-1231. | 3.0 | 150 |
| 14 | Dynamic variation in pleasure in children predicts nonlinear change in lateral frontal brain electrical activity.. <i>Developmental Psychology</i> , 2009, 45, 525-533. | 1.6 | 15 |