

Joseph P Dillard

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8000382/publications.pdf>

Version: 2024-02-01

65
papers

2,622
citations

186265

28
h-index

206112

48
g-index

68
all docs

68
docs citations

68
times ranked

2175
citing authors

#	ARTICLE	IF	CITATIONS
1	The AmiC/NlpD Pathway Dominates Peptidoglycan Breakdown in <i>Neisseria meningitidis</i> and Affects Cell Separation, NOD1 Agonist Production, and Infection. <i>Infection and Immunity</i> , 2022, 90, IA10048521.	2.2	4
2	Transcriptional and Translational Responsiveness of the <i>Neisseria gonorrhoeae</i> Type IV Secretion System to Conditions of Host Infections. <i>Infection and Immunity</i> , 2021, 89, e0051921.	2.2	5
3	Expression, Localization, and Protein Interactions of the Partitioning Proteins in the Gonococcal Type IV Secretion System. <i>Frontiers in Microbiology</i> , 2021, 12, 784483.	3.5	4
4	Hold It Right There! Gonococci Preserve Epithelium Integrity during Intimate Adherence. <i>Cell Host and Microbe</i> , 2020, 27, 685-686.	11.0	1
5	Protein interactions within and between two <i>Neisseria gonorrhoeae</i> type IV secretion systems. <i>Molecular Microbiology</i> , 2020, 114, 823-838.	2.5	11
6	The NtrYX Two-Component System Regulates the Bacterial Cell Envelope. <i>MBio</i> , 2020, 11, .	4.1	22
7	Antigenic Variation in <i>Neisseria gonorrhoeae</i> Occurs Independently of RecQ-Mediated Unwinding of the <i>pilE</i> G Quadruplex. <i>Journal of Bacteriology</i> , 2020, 202, .	2.2	9
8	Defective lytic transglycosylase disrupts cell morphogenesis by hindering cell wall de-O-acetylation in <i>Neisseria meningitidis</i> . <i>ELife</i> , 2020, 9, .	6.0	7
9	The Pathogenic <i>Neisseria</i> Use a Streamlined Set of Peptidoglycan Degradation Proteins for Peptidoglycan Remodeling, Recycling, and Toxic Fragment Release. <i>Frontiers in Microbiology</i> , 2019, 10, 73.	3.5	14
10	Transformation in <i>Neisseria gonorrhoeae</i> . <i>Methods in Molecular Biology</i> , 2019, 1997, 143-162.	0.9	8
11	Peptidoglycan Composition in <i>Neisseria</i> . <i>Methods in Molecular Biology</i> , 2019, 1997, 111-120.	0.9	0
12	Mucus Is a Key Factor in <i>Neisseria meningitidis</i> Commensalism. <i>MSphere</i> , 2019, 4, .	2.9	1
13	<i>Neisseria gonorrhoeae</i> PBP3 and PBP4 Facilitate NOD1 Agonist Peptidoglycan Fragment Release and Survival in Stationary Phase. <i>Infection and Immunity</i> , 2019, 87, .	2.2	6
14	The low-molecular-mass, penicillin-binding proteins DacB and DacC combine to modify peptidoglycan cross-linking and allow stable Type IV pilus expression in <i>Neisseria gonorrhoeae</i> . <i>Molecular Microbiology</i> , 2018, 109, 135-149.	2.5	11
15	Pathogenesis of <i>Neisseria gonorrhoeae</i> and the Host Defense in Ascending Infections of Human Fallopian Tube. <i>Frontiers in Immunology</i> , 2018, 9, 2710.	4.8	61
16	Selective Inhibition of <i>Neisseria gonorrhoeae</i> by a Dithiazoline in Mixed Infections with <i>Lactobacillus gasseri</i> . <i>Antimicrobial Agents and Chemotherapy</i> , 2018, 62, .	3.2	5
17	Antibiotic Targets in Gonococcal Cell Wall Metabolism. <i>Antibiotics</i> , 2018, 7, 64.	3.7	3
18	<i>Neisseria gonorrhoeae</i> Lytic Transglycosylases LtgA and LtgD Reduce Host Innate Immune Signaling through TLR2 and NOD2. <i>ACS Infectious Diseases</i> , 2017, 3, 624-633.	3.8	27

#	ARTICLE	IF	CITATIONS
19	A Single Dual-Function Enzyme Controls the Production of Inflammatory NOD Agonist Peptidoglycan Fragments by <i>Neisseria gonorrhoeae</i> . MBio, 2017, 8, .	4.1	28
20	Attention Seeker: Production, Modification, and Release of Inflammatory Peptidoglycan Fragments in <i>Neisseria</i> Species. Journal of Bacteriology, 2017, 199, .	2.2	21
21	Two lytic transglycosylases in <i>Neisseria gonorrhoeae</i> impart resistance to killing by lysozyme and human neutrophils. Cellular Microbiology, 2017, 19, e12662.	2.1	52
22	Secretion of Chromosomal DNA by the <i>Neisseria gonorrhoeae</i> Type IV Secretion System. Current Topics in Microbiology and Immunology, 2017, 413, 323-345.	1.1	20
23	Digestion of Peptidoglycan and Analysis of Soluble Fragments. Bio-protocol, 2017, 7, .	0.4	30
24	nagZ Triggers Gonococcal Biofilm Disassembly. Scientific Reports, 2016, 6, 22372.	3.3	27
25	Lytic transglycosylases LtgA and LtgD perform distinct roles in remodeling, recycling and releasing peptidoglycan in <i>Neisseria gonorrhoeae</i> . Molecular Microbiology, 2016, 102, 865-881.	2.5	38
26	<i>Neisseria gonorrhoeae</i> Crippled Its Peptidoglycan Fragment Permease To Facilitate Toxic Peptidoglycan Monomer Release. Journal of Bacteriology, 2016, 198, 3029-3040.	2.2	24
27	Genomic analyses of <i>Neisseria gonorrhoeae</i> reveal an association of the gonococcal genetic island with antimicrobial resistance. Journal of Infection, 2016, 73, 578-587.	3.3	54
28	Type I Interferon Induction by <i>Neisseria gonorrhoeae</i> : Dual Requirement of Cyclic GMP-AMP Synthase and Toll-like Receptor 4. Cell Reports, 2016, 15, 2438-2448.	6.4	66
29	Amidase Activity of AmiC Controls Cell Separation and Stem Peptide Release and Is Enhanced by NlpD in <i>Neisseria gonorrhoeae</i> . Journal of Biological Chemistry, 2016, 291, 10916-10933.	3.4	26
30	Analysis of Peptidoglycan Fragment Release. Methods in Molecular Biology, 2016, 1440, 185-200.	0.9	9
31	The Gonococcal NlpD Protein Facilitates Cell Separation by Activating Peptidoglycan Cleavage by AmiC. Journal of Bacteriology, 2016, 198, 615-622.	2.2	25
32	Targeted mutagenesis of intergenic regions in the <i>Neisseria gonorrhoeae</i> gonococcal genetic island reveals multiple regulatory mechanisms controlling type IV secretion. Molecular Microbiology, 2015, 97, 1168-1185.	2.5	9
33	Structural and Functional Features of a Developmentally Regulated Lipopolysaccharide-Binding Protein. MBio, 2015, 6, e01193-15.	4.1	16
34	TraK and TraB Are Conserved Outer Membrane Proteins of the <i>Neisseria gonorrhoeae</i> Type IV Secretion System and Are Expressed at Low Levels in Wild-Type Cells. Journal of Bacteriology, 2014, 196, 2954-2968.	2.2	17
35	Functional Analysis of the Gonococcal Genetic Island of <i>Neisseria gonorrhoeae</i> . PLoS ONE, 2014, 9, e109613.	2.5	26
36	Peptidoglycan Fragment Release from <i>Neisseria meningitidis</i> . Infection and Immunity, 2013, 81, 3490-3498.	2.2	43

#	ARTICLE	IF	CITATIONS
37	Mating Pair Formation Homologue TraG Is a Variable Membrane Protein Essential for Contact-Independent Type IV Secretion of Chromosomal DNA by <i>Neisseria gonorrhoeae</i> . <i>Journal of Bacteriology</i> , 2013, 195, 1666-1679.	2.2	23
38	Prevalence and Detailed Mapping of the Gonococcal Genetic Island in <i>Neisseria meningitidis</i> . <i>Journal of Bacteriology</i> , 2012, 194, 2275-2285.	2.2	31
39	<i>Neisseria gonorrhoeae</i> Virulence Factor NG1686 Is a Bifunctional M23B Family Metallopeptidase That Influences Resistance to Hydrogen Peroxide and Colony Morphology. <i>Journal of Biological Chemistry</i> , 2012, 287, 11222-11233.	3.4	27
40	New Complementation Constructs for Inducible and Constitutive Gene Expression in <i>Neisseria gonorrhoeae</i> and <i>Neisseria meningitidis</i> . <i>Applied and Environmental Microbiology</i> , 2012, 78, 3068-3078.	3.1	44
41	The Lytic Transglycosylases of <i>Neisseria gonorrhoeae</i> . <i>Microbial Drug Resistance</i> , 2012, 18, 271-279.	2.0	46
42	Characterization of the Single Stranded DNA Binding Protein SsbB Encoded in the Gonococcal Genetic Island. <i>PLoS ONE</i> , 2012, 7, e35285.	2.5	16
43	Genetic Manipulation of <i>Neisseria gonorrhoeae</i> . <i>Current Protocols in Microbiology</i> , 2011, 23, Unit4A.2.	6.5	82
44	The Gonococcal Genetic Island and Type IV Secretion in the Pathogenic <i>Neisseria</i> . <i>Frontiers in Microbiology</i> , 2011, 2, 61.	3.5	54
45	XerCD-Mediated Site-Specific Recombination Leads to Loss of the 57-Kilobase Gonococcal Genetic Island. <i>Journal of Bacteriology</i> , 2011, 193, 377-388.	2.2	41
46	Increased Expression of the Type IV Secretion System in Piliated <i>Neisseria gonorrhoeae</i> Variants. <i>Journal of Bacteriology</i> , 2010, 192, 1912-1920.	2.2	31
47	Type IV Secretion Machinery Promotes Ton-Independent Intracellular Survival of <i>Neisseria gonorrhoeae</i> within Cervical Epithelial Cells. <i>Infection and Immunity</i> , 2010, 78, 2429-2437.	2.2	22
48	RecQ DNA helicase HRDC domains are critical determinants in <i>Neisseria gonorrhoeae</i> pilin antigenic variation and DNA repair. <i>Molecular Microbiology</i> , 2009, 71, 158-171.	2.5	11
49	<i>Neisseria gonorrhoeae</i> Uses Two Lytic Transglycosylases To Produce Cytotoxic Peptidoglycan Monomers. <i>Journal of Bacteriology</i> , 2008, 190, 5989-5994.	2.2	49
50	Mutations in <i>ampG</i> or <i>ampD</i> Affect Peptidoglycan Fragment Release from <i>Neisseria gonorrhoeae</i> . <i>Journal of Bacteriology</i> , 2008, 190, 3799-3807.	2.2	43
51	AtIA Functions as a Peptidoglycan Lytic Transglycosylase in the <i>Neisseria gonorrhoeae</i> Type IV Secretion System. <i>Journal of Bacteriology</i> , 2007, 189, 5421-5428.	2.2	48
52	A novel relaxase homologue is involved in chromosomal DNA processing for type IV secretion in <i>Neisseria gonorrhoeae</i> . <i>Molecular Microbiology</i> , 2007, 66, 930-947.	2.5	47
53	Natural transformation of <i>Neisseria gonorrhoeae</i> : from DNA donation to homologous recombination. <i>Molecular Microbiology</i> , 2006, 59, 376-385.	2.5	187
54	AmiC Functions as an <i>N</i> -Acetylmuramyl- Alanine Amidase Necessary for Cell Separation and Can Promote Autolysis in <i>Neisseria gonorrhoeae</i> . <i>Journal of Bacteriology</i> , 2006, 188, 7211-7221.	2.2	69

#	ARTICLE	IF	CITATIONS
55	Genetic Manipulation of <i>Neisseria gonorrhoeae</i> . , 2006, Chapter 4, Unit 4A.2.		49
56	<i>Neisseria gonorrhoeae</i> secretes chromosomal DNA via a novel type IV secretion system. <i>Molecular Microbiology</i> , 2005, 55, 1704-1721.	2.5	254
57	Mutations Affecting Peptidoglycan Acetylation in <i>Neisseria gonorrhoeae</i> and <i>Neisseria meningitidis</i> . <i>Infection and Immunity</i> , 2005, 73, 5697-5705.	2.2	59
58	Mutation of a Single Lytic Transglycosylase Causes Aberrant Septation and Inhibits Cell Separation of <i>Neisseria gonorrhoeae</i> . <i>Journal of Bacteriology</i> , 2004, 186, 7811-7814.	2.2	44
59	A Lytic Transglycosylase of <i>Neisseria gonorrhoeae</i> Is Involved in Peptidoglycan-Derived Cytotoxin Production. <i>Infection and Immunity</i> , 2002, 70, 2752-2757.	2.2	65
60	A variable genetic island specific for <i>Neisseria gonorrhoeae</i> is involved in providing DNA for natural transformation and is found more often in disseminated infection isolates. <i>Molecular Microbiology</i> , 2001, 41, 263-277.	2.5	173
61	Insertion-Duplication Mutagenesis of <i>Neisseria</i> : Use in Characterization of DNA Transfer Genes in the Gonococcal Genetic Island. <i>Journal of Bacteriology</i> , 2001, 183, 4718-4726.	2.2	76
62	The pathogenic neisseriae contain an inactive <i>rpoN</i> gene and do not utilize the <i>pilE</i> promoter. <i>Gene</i> , 1998, 208, 95-102.	2.2	38
63	Pneumococcal Diversity: Considerations for New Vaccine Strategies with Emphasis on Pneumococcal Surface Protein A (PspA). <i>Clinical Microbiology Reviews</i> , 1998, 11, 645-657.	13.6	139
64	A peptidoglycan hydrolase similar to bacteriophage endolysins acts as an autolysin in <i>Neisseria gonorrhoeae</i> . <i>Molecular Microbiology</i> , 1997, 25, 893-901.	2.5	39
65	Genetic and molecular characterization of capsular polysaccharide biosynthesis in <i>Streptococcus pneumoniae</i> type 3. <i>Molecular Microbiology</i> , 1994, 12, 959-972.	2.5	79