

Andrzej Witkowski

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8001527/publications.pdf>

Version: 2024-02-01

127
papers

1,973
citations

304743
22
h-index

345221
36
g-index

129
all docs

129
docs citations

129
times ranked

2394
citing authors

#	ARTICLE	IF	CITATIONS
1	A fossil diatom-based reconstruction of sea-level changes for the Late Pleistocene and Holocene period in the NW South China Sea. <i>Oceanologia</i> , 2023, 65, 211-229.	2.2	2
2	Assessment of marine benthic diatom communities: insights from a combined morphologicalâ€“metabarcoding approach in Mediterranean shallow coastal waters. <i>Marine Pollution Bulletin</i> , 2022, 174, 113183.	5.0	13
3	Life History of the Diatom <i>Schizostauron trachyderma</i> : Cell Size and Lipid Accumulation. <i>Frontiers in Marine Science</i> , 2022, 8, .	2.5	0
4	What Was Old Is New Again: The Pennate Diatom <i>Haslea ostrearia</i> (Gaiillon) Simonsen in the Multi-Omic Age. <i>Marine Drugs</i> , 2022, 20, 234.	4.6	5
5	Lipid Constituents of Diatoms (Halamphora) as Components for Production of Lipid Nanoparticles. <i>Pharmaceutics</i> , 2022, 14, 1171.	4.5	3
6	Ripe for reassessment: A synthesis of available molecular data for the speciose diatom family Bacillariaceae. <i>Molecular Phylogenetics and Evolution</i> , 2021, 158, 106985.	2.7	34
7	Morphological and molecular identification reveals that waters from an isolated oasis in Tamanrasset (extreme South of Algerian Sahara) are colonized by opportunistic and pollution-tolerant diatom species. <i>Ecological Indicators</i> , 2021, 121, 107104.	6.3	9
8	A diatom-based Holocene record of sedimentary and oceanographic environmental changes within the Beibu Gulf, NW South China Sea. <i>Marine Geology</i> , 2021, 432, 106395.	2.1	3
9	<i>Cocconeis vaiamanuensis</i> sp. nov. (Bacillariophyceae) from Raivavae (South Pacific) and allied taxa: ultrastructural specificities and remarks about the polyphyletic genus <i>Cocconeis</i> Ehrenberg. <i>Marine Biodiversity</i> , 2021, 51, 1.	1.0	2
10	Three new <i>Luticola</i> D.G.Mann (Bacillariophyta) species from Rapa Nui (Easter Island) found in terrestrial diatom assemblages dominated by widely distributed taxa. <i>PeerJ</i> , 2021, 9, e11142.	2.0	6
11	<i>Haslea silbo</i> , A Novel Cosmopolitan Species of Blue Diatoms. <i>Biology</i> , 2021, 10, 328.	2.8	12
12	Diatom Genus <i>Hyalosira</i> (Rhabdonematales emend.) and Resolution of its Polyphyly in Grammatophoraceae and Rhabdonemataceae with a New Genus, <i>Placosira</i> , and Five New <i>Hyalosira</i> Species. <i>Protist</i> , 2021, 172, 125816.	1.5	10
13	Extreme Enlargement of the Inverted Repeat Region in the Plastid Genomes of Diatoms from the Genus <i>Climaconeis</i> . <i>International Journal of Molecular Sciences</i> , 2021, 22, 7155.	4.1	8
14	Reducing Efficiency of Fucoxanthin in Diatom Mediated Biofabrication of Gold Nanoparticles. <i>Materials</i> , 2021, 14, 4094.	2.9	14
15	Multigene phylogenetic data place monoraphid diatoms <i>Schizostauron</i> and <i>Astartiella</i> along with other fistulaâ€“bearing genera in the Stauroneidaceae 1. <i>Journal of Phycology</i> , 2021, 57, 1472-1491.	2.3	5
16	Insight into diatom frustule structures using various imaging techniques. <i>Scientific Reports</i> , 2021, 11, 14555.	3.3	12
17	Indonesian coral reef habitats reveal exceptionally high species richness and biodiversity of diatom assemblages. <i>Estuarine, Coastal and Shelf Science</i> , 2021, 261, 107551.	2.1	21
18	Novel Diatoms (Bacillariophyta) from tropical and temperate marine littoral habitats with the description of <i>Catenulopsis</i> gen. nov., and two <i>Catenula</i> species. <i>Diatom Research</i> , 2021, 36, 265-280.	1.2	4

#	ARTICLE	IF	CITATIONS
19	<i>Nitzschia anatoliensis</i> sp. nov., a cryptic diatom species from the highly alkaline Van Lake (Turkey). PeerJ, 2021, 9, e12220.	2.0	4
20	Mitochondrial and Plastid Genomes of the Monoraphid Diatom <i>Schizostauron trachyderma</i> . International Journal of Molecular Sciences, 2021, 22, 11139.	4.1	5
21	Diatom Mediated Production of Fluorescent Flower Shaped Silver-Silica Nanohybrid. Materials, 2021, 14, 7284.	2.9	6
22	Marine diatom assemblages of the Nosy Be Island coasts, NW Madagascar: species composition and biodiversity using molecular and morphological taxonomy. Systematics and Biodiversity, 2020, 18, 161-180.	1.2	8
23	A hybrid biomaterial of biosilica and C-phycocyanin for enhanced photodynamic effect towards tumor cells. Biochemical and Biophysical Research Communications, 2020, 533, 573-579.	2.1	11
24	Marine diatom response to oceanographic and climatic changes in the NW South China Sea since the penultimate glacial interval. Journal of Asian Earth Sciences, 2020, 204, 104553.	2.3	4
25	Austral winter marine epilithic diatoms: Community composition and distribution on intertidal rocky substrate around the coast of South Africa. Estuarine, Coastal and Shelf Science, 2020, 242, 106837.	2.1	5
26	â€œOutsourcingâ€œDiatoms in Fabrication of Metal-Doped 3D Biosilica. Materials, 2020, 13, 2576.	2.9	13
27	Exploring Diversity, Taxonomy and Phylogeny of Diatoms (Bacillariophyta) from Marine Habitats. Novel Taxa with Internal Costae. Protist, 2020, 171, 125713.	1.5	11
28	Morphology, phylogeny, and molecular dating in Plagiogrammaceae family focused on Plagiogramma-Dimeregramma complex (Urneidophycidae, Bacillariophyceae). Molecular Phylogenetics and Evolution, 2020, 148, 106808.	2.7	2
29	The Taxonomy and Diversity of <i>Proschkinia</i> (Bacillariophyta), A Common But Enigmatic Genus from Marine Coasts. Journal of Phycology, 2020, 56, 953-978.	2.3	5
30	<i>Navicula dermochelycola</i> sp. nov., presumably an exclusively epizoic diatom on sea turtles <i>Dermochelys coriacea</i> and <i>Lepidochelys olivacea</i> from French Guiana. Oceanological and Hydrobiological Studies, 2020, 49, 132-139.	0.7	6
31	Majewskaea gen. nov. (Bacillariophyta), a new marine benthic diatom genus from the Adriatic Sea. Fottea, 2020, 20, 112-120.	0.9	5
32	Biodiversity of carapace epibiont diatoms in loggerhead sea turtles (<i>Caretta caretta</i> Linnaeus) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 20		
33	Morphology and molecular phylogeny of Gomphonemopsis sieminskae sp. nov. isolated from brackish waters of the East China Sea coast. Plant and Fungal Systematics, 2019, 64, 17-24.	0.5	1
34	Discovery of a kleptoplastic â€œdinotomâ€™ dinoflagellate and the unique nuclear dynamics of converting kleptoplastids to permanent plastids. Scientific Reports, 2019, 9, 10474.	3.3	25
35	Complete chloroplast genome of the tiny marine diatom <i>Nanofrustulum shiloii</i> (Bacillariophyta) from the Adriatic Sea. Mitochondrial DNA Part B: Resources, 2019, 4, 3374-3376.	0.4	5
36	Cocconeis nosybetiana sp. nov. from Nosy Be Island (Madagascar) and allied taxa. Nova Hedwigia, 2019, 108, 321-338.	0.4	2

#	ARTICLE	IF	CITATIONS
37	< i>Cocconeis carinata</i> sp. nov. (Bacillariophyceae) and re-examination of < i>Cocconeis orbicularis</i> Frenguelli & H.A.Orlando and < i>Cocconeis reticulata</i> var. < i>deceptionis</i> Frenguelli & H.A.Orlando. Diatom Research, 2019, 34, 149-163.	1.2	2
38	<p>Diatom phenotypic plasticity: Olifiantiella gorandiana epizoic on â€˜G5-Manahereâ€™ (Society Archipelago, South Pacific), a case study</p>. Phytotaxa, 2019, 415, 89-104.	0.3	5
39	The biogeography and ecology of common diatom species in the northern North Atlantic, and their implications for paleoceanographic reconstructions. Marine Micropaleontology, 2019, 148, 1-28.	1.2	23
40	Epiphytic diatom assemblages on invasive < i>Caulerpa taxifolia</i> and autochthonous < i>Halimeda tuna</i> and < i>Padina</i> sp. seaweeds in the Adriatic Sea â€“ summer/autumn aspect. Oceanological and Hydrobiological Studies, 2019, 48, 209-226.	0.7	4
41	Toward a multigene phylogeny of the Cymatosiraceae (Bacillariophyta, Mediophyceae) II: morphological and molecular insights into the taxonomy of the forgotten species < i>Campylosira africana and < i>Extubocellulus, with a description of two new taxa. Journal of Phycology, 2019, 55, 425-441.	2.3	8
42	Complete mitochondrial genome of a rare diatom (Bacillariophyta) < i>Proschkinia</i> and its phylogenetic and taxonomic implications. Mitochondrial DNA Part B: Resources, 2019, 4, 25-26.	0.4	11
43	Taxonomy and diversity of a little-known diatom genus < i>Simonsenia (Bacillariaceae) in the marine littoral: novel taxa from the Yellow Sea and the Gulf of Mexico. Plant Ecology and Evolution, 2019, 152, 248-261.	0.7	6
44	Nitzschia omanensis sp. nov., a new diatom species from the marine coast of Oman, characterized by valve morphology and molecular data. Fottea, 2019, 19, 175-184.	0.9	3
45	Isolation and identification of indigenous marine diatoms (Bacillariophyta) for biomass production in open raceway ponds. Aquaculture Research, 2018, 49, 928-938.	1.8	8
46	Multiproxy analysis of tsunami depositsâ€”The TirÃ³a example, central Chile. , 2018, 14, 1067-1086.		11
47	Achnanthales from historical Grunow collection in Porto Subzanski, Croatia. Botanica Marina, 2018, 61, 573-593.	1.2	1
48	Cocconeis kurakakea, a new diatom species from Nukutavake (Tuamotuâ€œArchipelago, South Pacific): description and comparison with < i>C. diruptoides and < i>C. pseudodiruptoides. Phytotaxa, 2018, 349, 115.	0.3	1
49	The complete mitochondrial DNA of the tropical oyster < i>Crassostrea belcheri</i> from the Cátòn GiÃ²â€™ mangrove in Vietnam. Mitochondrial DNA Part B: Resources, 2018, 3, 462-463.	0.4	4
50	The morphology and molecular phylogenetics of some marine diatom taxa within the Fragilariaeae, including twenty undescribed species and their relationship to < i>Nanofrustulum, Opephora and Pseudostaurosira. Phytotaxa, 2018, 355, 1.	0.3	35
51	Planothidium juandenovense sp. nov. (Bacillariophyta) from Juan de Nova (Scattered Islands,) Tj ETQq1 1 0.784314 rgBT /Overlock 10 T... delicatulum complex. Fottea, 2018, 18, 106-119.	0.9	9
52	A new sediment dwelling and epizoic species of < i>Olifiantiella (Bacillariophyceae), with an account on the genus ultrastructure based on Focused Ion Beam nanocuts. Fottea, 2018, 18, 212-226.	0.9	14
53	New epizoic diatom (Bacillariophyta) species from sea turtles in the Eastern Caribbean and South Pacific. Diatom Research, 2017, 32, 109-125.	1.2	18
54	Towards a multigene phylogeny of the Cymatosiraceae (Bacillariophyta, Mediophyceae) I: novel taxa within the subfamily cymatosiroideae based on molecular and morphological data. Journal of Phycology, 2017, 53, 342-360.	2.3	14

#	ARTICLE	IF	CITATIONS
55	Cocconeis subantarctica sp. nov. from Kerguelen Archipelago (Austral Ocean) and comparison with Cocconeis stauroneiformis (W.Smith) Okuno. <i>Oceanological and Hydrobiological Studies</i> , 2017, 46, 350-362.	0.7	1
56	Visualization of the internal structure of <i>Didymosphenia geminata</i> frustules using nano X-ray tomography. <i>Scientific Reports</i> , 2017, 7, 9086.	3.3	21
57	Sexual reproduction in <i>< i>Schizostauron</i></i> (Bacillariophyta) and a preliminary phylogeny of the genus. <i>Phycologia</i> , 2017, 56, 77-93.	1.4	19
58	Molecular and Morphological Investigations of the Stauros-bearing, Raphid Pennate Diatoms (Bacillariophyceae): <i>Craspedostauros</i> E.J. Cox, and <i>Staurotropis</i> T.B.B. Paddock, and their Relationship to the Rest of the Mastogloiales. <i>Protist</i> , 2017, 168, 48-70.	1.5	30
59	Novel diatom species (Bacillariophyta) from the freshwater discharge site of Laguna Diablas (Island) Tj ETQq1 1 0.784314 rgBT /Overloc	0.5	16
60	Two new <i>Tursiocola</i> species (Bacillariophyta) epizoic on green turtles (<i>Chelonia mydas</i>) in French Guiana and Eastern Caribbean. <i>Fottea</i> , 2017, 17, 150-163.	0.9	15
61	Postglacial Evolution of the Odra River Mouth, Poland-Germany. <i>Coastal Research Library</i> , 2017, , 193-217.	0.4	5
62	Late Glacial to Holocene Environmental Changes (with Particular Reference to Salinity) in the Southern Baltic Reconstructed from Shallow Water Lagoon Sediments. <i>Coastal Research Library</i> , 2017, , 175-192.	0.4	2
63	Multiphase Biominerization: Enigmatic Invasive Siliceous Diatoms Produce Crystalline Calcite. <i>Advanced Functional Materials</i> , 2016, 26, 2503-2510.	14.9	37
64	Multigene Assessment of Biodiversity of Diatom(Bacillariophyceae) Assemblages from the Littoral Zone of the Bohai and Yellow Seas in Yantai Region of Northeast China with some Remarks on Ubiquitous Taxa. <i>Journal of Coastal Research</i> , 2016, 74, 166-195.	0.3	32
65	Ultrastructural and molecular characterization of diversity among small araphid diatoms all lacking rimoportulae. I. Five new genera, eight new species. <i>Journal of Phycology</i> , 2016, 52, 1018-1036.	2.3	15
66	Significance of the <i>< i>Paralia sulcata</i></i> fossil record in palaeoenvironmental reconstructions of the SE Asia marginal seas over the Last Glacial Cycle. <i>Geological Society Special Publication</i> , 2016, 429, 211-221.	1.3	6
67	Surface and sub-surface multi-proxy reconstruction of middle to late Holocene palaeoceanographic changes in Disko Bugt, West Greenland. <i>Quaternary Science Reviews</i> , 2016, 132, 146-160.	3.0	48
68	<i>Simonsenia aveniformis</i> sp. nov. (Bacillariophyceae), molecular phylogeny and systematics of the genus and a new type of canal raphe system. <i>Scientific Reports</i> , 2015, 5, 17115.	3.3	16
69	<p>Small-sized and discoid species of the genus Cocconeiospsis (Bacillariophyta) on Holothuria atra (Juan de Nova, Mozambique Channel)</p>. <i>Phytotaxa</i> , 2015, 54, 43.	0.3	11
70	Taxonomy, frustular morphology and systematics of <i>Platichthys</i> , a new genus of canal raphe bearing diatoms within the Entomoneidaceae. <i>Phytotaxa</i> , 2015, 236, 135.	0.3	6
71	<i>Pseudachnanthidium megapteropsis</i> gen. nov. and sp. nov. (Bacillariophyta): A Widespread Indo-Pacific Elusive Taxon. <i>Cryptogamie, Algologie</i> , 2015, 36, 291-304.	0.9	6
72	New species of <i>Eunotia</i> (Bacillariophyta) from Lake Baikal with comments on morphology and biogeography of the genus. <i>Phycologia</i> , 2015, 54, 248-260.	1.4	17

#	ARTICLE	IF	CITATIONS
73	New Insights into Plagiogrammaceae (Bacillariophyta) Based on Multigene Phylogenies and Morphological Characteristics with the Description of a New Genus and Three New Species. PLoS ONE, 2015, 10, e0139300.	2.5	29
74	Cocconeis Ehrenberg taxa (Bacillariophyta) with a marginal row of simple processes: relationship with the valvocopula system and distinctive features of related taxa. Fottea, 2015, 15, 139-154.	0.9	6
75	Diatoms from isolated islands II:Pseudostaurosira diablarum, a new species from a mangrove ecosystem in the GalÃ¡pagos Islands.. Diatom Research, 2014, 29, 201-211.	1.2	7
76	< i>Madinithidium gen. nov</i>. (Bacillariophyceae), a new monoraphid diatom genus from the tropical marine coastal zone. Phycologia, 2014, 53, 583-592.	1.4	16
77	Morphology, ecology and distribution of the diatom (Bacillariophyceae) species Simonsenia delognei (Grunow) Lange-Bertalot. Oceanological and Hydrobiological Studies, 2014, 43, 393-401.	0.7	7
78	Sea surface temperatures in Disko Bay during the Little Ice Age – caution needs to be exercised before assigning Thalassiosira kushirensis resting spore as a warm-water indicator in palaeoceanographic studies. Quaternary Science Reviews, 2014, 101, 234-237.	3.0	9
79	Minutocellus africana DÄ...bek & Witkowski sp. nov.: a new marine benthic diatom (Bacillariophyta,) Tj ETQql 1 0.784314 rgBT / Ov 223-232.	0.4	7
80	Looking forward through the past: identification of 50 priority research questions in palaeoecology. Journal of Ecology, 2014, 102, 256-267.	4.0	212
81	Fogedia giffeniana (Foged) Witkowski, Lange-Bertalot, Metzeltin & Bafana a benthic diatom new to the Turkish Aegean Sea. Su ÃœçerÃ¼nleri Dergisi, 2014, 31, 133-136.	0.3	0
82	A quantitative framework for analysis of regime shifts in a GalÃ¡pagos coastal lagoon. Ecology, 2014, 95, 3046-3055.	3.2	49
83	Description of diatoms from the Southwest to West Greenland coastal and open marine waters. Polar Biology, 2014, 37, 1589-1606.	1.2	23
84	A Description of Biremis panamae sp. nov., a New Diatom Species from the Marine Littoral, with an Account of the Phylogenetic Position of Biremis D.G. Mann et E.J. Cox (Bacillariophyceae). PLoS ONE, 2014, 9, e114508.	2.5	12
85	Diatom-based estimation of sea surface salinity in the south Baltic Sea and Kattegat. Baltica, 2014, 27, 131-140.	0.3	3
86	Late-Holocene diatom derived seasonal variability in hydrological conditions off Disko Bay, West Greenland. Quaternary Science Reviews, 2013, 67, 93-104.	3.0	21
87	An emended description of the genus Fogedia(Bacillariophyceae) with reports of four species new to science from a Korean sand flat. Phycologia, 2013, 52, 437-446.	1.4	6
88	An account of < i>Astartiella</i> species from tropical areas with a description of < i>A. societatis</i> sp. nov. and nomenclatural notes. Diatom Research, 2013, 28, 419-430.	1.2	8
89	<i>Gliwiczia gen. nov.</i> a new monoraphid diatom genus from Lake Baikal with a description of four species new for science. Phytotaxa, 2013, 109, 1.	0.3	53
90	Cymatosirella DÄ...bek, Witkowski & Sabbe gen. nov., a new marine benthic diatom genus (Bacillariophyta) belonging to the family Cymatosiraceae. Phytotaxa, 2013, 121, 42.	0.3	9

#	ARTICLE	IF	CITATIONS
91	Syvertsenia iberica (Cymatosiraceae): a new estuarine diatom genus characterized by the position of its process. <i>Phytotaxa</i> , 2013, 142, 25.	0.3	4
92	Description of a new marine diatom, <i>Coccconeis caulerpacola</i> sp. nov. (Bacillariophyceae), epiphytic on invasive Caulerpaspecies. <i>European Journal of Phycology</i> , 2012, 47, 433-448.	2.0	9
93	DESCRIPTION OF A NEW NAVICULOID DIATOM GENUS <i>< i>MORENEIS</i></i> GEN. NOV. (BACILLARIOPHYCEAE) FROM SAND FLATS IN KOREA ¹ . <i>Journal of Phycology</i> , 2012, 48, 186-195.	2.3	5
94	Reinterpretation of two diatom species from the West Greenland margin – Thalassiosira kushirensis and Thalassiosira antarctica var. borealis – hydrological consequences. <i>Marine Micropaleontology</i> , 2012, 88-89, 1-14.	1.2	14
95	Scalariella a new genus of monoraphid diatom (Bacillariophyta) with a bipolar distribution.. <i>Fottea</i> , 2012, 12, 13-25.	0.9	15
96	Valve ultrastructure of two new genera of marine canal-bearing diatoms (Bacillariophyceae). <i>Phycologia</i> , 2011, 50, 170-181.	1.4	12
97	Achnanthidium sibiricum (Bacillariophyceae), a new species from bottom sediments in Lake Baikal. <i>Algological Studies</i> (Stuttgart, Germany: 2007), 2011, 136-137, 77-87.	0.4	20
98	DIATOMS (BACILLARIOPHYTA) OF ISOLATED ISLANDS: NEW TAXA IN THE GENUS NAVICULA SENSU STRICTO FROM THE GALÃ¡PAGOS ISLANDS1. <i>Journal of Phycology</i> , 2011, 47, 861-879.	2.3	12
99	<i>< i>Planothidium iberense</i></i> sp. nov., a new brackish diatom of the Ebro Estuary, northeast Spain. <i>Diatom Research</i> , 2011, 26, 99-107.	1.2	5
100	Late Quaternary Climate Variations Reflected in Baltic Sea Sediments. <i>Central and Eastern European Development Studies</i> , 2011, , 99-132.	0.6	23
101	The genus Navicula in ancient basins. I. Two novelties from the Black Sea. <i>Plant Ecology and Evolution</i> , 2010, 143, 307-317.	0.7	12
102	Diatoms as a proxy in reconstructing the Holocene environmental changes in the south-western Baltic Sea: the lower Rega River Valley sedimentary record. <i>Hydrobiologia</i> , 2009, 631, 155-172.	2.0	24
103	Diatoms as a proxy in reconstructing the Holocene environmental changes in the south-western Baltic Sea: the lower Rega River Valley sedimentary record. , 2009, , 155-172.	0	
104	<i>< i>COCCONEIS GERMAINII</i></i> SP. NOV. AND A RELATED TAXON FROM KERGUELEN ARCHIPELAGO (AUSTRAL) Tj ETQq0 0 0 rgBT /Overlo	1.2	
105	Holocene marine diatoms from the Faeroe Islands and their paleoceanographic implications. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2006, 239, 487-509.	2.3	15
106	HOLOCENE DIATOMS (BACILLARIOPHYCEAE) FROM FAEROE ISLANDS FJORDS, NORTHERN ATLANTIC OCEAN. II. DISTRIBUTION AND TAXONOMY OF MARINE TAXA WITH SPECIAL REFERENCE TO BENTHIC FORMS. <i>Diatom Research</i> , 2006, 21, 175-215.	1.2	8
107	Palaeolimnology of Lake Zeribar, Iran, and its Climatic Implications. <i>Quaternary Research</i> , 2006, 66, 477-493.	1.7	58
108	Holocene North Atlantic surface circulation and climatic variability: evidence from diatom records. <i>Holocene</i> , 2005, 15, 85-96.	1.7	31

#	ARTICLE	IF	CITATIONS
109	Late Glacial and Holocene depositional history in the eastern part of the Szczecin Lagoon (Great) Tj ETQq1 1 0.784314 rgBT /Overlock 1.5 36		
110	Darss Sill as a biological border in the fossil record of the Baltic Sea: evidence from diatoms. Quaternary International, 2005, 130, 97-109.	1.5	41
111	HIPPONOTA SUBCOSTULATA(HUSTEDT) LANGE-BERTALOT, METZELTIN ET WITKOWSKI AND SOME FRAGILARIOID DIATOM TAXA FROM THE HOLOCENE LACUSTRINE SEDIMENTS OF THE FAEROE ISLANDS. Diatom Research, 2004, 19, 123-134.	1.2	1
112	EHRENBERGIULVAWITKOWSKI, LANGE-BERTALOT ET METZELTIN NOM. NOV."A NEW NAME FOREHRENBERGIAWITKOWSKIET AL.. Diatom Research, 2004, 19, 143-144.	1.2	1
113	Four new species of <i>Nitzschia</i> sect. <i>Tryblionella</i> (Bacillariophyceae) resembling <i>N. parvula</i> . Phycologia, 2004, 43, 579-595.	1.4	10
114	INFERRING SEA LEVEL VARIATION FROM RELATIVE PERCENTAGES OF <i>PSEUDOPODOSIRA KOSUGII</i> IN ROCHA LAGOON, SE URUGUAY. Diatom Research, 2003, 18, 49-59.	1.2	26
115	Diatom (Bacillariophyceae) flora of early Holocene freshwater sediments from Skalafjord, Faeroe Islands. Journal of Micropalaeontology, 2003, 22, 183-208.	3.6	13
116	A multi-proxy study of Pliocene sediments from Åžle de France, North-East Greenland. Palaeogeography, Palaeoclimatology, Palaeoecology, 2002, 186, 1-23.	2.3	49
117	Early Holocene history of the southwestern Baltic Sea: the Ancylus Lake stage. Boreas, 1999, 28, 437-453.	2.4	77
118	Early Holocene history of the southwestern Baltic Sea: the Ancylus Lake stage. Boreas, 1999, 28, 437-453.	2.4	15
119	The Baltic Ice Lake in the southwestern Baltic: sequenceâ€¢; chronoâ€¢and biostratigraphy. Boreas, 1997, 26, 217-236.	2.4	74
120	Fogedia gen. nov. (Bacillariophyceae), a new naviculoid genus from the marine littoral. Nova Hedwigia, 1997, 65, 79-98.	0.4	6
121	The Diatom Species <i>Fragilaria martyi</i> (Heribaud) Lange-Bertalot, Identity and Ecology. Archiv FÄ¼r Protistenkunde, 1996, 146, 281-292.	0.8	13
122	<i>Cocconeis hauniensis</i> sp. nov., a new epipsammic diatom from Puck Bay (Southern Baltic Sea), Poland. Nordic Journal of Botany, 1993, 13, 467-471.	0.5	15
123	Diatoms of the Puck Bay coastal shallows (Poland, Southern Baltic). Nordic Journal of Botany, 1991, 11, 689-701.	0.5	19
124	Epizoic diatoms on sea turtles and their relationship to host species, behaviour and biogeography: a morphological approach. European Journal of Phycology, 0, , 1-14.	2.0	4
125	Epilithic diatom communities from areas of invasive <i>Caulerpa</i> species (<i>Caulerpa taxifolia</i> and <i>Caulerpa</i>) Tj ETQq1 1 0.784314 rgBT /Overlock 1.6 2		
126	Chinia gen. nov."the second diatom genus simonsenoid raphe from mangroves in Fujian, China. Journal of Oceanology and Limnology, 0, , 1.	1.3	1

ARTICLE

IF CITATIONS

- | | | | |
|-----|--|-----|---|
| 127 | The sub-fossil diatom distribution in the Beibu Gulf (northwest South China Sea) and related environmental interpretation. PeerJ, 0, 10, e13115. | 2.0 | 3 |
|-----|--|-----|---|