

Erica Liberto

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/802010/publications.pdf>

Version: 2024-02-01

75
papers

3,029
citations

126708

33
h-index

174990

52
g-index

81
all docs

81
docs citations

81
times ranked

2621
citing authors

#	ARTICLE	IF	CITATIONS
1	Exploring food volatilome by advanced chromatographic fingerprinting based on comprehensive two-dimensional gas chromatographic patterns. <i>Comprehensive Analytical Chemistry</i> , 2022, 96, 261-261.	0.7	0
2	<i>Corylus avellana</i> L. Aroma Blueprint: Potent Odorants Signatures in the Volatilome of High Quality Hazelnuts. <i>Frontiers in Plant Science</i> , 2022, 13, 840028.	1.7	10
3	Exploring extra dimensions to capture saliva metabolite fingerprints from metabolically healthy and unhealthy obese patients by comprehensive two-dimensional gas chromatography featuring Tandem ionization mass spectrometry. <i>Analytical and Bioanalytical Chemistry</i> , 2021, 413, 403-418.	1.9	14
4	Exploring the Extra-Virgin Olive Oil Volatilome by Adding Extra Dimensions to Comprehensive Two-Dimensional Gas Chromatography and Time-of-Flight Mass Spectrometry Featuring Tandem Ionization: Validation of Ripening Markers in Headspace Linearity Conditions. <i>Journal of AOAC INTERNATIONAL</i> , 2021, 104, 274-287.	0.7	9
5	Cocoa smoky off-flavour: A MS-based analytical decision maker for routine controls. <i>Food Chemistry</i> , 2021, 336, 127691.	4.2	16
6	An effective chromatographic fingerprinting workflow based on comprehensive two-dimensional gas chromatography + Mass spectrometry to establish volatiles patterns discriminative of spoiled hazelnuts (<i>Corylus avellana</i> L.). <i>Food Chemistry</i> , 2021, 340, 128135.	4.2	23
7	Chromatographic Fingerprinting Strategy to Delineate Chemical Patterns Correlated to Coffee Odor and Taste Attributes. <i>Journal of Agricultural and Food Chemistry</i> , 2021, 69, 4550-4560.	2.4	18
8	Chromatographic Fingerprinting Enables Effective Discrimination and Identification of High-Quality Italian Extra-Virgin Olive Oils. <i>Journal of Agricultural and Food Chemistry</i> , 2021, 69, 8874-8889.	2.4	10
9	Adulteration of Essential Oils: A Multitask Issue for Quality Control. Three Case Studies: <i>Lavandula angustifolia</i> Mill., <i>Citrus limon</i> (L.) Osbeck and <i>Melaleuca alternifolia</i> (Maiden & Betche) Cheel. <i>Molecules</i> , 2021, 26, 5610.	1.7	19
10	Separation of stereoisomers by gas chromatography. , 2021, , 581-614.		4
11	Cocoa smoky off-flavor: Chemical characterization and objective evaluation for quality control. <i>Food Chemistry</i> , 2020, 309, 125561.	4.2	20
12	Sedentariness and Urinary Metabolite Profile in Type 2 Diabetic Patients, a Cross-Sectional Study. <i>Metabolites</i> , 2020, 10, 205.	1.3	7
13	Highly Informative Fingerprinting of Extra-Virgin Olive Oil Volatiles: The Role of High Concentration-Capacity Sampling in Combination with Comprehensive Two-Dimensional Gas Chromatography. <i>Separations</i> , 2019, 6, 34.	1.1	33
14	4th Conference on Cocoa Coffee and Tea (CoCoTea 2017) –“ The world in a cup. <i>Food Research International</i> , 2019, 115, 302.	2.9	1
15	Untargeted and Targeted Fingerprinting of Extra Virgin Olive Oil Volatiles by Comprehensive Two-Dimensional Gas Chromatography with Mass Spectrometry: Challenges in Long-Term Studies. <i>Journal of Agricultural and Food Chemistry</i> , 2019, 67, 5289-5302.	2.4	41
16	Robust Markers of Coffee Consumption Identified Among the Volatile Organic Compounds in Human Urine. <i>Molecular Nutrition and Food Research</i> , 2019, 63, e1801060.	1.5	16
17	Comprehensive two-dimensional gas chromatography coupled with time of flight mass spectrometry featuring tandem ionization: Challenges and opportunities for accurate fingerprinting studies. <i>Journal of Chromatography A</i> , 2019, 1597, 132-141.	1.8	30
18	Evaluation of volatile bioactive secondary metabolites transfer from medicinal and aromatic plants to herbal teas: Comparison of different methods for the determination of transfer rate and human intake. <i>Journal of Chromatography A</i> , 2019, 1594, 173-180.	1.8	14

#	ARTICLE	IF	CITATIONS
19	HS-SPME-MS-Enose Coupled with Chemometrics as an Analytical Decision Maker to Predict In-Cup Coffee Sensory Quality in Routine Controls: Possibilities and Limits. <i>Molecules</i> , 2019, 24, 4515.	1.7	11
20	Strategies for Accurate Quantitation of Volatiles from Foods and Plant-Origin Materials: A Challenging Task. <i>Journal of Agricultural and Food Chemistry</i> , 2019, 67, 1619-1630.	2.4	34
21	Ionic liquids as water-compatible GC stationary phases for the analysis of fragrances and essential oils. <i>Analytical and Bioanalytical Chemistry</i> , 2018, 410, 4657-4668.	1.9	24
22	Fructose liquid and solid formulations differently affect gut integrity, microbiota composition and related liver toxicity: a comparative in vivo study. <i>Journal of Nutritional Biochemistry</i> , 2018, 55, 185-199.	1.9	53
23	Evolution of potent odorants within the volatile metabolome of high-quality hazelnuts (<i>Corylus</i>) Tj ETQq1 1 0.784314 rgBT /Overlock spectrometry. <i>Analytical and Bioanalytical Chemistry</i> , 2018, 410, 3491-3506.	1.9	32
24	Combined untargeted and targeted fingerprinting by comprehensive two-dimensional gas chromatography: revealing fructose-induced changes in mice urinary metabolic signatures. <i>Analytical and Bioanalytical Chemistry</i> , 2018, 410, 2723-2737.	1.9	23
25	Advanced fingerprinting of high-quality cocoa: Challenges in transferring methods from thermal to differential-flow modulated comprehensive two dimensional gas chromatography. <i>Journal of Chromatography A</i> , 2018, 1536, 122-136.	1.8	35
26	Chemometric Modeling of Coffee Sensory Notes through Their Chemical Signatures: Potential and Limits in Defining an Analytical Tool for Quality Control. <i>Journal of Agricultural and Food Chemistry</i> , 2018, 66, 7096-7109.	2.4	40
27	Black tea volatiles fingerprinting by comprehensive two-dimensional gas chromatography " Mass spectrometry combined with high concentration capacity sample preparation techniques: Toward a fully automated sensomic assessment. <i>Food Chemistry</i> , 2017, 225, 276-287.	4.2	65
28	Fractionated dynamic headspace sampling in the analysis of matrices of vegetable origin in the food field. <i>Journal of Chromatography A</i> , 2017, 1489, 18-28.	1.8	11
29	Analysis of essential oils and fragrances with a new generation of highly inert gas chromatographic columns coated with ionic liquids. <i>Journal of Chromatography A</i> , 2017, 1495, 64-75.	1.8	29
30	Comprehensive Chemical Fingerprinting of High-Quality Cocoa at Early Stages of Processing: Effectiveness of Combined Untargeted and Targeted Approaches for Classification and Discrimination. <i>Journal of Agricultural and Food Chemistry</i> , 2017, 65, 6329-6341.	2.4	58
31	Coffee aroma: Chemometric comparison of the chemical information provided by three different samplings combined with GC-MS to describe the sensory properties in cup. <i>Food Chemistry</i> , 2017, 214, 218-226.	4.2	91
32	Enantioselective Gas Chromatography with Cyclodextrin in Odorant Analysis. , 2017, , 51-52.		3
33	Oxygenated heterocyclic compounds to differentiate Citrus spp. essential oils through metabolomic strategies. <i>Food Chemistry</i> , 2016, 206, 223-233.	4.2	25
34	Enantioselective Gas Chromatography with Derivatized Cyclodextrins in the Flavour and Fragrance Field. <i>Israel Journal of Chemistry</i> , 2016, 56, 925-939.	1.0	26
35	Chromatography: Focus on Multidimensional GC. , 2016, , 85-92.		0
36	Parallel dual secondary-column dual detection comprehensive two-dimensional gas chromatography: a flexible and reliable analytical tool for essential oils quantitative profiling. <i>Flavour and Fragrance Journal</i> , 2015, 30, 366-380.	1.2	29

#	ARTICLE	IF	CITATIONS
37	Cyclodextrin Derivatives as Stationary Phases for the GC Separation of Enantiomers in the Flavor and Fragrance Field. ACS Symposium Series, 2015, , 15-34.	0.5	6
38	Alignment for Comprehensive Two-Dimensional Gas Chromatography with Dual Secondary Columns and Detectors. Analytical Chemistry, 2015, 87, 10056-10063.	3.2	33
39	Herbs and spices: Characterization and quantitation of biologically-active markers for routine quality control by multiple headspace solid-phase microextraction combined with separative or non-separative analysis. Journal of Chromatography A, 2015, 1376, 9-17.	1.8	47
40	High-quality Italian rice cultivars: Chemical indices of ageing and aroma quality. Food Chemistry, 2015, 172, 305-313.	4.2	79
41	A metabolomic approach to quality determination and authentication of raw plant material in the fragrance field. Iris rhizomes: A case study. Journal of Chromatography A, 2014, 1368, 143-154.	1.8	22
42	Toward a definition of blueprint of virgin olive oil by comprehensive two-dimensional gas chromatography. Journal of Chromatography A, 2014, 1334, 101-111.	1.8	89
43	Urinary metabolic fingerprinting of mice with diet-induced metabolic derangements by parallel dual secondary column-dual detection two-dimensional comprehensive gas chromatography. Journal of Chromatography A, 2014, 1361, 265-276.	1.8	26
44	Parallel dual secondary column-dual detection: A further way of enhancing the informative potential of two-dimensional comprehensive gas chromatography. Journal of Chromatography A, 2014, 1360, 264-274.	1.8	30
45	Gas Chromatography in the Analysis of Flavours and Fragrances. , 2014, , 717-743.		1
46	Volatile profiling of high quality hazelnuts (Corylus avellana L.): Chemical indices of roasting. Food Chemistry, 2013, 138, 1723-1733.	4.2	53
47	Quantitative fingerprinting by headspaceâ€”Two-dimensional comprehensive gas chromatographyâ€”mass spectrometry of solid matrices: Some challenging aspects of the exhaustive assessment of food volatiles. Analytica Chimica Acta, 2013, 798, 115-125.	2.6	40
48	High concentration capacity sample preparation techniques to improve the informative potential of two-dimensional comprehensive gas chromatographyâ€”mass spectrometry: Application to sensomics. Journal of Chromatography A, 2013, 1318, 1-11.	1.8	29
49	Non-separative Headspace Solid Phase Microextractionâ€”Mass Spectrometry Profile as a Marker To Monitor Coffee Roasting Degree. Journal of Agricultural and Food Chemistry, 2013, 61, 1652-1660.	2.4	44
50	New medium-to-high polarity twister coatings for liquid and vapour phase sorptive extraction of matrices of vegetable origin. Journal of Chromatography A, 2012, 1265, 39-45.	1.8	36
51	Room temperature ionic liquids: New GC stationary phases with a novel selectivity for flavor and fragrance analyses. Journal of Chromatography A, 2012, 1268, 130-138.	1.8	43
52	A Further Tool To Monitor the Coffee Roasting Process: Aroma Composition and Chemical Indices. Journal of Agricultural and Food Chemistry, 2012, 60, 11283-11291.	2.4	46
53	Headspace Sampling in Flavor and Fragrance Field. , 2012, , 1-25.		8
54	Fast headspace-enantioselective GCâ€”mass spectrometric-multivariate statistical method for routine authentication of flavoured fruit foods. Food Chemistry, 2012, 132, 1071-1079.	4.2	56

#	ARTICLE	IF	CITATIONS
55	Quantitative analysis of volatiles from solid matrices of vegetable origin by high concentration capacity headspace techniques: Determination of furan in roasted coffee. <i>Journal of Chromatography A</i> , 2011, 1218, 753-762.	1.8	72
56	Profiling food volatiles by comprehensive two-dimensional gas chromatography coupled with mass spectrometry: Advanced fingerprinting approaches for comparative analysis of the volatile fraction of roasted hazelnuts (<i>Corylus avellana</i> L.) from different origins. <i>Journal of Chromatography A</i> , 2010, 1217, 5848-5858.	1.8	100
57	Essential oils and volatiles: sample preparation and analysis. A review.. <i>Flavour and Fragrance Journal</i> , 2010, 25, 282-290.	1.2	132
58	Solvent-enhanced headspace sorptive extraction in the analysis of the volatile fraction of matrices of vegetable origin. <i>Journal of Separation Science</i> , 2010, 33, 2191-2199.	1.3	16
59	New asymmetrical per-substituted cyclodextrins (2-O-methyl-3-O-ethyl- and) Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 50 587 Td (2-O-... chromatography in the flavour and fragrance field. <i>Journal of Chromatography A</i> , 2010, 1217, 1106-1113.	1.8	30
60	Quantitative determination of some volatile suspected allergens in cosmetic creams spread on skin by direct contact sorptive tape extraction-gas chromatography-mass spectrometry. <i>Journal of Chromatography A</i> , 2010, 1217, 2599-2605.	1.8	44
61	Development of fast enantioselective gas-chromatographic analysis using gas-chromatographic method-translation software in routine essential oil analysis (lavender essential oil). <i>Journal of Chromatography A</i> , 2010, 1217, 1530-1536.	1.8	40
62	Targeted and Non-Targeted Approaches for Complex Natural Sample Profiling by GCxGC-qMS. <i>Journal of Chromatographic Science</i> , 2010, 48, 251-261.	0.7	71
63	Fast-conventional quadrupole mass spectrometry in essential oil analysis. <i>Journal of Separation Science</i> , 2008, 31, 1074-1084.	1.3	34
64	Quantitative analysis of essential oils: a complex task. <i>Flavour and Fragrance Journal</i> , 2008, 23, 382-391.	1.2	163
65	Headspace sampling of the volatile fraction of vegetable matrices. <i>Journal of Chromatography A</i> , 2008, 1184, 220-233.	1.8	132
66	Enantiomer identification in the flavour and fragrance fields by interactive-combination of linear retention indices from enantioselective gas chromatography and mass spectrometry. <i>Journal of Chromatography A</i> , 2008, 1195, 117-126.	1.8	62
67	Conventional and narrow bore short capillary columns with cyclodextrin derivatives as chiral selectors to speed-up enantioselective gas chromatography and enantioselective gas chromatography-mass spectrometry analyses. <i>Journal of Chromatography A</i> , 2008, 1212, 114-123.	1.8	43
68	Reliability of fibres in solid-phase microextraction for routine analysis of the headspace of aromatic and medicinal plants. <i>Journal of Chromatography A</i> , 2007, 1152, 138-149.	1.8	57
69	Sorptive tape extraction in the analysis of the volatile fraction emitted from biological solid matrices. <i>Journal of Chromatography A</i> , 2007, 1148, 137-144.	1.8	55
70	Influence of polydimethylsiloxane outer coating and packing material on analyte recovery in dual-phase headspace sorptive extraction. <i>Journal of Chromatography A</i> , 2007, 1164, 33-39.	1.8	13
71	Headspace-Solid-Phase Microextraction in the Analysis of the Volatile Fraction of Aromatic and Medicinal Plants. <i>Journal of Chromatographic Science</i> , 2006, 44, 416-429.	0.7	73
72	Headspace-solid-phase microextraction fast GC in combination with principal component analysis as a tool to classify different chemotypes of chamomile flower-heads (<i>Matricaria recutita</i> L.). <i>Phytochemical Analysis</i> , 2006, 17, 217-225.	1.2	40

#	ARTICLE	IF	CITATIONS
73	Impact of phase ratio, polydimethylsiloxane volume and size, and sampling temperature and time on headspace sorptive extraction recovery of some volatile compounds in the essential oil field. <i>Journal of Chromatography A</i> , 2005, 1071, 111-118.	1.8	35
74	Dual-phase twistlers: A new approach to headspace sorptive extraction and stir bar sorptive extraction. <i>Journal of Chromatography A</i> , 2005, 1094, 9-16.	1.8	124
75	Automated headspace solid-phase dynamic extraction to analyse the volatile fraction of food matrices. <i>Journal of Chromatography A</i> , 2004, 1024, 217-226.	1.8	109