

Ignacio Romagosa

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/802186/publications.pdf>

Version: 2024-02-01

94
papers

3,597
citations

94433

37
h-index

161849

54
g-index

94
all docs

94
docs citations

94
times ranked

2819
citing authors

#	ARTICLE	IF	CITATIONS
1	Impact of Rising Temperature in the Deposition Patterns of Bioactive Compounds in Field Grown Food Barley Grains. <i>Plants</i> , 2021, 10, 598.	3.5	4
2	Bioactive Compounds and Antioxidant Capacity in Pearling Fractions of Hulled, Partially Hull-Less and Hull-Less Food Barley Genotypes. <i>Foods</i> , 2021, 10, 565.	4.3	7
3	Genetic and Management Effects on Barley Yield and Phenology in the Mediterranean Basin. <i>Frontiers in Plant Science</i> , 2021, 12, 655406.	3.6	12
4	Post-anthesis thermal stress induces differential accumulation of bioactive compounds in field-grown barley. <i>Journal of the Science of Food and Agriculture</i> , 2021, 101, 6496-6504.	3.5	1
5	Purple, high β -glucan, hulless barley as valuable ingredient for functional food. <i>LWT - Food Science and Technology</i> , 2020, 131, 109582.	5.2	26
6	CRISPR/Cas9 mutations in the rice <i>Waxy</i> / <i>GBSSI</i> gene induce allele-specific and zygosity-dependent feedback effects on endosperm starch biosynthesis. <i>Plant Cell Reports</i> , 2019, 38, 417-433.	5.6	45
7	The impact of climate change on barley yield in the Mediterranean basin. <i>European Journal of Agronomy</i> , 2019, 106, 1-11.	4.1	93
8	Genotype by Environment Interaction and Adaptation. , 2019, , 29-71.		5
9	ZmPBF and ZmGAMYB transcription factors independently transactivate the promoter of the maize (<i>Zea mays</i>) β -carotene hydroxylase 2 gene. <i>New Phytologist</i> , 2019, 222, 793-804.	7.3	20
10	Genotype by Environment Interaction and Adaptation. , 2018, , 1-44.		10
11	Effects of Barley (<i>Hordeum Vulgare</i> L.) Variety and Growing Environment on Beer Flavor. <i>Journal of the American Society of Brewing Chemists</i> , 2017, 75, 345-353.	1.1	30
12	Malt Modification and its Effects on the Contributions of Barley Genotype to Beer Flavor. <i>Journal of the American Society of Brewing Chemists</i> , 2017, 75, 354-362.	1.1	19
13	Biotecnología agrícola. <i>Arbor</i> , 2014, 190, a152.	0.3	0
14	QTLs for barley yield adaptation to Mediterranean environments in the "Nure"–"Tremois" biparental population. <i>Euphytica</i> , 2014, 197, 73-86.	1.2	74
15	Building bridges: an integrated strategy for sustainable food production throughout the value chain. <i>Molecular Breeding</i> , 2013, 32, 743-770.	2.1	28
16	Genome-wide association mapping of frost tolerance in barley (<i>Hordeum vulgare</i> L.). <i>BMC Genomics</i> , 2013, 14, 424.	2.8	101
17	Determinants of barley grain yield in drought-prone Mediterranean environments. <i>Italian Journal of Agronomy</i> , 2013, 8, 1.	1.0	17
18	Genetic control of duration of pre-anthesis phases in wheat (<i>Triticum aestivum</i> L.) and relationships to leaf appearance, tillering, and dry matter accumulation. <i>Journal of Experimental Botany</i> , 2012, 63, 69-89.	4.8	80

#	ARTICLE	IF	CITATIONS
19	Genetic characterization of a reciprocal translocation present in a widely grown barley variety. <i>Molecular Breeding</i> , 2012, 30, 1109-1119.	2.1	17
20	Agronomic effects of a reciprocal translocation in a widely grown Spanish barley variety. <i>Euphytica</i> , 2012, 185, 119-122.	1.2	0
21	Genetic control of pre-heading phases in the Steptoe \times Morex barley population under different conditions of photoperiod and temperature. <i>Euphytica</i> , 2012, 183, 303-321.	1.2	32
22	Determinants of barley grain yield in a wide range of Mediterranean environments. <i>Field Crops Research</i> , 2011, 120, 169-178.	5.1	73
23	Development of wild barley-derived DArT markers and their integration into a barley consensus map. <i>Molecular Breeding</i> , 2011, 27, 77-92.	2.1	32
24	Linkage map construction involving a reciprocal translocation. <i>Theoretical and Applied Genetics</i> , 2011, 122, 1029-1037.	3.6	35
25	Mixed model association scans of multi-environmental trial data reveal major loci controlling yield and yield related traits in <i>Hordeum vulgare</i> in Mediterranean environments. <i>Theoretical and Applied Genetics</i> , 2011, 122, 1363-1373.	3.6	75
26	Comparative mapping of the Oregon Wolfe Barley using doubled haploid lines derived from female and male gametes. <i>Theoretical and Applied Genetics</i> , 2011, 122, 1399-1410.	3.6	13
27	Genetic control of pre-heading phases and other traits related to development in a double-haploid barley (<i>Hordeum vulgare</i> L.) population. <i>Field Crops Research</i> , 2010, 119, 36-47.	5.1	51
28	Genetic relationships between preharvest sprouting and dormancy in barley. <i>Euphytica</i> , 2009, 168, 331-345.	1.2	31
29	Patterns of genetic diversity and linkage disequilibrium in a highly structured <i>Hordeum vulgare</i> association-mapping population for the Mediterranean basin. <i>Theoretical and Applied Genetics</i> , 2009, 119, 175-187.	3.6	99
30	Improvements in the production of doubled haploids in durum wheat (<i>Triticum turgidum</i> L.) through isolated microspore culture. <i>Plant Cell Reports</i> , 2009, 28, 727-735.	5.6	36
31	Genetic variability in duration of pre-heading phases and relationships with leaf appearance and tillering dynamics in a barley population. <i>Field Crops Research</i> , 2009, 113, 95-104.	5.1	68
32	Transgenic wheat plants expressing an oat arginine decarboxylase cDNA exhibit increases in polyamine content in vegetative tissue and seeds. <i>Molecular Breeding</i> , 2008, 22, 39-50.	2.1	21
33	Mapping adaptation of barley to droughted environments. <i>Euphytica</i> , 2008, 161, 35-45.	1.2	44
34	Changes in allele frequencies in landraces, old and modern barley cultivars of marker loci close to QTL for grain yield under high and low input conditions. <i>Euphytica</i> , 2008, 163, 435-447.	1.2	32
35	Barley adaptation and improvement in the Mediterranean basin. <i>Plant Breeding</i> , 2008, 127, 554-560.	1.9	40
36	Genetic markers for doubled haploid response in barley. <i>Euphytica</i> , 2007, 158, 287-294.	1.2	33

#	ARTICLE	IF	CITATIONS
37	Segregation distortion for agronomic traits in doubled haploid lines of barley. <i>Plant Breeding</i> , 2005, 124, 546-550.	1.9	10
38	Morphological and Agronomical Diversity Patterns in the Spanish Barley Core Collection. <i>Hereditas</i> , 2004, 135, 217-225.	1.4	33
39	Growth and yield responses of spring and winter triticale cultivated under Mediterranean conditions. <i>European Journal of Agronomy</i> , 2004, 20, 281-292.	4.1	27
40	Mixed models including environmental covariables for studying QTL by environment interaction. <i>Euphytica</i> , 2004, 137, 139-145.	1.2	128
41	Malting Behaviour of Barleys Grown in Canada and Spain as Related to Hordein and Enzyme Content. <i>Journal of the Institute of Brewing</i> , 2004, 110, 34-42.	2.3	15
42	Genetic control of dormancy in a Triumph/Morex cross in barley. <i>Theoretical and Applied Genetics</i> , 2004, 109, 62-70.	3.6	60
43	Use of new EST markers to elucidate the genetic differences in grain protein content between European and North American two-rowed malting barleys. <i>Theoretical and Applied Genetics</i> , 2004, 110, 116-125.	3.6	31
44	A centromeric region on chromosome 6(6H) affects dormancy in an induced mutant in barley. <i>Journal of Experimental Botany</i> , 2004, 56, 47-54.	4.8	11
45	Quantitative genetic analysis of acid detergent fibre content in barley grain. <i>Journal of Cereal Science</i> , 2003, 38, 167-172.	3.7	21
46	Productivity in prehistoric agriculture: physiological models for the quantification of cereal yields as an alternative to traditional approaches. <i>Journal of Archaeological Science</i> , 2003, 30, 681-693.	2.4	62
47	Patterns of grain filling of spring and winter hexaploid triticales. <i>European Journal of Agronomy</i> , 2002, 16, 219-230.	4.1	32
48	FOCUS: Estimated Wheat Yields During the Emergence of Agriculture Based on the Carbon Isotope Discrimination of Grains: Evidence from a 10th Millennium BP Site on the Euphrates. <i>Journal of Archaeological Science</i> , 2001, 28, 341-350.	2.4	41
49	Efficient production of androgenic doubled-haploid mutants in barley by the application of sodium azide to anther and microspore cultures. <i>Plant Cell Reports</i> , 2001, 20, 105-111.	5.6	50
50	Dormancy, ABA content and sensitivity of a barley mutant to ABA application during seed development and after ripening. <i>Journal of Experimental Botany</i> , 2001, 52, 1499-1506.	4.8	47
51	Low responsiveness of six-rowed genotypes to androgenesis in barley does not have a pleiotropic basis. <i>Genome</i> , 2001, 44, 936-940.	2.0	4
52	Mechanisms of Malt Extract Development in Barleys from Different European Regions: II. Effect of Barley Hordein Fractions on Malt Extract Yield. <i>Journal of the Institute of Brewing</i> , 2000, 106, 117-124.	2.3	30
53	Remobilization of Pre-Anthesis Assimilates to the Grain for Grain Only and Dual-Purpose (Forage and) Tj ETQq1 1 0.784314 rgBT / Ov	1.8	53
54	Individual Locus Effects on Dormancy during Seed Development and After Ripening in Barley. <i>Crop Science</i> , 1999, 39, 74-79.	1.8	44

#	ARTICLE	IF	CITATIONS
55	Crop water availability in early agriculture: evidence from carbon isotope discrimination of seeds from a tenth millennium BP site on the Euphrates. <i>Global Change Biology</i> , 1999, 5, 201-212.	9.5	81
56	RFLP markers associated with major genes controlling heading date evaluated in a barley germ plasm pool. <i>Heredity</i> , 1999, 83, 551-559.	2.6	22
57	Title is missing!. <i>Molecular Breeding</i> , 1999, 5, 143-152.	2.1	80
58	A mutant induced in the malting barley cv Triumph with reduced dormancy and ABA response. <i>Theoretical and Applied Genetics</i> , 1999, 98, 347-355.	3.6	27
59	Further evidence supporting Morocco as a centre of origin of barley. <i>Theoretical and Applied Genetics</i> , 1999, 98, 913-918.	3.6	49
60	Integrating statistical and ecophysiological analyses of genotype by environment interaction for grain filling of barley I.. <i>Field Crops Research</i> , 1999, 62, 63-74.	5.1	49
61	Integrating statistical and ecophysiological analyses of genotype by environment interaction for grain filling of barley II.. <i>Field Crops Research</i> , 1999, 62, 75-84.	5.1	39
62	Inheritance and fine mapping of a major barley seed dormancy QTL. <i>Plant Science</i> , 1999, 143, 113-118.	3.6	60
63	Genotype by environment interaction for grain yield and carbon isotope discrimination of barley in Mediterranean Spain. <i>Australian Journal of Agricultural Research</i> , 1999, 50, 1263.	1.5	102
64	The Spanish barley core collection. <i>Genetic Resources and Crop Evolution</i> , 1998, 45, 475-481.	1.6	61
65	Changes over time in the adaptation of barley releases in north-eastern Spain. <i>Plant Breeding</i> , 1998, 117, 531-535.	1.9	53
66	Mineral accumulation, carbon isotope discrimination and indirect selection for grain yield in two-rowed barley grown under semiarid conditions. <i>European Journal of Agronomy</i> , 1998, 9, 147-155.	4.1	57
67	Growth and Final Weight of Central and Lateral Barley Grains under Mediterranean Conditions as Influenced by Sink Strength. <i>Crop Science</i> , 1998, 38, 84-89.	1.8	23
68	The Development of Î ² -Glucanase and Degradation of Î ² -Glucan in Barley Grown in Scotland and Spain. <i>Journal of Cereal Science</i> , 1997, 26, 75-82.	3.7	34
69	Grain size and nitrogen accumulation in sink-reduced barley under Mediterranean conditions. <i>Field Crops Research</i> , 1997, 52, 117-126.	5.1	39
70	Molecular marker-assisted selection for malting quality traits in barley. <i>Molecular Breeding</i> , 1997, 3, 427-437.	2.1	96
71	Changes in carbon isotope discrimination in grain cereals from different regions of the western Mediterranean Basin during the past seven millennia. Palaeoenvironmental evidence of a differential change in aridity during the late Holocene. <i>Global Change Biology</i> , 1997, 3, 107-118.	9.5	100
72	Verification of barley seed dormancy loci via linked molecular markers. <i>Theoretical and Applied Genetics</i> , 1996, 92, 87-91.	3.6	72

#	ARTICLE	IF	CITATIONS
73	Use of the additive main effects and multiplicative interaction model in QTL mapping for adaptation in barley. <i>Theoretical and Applied Genetics</i> , 1996, 93-93, 30-37.	3.6	96
74	Effect of forage removal at the first detectable node stage on the growth of winter and spring triticale. <i>Grass and Forage Science</i> , 1996, 51, 170-179.	2.9	5
75	Agronomical and morphological differentiation among winter and spring triticales. <i>Plant Breeding</i> , 1995, 114, 413-416.	1.9	19
76	Growth and yield responses of two contrasting barley cultivars in a Mediterranean environment. <i>European Journal of Agronomy</i> , 1995, 4, 317-326.	4.1	24
77	Nitrogen fertilization of barley under semi-arid rainfed conditions. <i>European Journal of Agronomy</i> , 1995, 4, 309-316.	4.1	33
78	On the origin of Spanish two-rowed barleys. <i>Theoretical and Applied Genetics</i> , 1994, 87, 829-836.	3.6	22
79	Production of large number of doubled haploid plants from barley anthers pretreated with high concentrations of mannitol. <i>Plant Cell Reports</i> , 1994, 13, 709-12.	5.6	44
80	Integration of statistical and physiological analyses of adaptation of near-isogenic barley lines. <i>Theoretical and Applied Genetics</i> , 1993, 86, 822-826.	3.6	42
81	Differential Adaptation of Complete and Substituted Triticale. <i>Plant Breeding</i> , 1993, 111, 113-119.	1.9	28
82	Barley Tetrameric Inhibitor of Insect α -amylases. Characterization of an Allelic Variant of the BTAI-CMb Subunit. <i>Journal of Cereal Science</i> , 1993, 17, 107-113.	3.7	7
83	Genetic variants of the trypsin inhibitor from barley endosperm show different inhibitory activities. <i>Plant Science</i> , 1993, 89, 23-29.	3.6	17
84	Carbon Isotope Ratios in Ear Parts of Triticale. <i>Plant Physiology</i> , 1992, 100, 1033-1035.	4.8	32
85	Water Status Measurements of Excised Wheat Leaves: Position and Age Effects. <i>Crop Science</i> , 1991, 31, 1583-1588.	1.8	20
86	Screening Durum Wheat Germplasm for Dry Growing Conditions: Morphological and Physiological Criteria. <i>Crop Science</i> , 1991, 31, 770-775.	1.8	18
87	Quantitative phenotypical expression of three mutant genes in barley and the basis for defining an ideotype for Mediterranean environments. <i>Theoretical and Applied Genetics</i> , 1990, 80, 762-768.	3.6	9
88	Measurement and Use of Excised Leaf Water Status in Wheat. <i>Crop Science</i> , 1989, 29, 1140-1145.	1.8	38
89	RELATIONSHIP OF EXCISED-LEAF WATER LOSS RATE AND YIELD OF DURUM WHEAT IN DIVERSE ENVIRONMENTS. <i>Canadian Journal of Plant Science</i> , 1989, 69, 1075-1081.	0.9	43
90	Primary Trisomics in Sugarbeet. II. Cytological Identification 1. <i>Crop Science</i> , 1987, 27, 435-439.	1.8	15

#	ARTICLE	IF	CITATIONS
91	Primary Trisomics in Sugarbeet. I. Isolation and Morphological Characterization 1. Crop Science, 1986, 26, 243-249.	1.8	24
92	Cytological identification of acrotrisomy in sugar beets. Journal of Heredity, 1985, 76, 227-228.	2.4	6
93	Karyotype Analysis in Haploid Sugarbeet. Botanical Gazette, 1985, 146, 259-263.	0.6	8
94	Small Mesh-Bags within Industrial Malting Batches as a Simple Non-Expensive Alternative Micro-Malting Technique. Journal of the American Society of Brewing Chemists, 0, , 1-6.	1.1	0