

Carlos P Vio

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8067184/publications.pdf>

Version: 2024-02-01

82
papers

2,753
citations

172457

29
h-index

189892

50
g-index

83
all docs

83
docs citations

83
times ranked

3139
citing authors

#	ARTICLE	IF	CITATIONS
1	Renin-angiotensin system activation and interstitial inflammation in human diabetic nephropathy. <i>Kidney International</i> , 2003, 64, S64-S70.	5.2	154
2	Ischemic acute renal failure induces the expression of a wide range of nephrogenic proteins. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2006, 290, R861-R870.	1.8	150
3	Restoration of muscle strength in dystrophic muscle by angiotensin-1-7 through inhibition of TGF- β 2 signalling. <i>Human Molecular Genetics</i> , 2014, 23, 1237-1249.	2.9	143
4	Renal Identification of Cyclooxygenase-2 in a Subset of Thick Ascending Limb Cells. <i>Hypertension</i> , 1997, 30, 687-692.	2.7	121
5	Immunoreactive kallikrein localization in the rat kidney: an immuno-electron-microscopic study.. <i>Journal of Histochemistry and Cytochemistry</i> , 1984, 32, 117-121.	2.5	106
6	Hypokalemia induces renal injury and alterations in vasoactive mediators that favor salt sensitivity. <i>American Journal of Physiology - Renal Physiology</i> , 2001, 281, F620-F629.	2.7	97
7	Identification of persistently altered gene expression in the kidney after functional recovery from ischemic acute renal failure. <i>American Journal of Physiology - Renal Physiology</i> , 2005, 288, F953-F963.	2.7	86
8	Subcellular localization of renal kallikrein by ultrastructural immunocytochemistry. <i>Kidney International</i> , 1985, 28, 36-42.	5.2	79
9	Evidence for a stimulatory effect of high potassium diet on renal kallikrein. <i>Kidney International</i> , 1987, 31, 1327-1334.	5.2	79
10	Angiotensin II receptor type 1 blockade decreases CTGF/CCN2-mediated damage and fibrosis in normal and dystrophic skeletal muscles. <i>Journal of Cellular and Molecular Medicine</i> , 2012, 16, 752-764.	3.6	72
11	Mesenchymal stem cell injection ameliorates chronic renal failure in a rat model. <i>Clinical Science</i> , 2011, 121, 489-499.	4.3	67
12	Inhibition of the angiotensin-converting enzyme decreases skeletal muscle fibrosis in dystrophic mice by a diminution in the expression and activity of connective tissue growth factor (CTGF/CCN-2). <i>Cell and Tissue Research</i> , 2013, 353, 173-187.	2.9	67
13	bFGF induces an earlier expression of nephrogenic proteins after ischemic acute renal failure. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2006, 291, R1677-R1687.	1.8	66
14	Angiotensin II-independent Upregulation of Cyclooxygenase-2 by Activation of the (Pro)Renin Receptor in Rat Renal Inner Medullary Cells. <i>Hypertension</i> , 2013, 61, 443-449.	2.7	63
15	Human mesenchymal stem cells derived from adipose tissue reduce functional and tissue damage in a rat model of chronic renal failure. <i>Clinical Science</i> , 2013, 125, 199-210.	4.3	62
16	Fructose consumption reduces hippocampal synaptic plasticity underlying cognitive performance. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2015, 1852, 2379-2390.	3.8	55
17	Megalyn/LRP2 Expression Is Induced by Peroxisome Proliferator-Activated Receptor -Alpha and -Gamma: Implications for PPARs' Roles in Renal Function. <i>PLoS ONE</i> , 2011, 6, e16794.	2.5	54
18	Denervation-induced skeletal muscle fibrosis is mediated by CTGF/CCN2 independently of TGF- β 2. <i>Matrix Biology</i> , 2019, 82, 20-37.	3.6	52

#	ARTICLE	IF	CITATIONS
19	Aquaporin-2, a regulated water channel, is expressed in apical membranes of rat distal colon epithelium. <i>American Journal of Physiology - Renal Physiology</i> , 2001, 281, G856-G863.	3.4	50
20	Local induction of angiotensin-converting enzyme in the kidney as a mechanism of progressive renal diseases. <i>Kidney International</i> , 2003, 64, S57-S63.	5.2	50
21	Renal angiotensin II up-regulation and myofibroblast activation in human membranous nephropathy. <i>Kidney International</i> , 2003, 64, S39-S45.	5.2	50
22	Angiotensin II increases fibronectin and collagen I through the β -catenin-dependent signaling in mouse collecting duct cells. <i>American Journal of Physiology - Renal Physiology</i> , 2015, 308, F358-F365.	2.7	49
23	Role of Wnt Signaling in Tissue Fibrosis, Lessons from Skeletal Muscle and Kidney. <i>Current Molecular Medicine</i> , 2014, 14, 510-522.	1.3	47
24	Induction of Cyclooxygenase-2 in Thick Ascending Limb Cells by Adrenalectomy. <i>Journal of the American Society of Nephrology: JASN</i> , 2001, 12, 649-658.	6.1	41
25	Localization of Immunoreactive Glandular Kallikrein in Lactotrophs of the Rat Anterior Pituitary. <i>Neuroendocrinology</i> , 1990, 51, 10-14.	2.5	38
26	Sterol carrier protein 2 gene transfer changes lipid metabolism and enterohepatic sterol circulation in mice. <i>Gastroenterology</i> , 2000, 119, 1708-1719.	1.3	38
27	Inhibition of bFGF-receptor type 2 increases kidney damage and suppresses nephrogenic protein expression after ischemic acute renal failure. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2008, 294, R819-R828.	1.8	37
28	Localization of Immunoreactive Tissue Kallikrein in Human Trachea. <i>American Journal of Respiratory Cell and Molecular Biology</i> , 1993, 8, 16-19.	2.9	35
29	Renal and Vascular Injury Induced by Exogenous Angiotensin II Is AT1 Receptor-Dependent. <i>Nephron</i> , 2001, 87, 66-74.	1.8	34
30	Effect of ischemic acute renal damage on the expression of COX-2 and oxidative stress-related elements in rat kidney. <i>American Journal of Physiology - Renal Physiology</i> , 2007, 292, F1364-F1371.	2.7	28
31	E Prostanoid-1 receptor regulates renal medullary β -ENaC in rats infused with angiotensin II. <i>Biochemical and Biophysical Research Communications</i> , 2009, 389, 372-377.	2.1	28
32	Postnatal development of cyclooxygenase-2 in the rat kidney. <i>Immunopharmacology</i> , 1999, 44, 205-210.	2.0	27
33	Cyclooxygenase-2 induction by bradykinin in aortic vascular smooth muscle cells. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2006, 290, H30-H36.	3.2	26
34	The increased potassium intake improves cognitive performance and attenuates histopathological markers in a model of Alzheimer's disease. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2015, 1852, 2630-2644.	3.8	26
35	Bradykinin Stimulates Renal Na ⁺ and K ⁺ Excretion by Inhibiting the K ⁺ Channel (Kir4.1) in the Distal Convolute Tubule. <i>Hypertension</i> , 2018, 72, 361-369.	2.7	25
36	Mesangial cells are able to produce catecholamines in vitro. <i>Journal of Cellular Biochemistry</i> , 2003, 89, 144-151.	2.6	24

#	ARTICLE	IF	CITATIONS
37	Bradykinin Regulates Cyclooxygenase-2 in Rat Renal Thick Ascending Limb Cells. <i>Hypertension</i> , 2004, 44, 230-235.	2.7	23
38	Characterization of a long-term rat mTAL cell line. <i>American Journal of Physiology - Renal Physiology</i> , 2007, 293, F1413-F1422.	2.7	22
39	Pregnant Rats Treated With a Serotonin Precursor Have Reduced Fetal Weight and Lower Plasma Volume and Kallikrein Levels. <i>Hypertension</i> , 2007, 50, 773-779.	2.7	22
40	Induction of Renal Kallikrein and Renin Gene Expression by Insulin and IGF-I in the Diabetic Rat. <i>Diabetes</i> , 1997, 46, 2049-2056.	0.6	21
41	Basic fibroblast growth factor reduces functional and structural damage in chronic kidney disease. <i>American Journal of Physiology - Renal Physiology</i> , 2014, 306, F430-F441.	2.7	21
42	NFAT5 Is Protective Against Ischemic Acute Kidney Injury. <i>Hypertension</i> , 2014, 63, e46-52.	2.7	21
43	Renal phenotype of low kallikrein rats. <i>Kidney International</i> , 2001, 59, 2233-2242.	5.2	20
44	Sildenafil Stimulates and Dexamethasone Inhibits Pulmonary Vascular Development in Congenital Diaphragmatic Hernia Rat Lungs. <i>Neonatology</i> , 2014, 106, 74-80.	2.0	19
45	β -Catenin-Dependent Signaling Pathway Contributes to Renal Fibrosis in Hypertensive Rats. <i>BioMed Research International</i> , 2015, 2015, 1-13.	1.9	18
46	Kallikrein Excretion: Relationship with Maturation and Renal Function in Human Neonates at Different Gestational Ages. <i>Neonatology</i> , 1987, 52, 121-126.	2.0	17
47	Diabetes induces changes of catecholamines in primary mesangial cells. <i>International Journal of Biochemistry and Cell Biology</i> , 2008, 40, 747-754.	2.8	17
48	Renin-angiotensin system may trigger kidney damage in NOD mice. <i>JRAAS - Journal of the Renin-Angiotensin-Aldosterone System</i> , 2011, 12, 15-22.	1.7	17
49	Blockade of Bradykinin receptors worsens the dystrophic phenotype of mdx mice: differential effects for B1 and B2 receptors. <i>Journal of Cell Communication and Signaling</i> , 2018, 12, 589-601.	3.4	17
50	Cellular Distribution of Exogenous Aprotinin in the Rat Kidney. <i>Biological Chemistry</i> , 1998, 379, 1271-1278.	2.5	16
51	Long-Term, Fructose-Induced Metabolic Syndrome-Like Condition Is Associated with Higher Metabolism, Reduced Synaptic Plasticity and Cognitive Impairment in <i>Octodon degus</i> . <i>Molecular Neurobiology</i> , 2018, 55, 9169-9187.	4.0	16
52	Fetal Programming of Renal Dysfunction and High Blood Pressure by Chronodisruption. <i>Frontiers in Endocrinology</i> , 2019, 10, 362.	3.5	16
53	Postnatal maturation of tissue kallikrein-producing cells (connecting tubule cells) in the rat kidney: a morphometric and immunohistochemical study. <i>Anatomy and Embryology</i> , 1995, 192, 407-14.	1.5	15
54	Potassium Intake Prevents the Induction of the Renin-Angiotensin System and Increases Medullary ACE2 and COX-2 in the Kidneys of Angiotensin II-Dependent Hypertensive Rats. <i>Frontiers in Pharmacology</i> , 2019, 10, 1212.	3.5	14

#	ARTICLE	IF	CITATIONS
55	Canonical Wnt Signaling Modulates the Expression of Pre- and Postsynaptic Components in Different Temporal Patterns. <i>Molecular Neurobiology</i> , 2020, 57, 1389-1404.	4.0	14
56	TNF α regulates renal COX-2 in the rat thick ascending limb (TAL). <i>Thrombosis Research</i> , 2003, 110, 277-280.	1.7	13
57	Renal and Hormonal Effects of Water Deprivation in Late-Term Pregnant Rats. <i>Hypertension</i> , 2004, 44, 334-339.	2.7	13
58	Prostaglandin E ₂ EP3 receptor regulates cyclooxygenase-2 expression in the kidney. <i>American Journal of Physiology - Renal Physiology</i> , 2012, 303, F449-F457.	2.7	13
59	Eicosanoids and tumor necrosis factor-alpha in the kidney. <i>Prostaglandins and Other Lipid Mediators</i> , 2012, 98, 101-106.	1.9	13
60	Pregnant Rats With 5/6 Nephrectomy Have Normal Volume Expansion Despite Lower Renin and Kallikrein. <i>Hypertension</i> , 2003, 42, 744-748.	2.7	12
61	Catecholamine Production Along the Nephron. <i>Cellular Physiology and Biochemistry</i> , 2007, 20, 919-924.	1.6	12
62	Synthesis and expression of functional angiotensin II receptors in <i>Xenopus</i> oocytes injected with rat brain mRNA. <i>Molecular Brain Research</i> , 1987, 2, 268-270.	2.3	11
63	PGE ₂ EP ₃ receptor downregulates COX-2 expression in the medullary thick ascending limb induced by hypertonic NaCl. <i>American Journal of Physiology - Renal Physiology</i> , 2014, 307, F736-F746.	2.7	11
64	Arachidonic Acid Stimulates Renal Kallikrein Release in Isolated Rat Kidney. <i>Clinical Science</i> , 1982, 63, 235s-237s.	0.0	10
65	Cellular distribution and fate of the bradykinin antagonist HOE 140 in the rat kidney. Colocalization with the bradykinin B2 receptor. <i>Immunopharmacology</i> , 1996, 33, 146-150.	2.0	10
66	Early expression of monocyte chemoattractant protein-1 correlates with the onset of isoproterenol-induced cardiac fibrosis in rats with distinct angiotensin-converting enzyme polymorphism. <i>JRAAS - Journal of the Renin-Angiotensin-Aldosterone System</i> , 2008, 9, 154-162.	1.7	10
67	Renal Cyclooxygenase-2. <i>Hypertension</i> , 2001, 38, 630-634.	2.7	9
68	Cellular mechanisms of estrogen-and dopamine-induced control of glandular kallikrein in the anterior pituitary of the rat. <i>Cell and Tissue Research</i> , 1993, 274, 421-427.	2.9	7
69	P2C-Type ATPases and Their Regulation. <i>Molecular Neurobiology</i> , 2016, 53, 1343-1354.	4.0	7
70	Imbalance in Renal Vasoactive Enzymes Induced by Mild Hypoxia: Angiotensin-Converting Enzyme Increases While Neutral Endopeptidase Decreases. <i>Frontiers in Physiology</i> , 2018, 9, 1791.	2.8	6
71	Dietary Potassium Downregulates Angiotensin-I Converting Enzyme, Renin, and Angiotensin Converting Enzyme 2. <i>Frontiers in Pharmacology</i> , 2020, 11, 920.	3.5	6
72	N-domain angiotensin-I converting enzyme is expressed in immortalized mesangial, proximal tubule and collecting duct cells. <i>International Journal of Biological Macromolecules</i> , 2015, 72, 380-390.	7.5	4

#	ARTICLE	IF	CITATIONS
73	Renal phenotype of low kallikrein rats. <i>Kidney International</i> , 2001, 59, 2233.	5.2	4
74	Kallikrein–Kinin and Renin–Angiotensin Systems in Renovascular Hypertension in Rats. <i>Experimental Biology and Medicine</i> , 1980, 163, 447-451.	2.4	3
75	Immunocytochemical Identification of Glandular Kallikrein in the Rat Anterior Pituitary. <i>Advances in Experimental Medicine and Biology</i> , 1989, 247B, 183-188.	1.6	3
76	Cyclooxygenase-2 and hypoxia-regulated proteins are modulated by basic fibroblast growth factor in acute renal failure. <i>Biological Research</i> , 2012, 45, 51-60.	3.4	2
77	Effects of enriched-potassium diet on cardiorespiratory outcomes in experimental non-ischemic chronic heart failure. <i>Biological Research</i> , 2021, 54, 43.	3.4	2
78	Effect of COXâ€2 inhibition on sodium excretion and ENaC expression in Angiotensin II induced hypertensive rats. <i>FASEB Journal</i> , 2010, 24, 605.12.	0.5	0
79	Hormonal Regulation of Pituitary Glandular Kallikrein: A Morphometric Study. , 1992, 38 (Pt 1), 603-608.		0
80	Could Kinins Contribute to the Vasculoprotective Effect of Potassium Supplementation?. , 1995, , 80-89.		0
81	Dietary Potassium a Neglected Factor in Hypertension: Morphofunctional Evidence for Regulation of Cyclooxygenaseâ€2 and Kallikrein by Potassium. <i>FASEB Journal</i> , 2015, 29, 666.2.	0.5	0
82	Kir4.1 is involved in Bradykininâ€induced inhibition of NCC and natriuresis. <i>FASEB Journal</i> , 2018, 32, .	0.5	0