

# Dayan Knox

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8082507/publications.pdf>

Version: 2024-02-01

28  
papers

978  
citations

516710

16  
h-index

580821

25  
g-index

29  
all docs

29  
docs citations

29  
times ranked

1073  
citing authors

#	ARTICLE	IF	CITATIONS
1	Fear learning-induced changes in AMPAR and NMDAR expression in the fear circuit. <i>Learning and Memory</i> , 2022, 29, 83-92.	1.3	2
2	PI3K-Akt Signaling in the Basolateral Amygdala Facilitates Traumatic Stress Enhancements in Fear Memory. <i>International Journal of Neuropsychopharmacology</i> , 2021, 24, 229-238.	2.1	9
3	Issues with Using Animal Models to Examine Sex Difference in Developing PTSD. <i>Biological Psychiatry</i> , 2021, 89, S48-S49.	1.3	0
4	Systematic Review and Methodological Considerations for the Use of Single Prolonged Stress and Fear Extinction Retention in Rodents. <i>Frontiers in Behavioral Neuroscience</i> , 2021, 15, 652636.	2.0	17
5	The Effect of Traumatic Stress on the Mu-Opioid Receptors and Connectivity Within Reward Circuits. <i>Biological Psychiatry</i> , 2021, 89, S197.	1.3	0
6	Maternal Separation Induces Sex-Specific Differences in Sensitivity to Traumatic Stress. <i>Frontiers in Behavioral Neuroscience</i> , 2021, 15, 766505.	2.0	12
7	Single prolonged stress alters neural activation in the periaqueductal gray and midline thalamic nuclei during emotional learning and memory. <i>Learning and Memory</i> , 2019, 26, 403-411.	1.3	8
8	Using Near-infrared Fluorescence and High-resolution Scanning to Measure Protein Expression in the Rodent Brain. <i>Journal of Visualized Experiments</i> , 2019, , .	0.3	1
9	Using c-Jun to identify fear extinction learning-specific patterns of neural activity that are affected by single prolonged stress. <i>Behavioural Brain Research</i> , 2018, 341, 189-197.	2.2	18
10	Characterizing changes in glucocorticoid receptor internalization in the fear circuit in an animal model of post traumatic stress disorder. <i>PLoS ONE</i> , 2018, 13, e0205144.	2.5	10
11	Disruption of medial septum and diagonal bands of Broca cholinergic projections to the ventral hippocampus disrupt auditory fear memory. <i>Neurobiology of Learning and Memory</i> , 2018, 152, 71-79.	1.9	21
12	Neural circuits via which single prolonged stress exposure leads to fear extinction retention deficits. <i>Learning and Memory</i> , 2016, 23, 689-698.	1.3	46
13	The role of basal forebrain cholinergic neurons in fear and extinction memory. <i>Neurobiology of Learning and Memory</i> , 2016, 133, 39-52.	1.9	62
14	Cholinergic neuronal lesions in the medial septum and vertical limb of the diagonal bands of Broca induce contextual fear memory generalization and impair acquisition of fear extinction. <i>Hippocampus</i> , 2016, 26, 718-726.	1.9	61
15	Sex differences in the single prolonged stress model. <i>Behavioural Brain Research</i> , 2015, 286, 29-32.	2.2	60
16	Inhibiting corticosterone synthesis during fear memory formation exacerbates cued fear extinction memory deficits within the single prolonged stress model. <i>Behavioural Brain Research</i> , 2015, 287, 182-186.	2.2	46
17	Single prolonged stress enhances hippocampal glucocorticoid receptor and phosphorylated protein kinase B levels. <i>Neuroscience Research</i> , 2013, 75, 130-137.	1.9	65
18	Early handling attenuates enhancement of glucocorticoid receptors in the prefrontal cortex in an animal model of post-traumatic stress disorder. <i>Biology of Mood &amp; Anxiety Disorders</i> , 2013, 3, 22.	4.7	26

#	ARTICLE	IF	CITATIONS
19	Altered locus coeruleus norepinephrine function following single prolonged stress. <i>European Journal of Neuroscience</i> , 2013, 37, 901-909.	2.6	92
20	Expanding Our Understanding of Neurobiological Mechanisms of Resilience by Using Animal Models. <i>Neuropsychopharmacology</i> , 2012, 37, 317-318.	5.4	46
21	Single prolonged stress disrupts retention of extinguished fear in rats. <i>Learning and Memory</i> , 2012, 19, 43-49.	1.3	181
22	Unconditioned freezing is enhanced in an appetitive context: Implications for the contextual dependency of unconditioned fear. <i>Neurobiology of Learning and Memory</i> , 2012, 97, 386-392.	1.9	9
23	Inactivation of the prelimbic cortex enhances freezing induced by trimethylthiazoline, a component of fox feces. <i>Behavioural Brain Research</i> , 2011, 221, 320-323.	2.2	19
24	Single prolonged stress decreases glutamate, glutamine, and creatine concentrations in the rat medial prefrontal cortex. <i>Neuroscience Letters</i> , 2010, 480, 16-20.	2.1	111
25	Cortical modulation by nucleus basalis magnocellularis corticopetal cholinergic neurons during anxiety-like states is reflected by decreases in delta. <i>Brain Research</i> , 2008, 1227, 142-152.	2.2	5
26	Nucleus basalis magnocellularis and substantia innominata corticopetal cholinergic lesions attenuate freezing induced by predator odor. <i>Behavioral Neuroscience</i> , 2008, 122, 601-610.	1.2	4
27	Effect of nucleus basalis magnocellularis cholinergic lesions on fear-like and anxiety-like behavior. <i>Behavioral Neuroscience</i> , 2006, 120, 307-312.	1.2	29
28	Visceral Afferent Bias on Cortical Processing: Role of Adrenergic Afferents to the Basal Forebrain Cholinergic System. <i>Behavioral Neuroscience</i> , 2004, 118, 1455-1459.	1.2	18